



National Comprehensive
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Adult Cancer Pain

Version 2.2022 — June 27, 2022

[NCCN.org](https://www.nccn.org)

[Continue](#)



***Robert A. Swarm, MD/Chair** ϕ
Siteman Cancer Center at Barnes-
Jewish Hospital and Washington
University School of Medicine

***Jeanie M. Youngwerth, MD/Vice-Chair** £
University of Colorado Cancer Center

Julia L. Agne, MD £
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Doralina L. Anghelescu, MD £ ϕ
St. Jude Children's Research Hospital/
The University of Tennessee
Health Science Center

Madhuri Are, MD £
Fred & Pamela Buffett Cancer Center

Sorin Buga, MD £
City of Hope National Medical Center

Marcin Chwistek, MD £ P
Fox Chase Cancer Center

David Copenhaver, MD, MPH ϕ
UC Davis Comprehensive Cancer Center

***David Craig, PharmD** Σ £
Moffitt Cancer Center

Heidi Finnes, PharmD, BCOP Σ ‡
Mayo Clinic Cancer Center

Heather Greenlee, PhD, ND ¶
Fred Hutchinson Cancer Research Center/
Seattle Cancer Care Alliance

Eric Hansen, MD £
Roswell Park Comprehensive Cancer Center

Arif H. Kamal, MD, MBA, MHS †
Duke Cancer Institute

Mihir M. Kamdar, MD £ P
Massachusetts General Hospital Cancer Center

Enas Kandil, MD ϕ
UT Southwestern Simmons
Comprehensive Cancer Center

Sean Mackey, MD, PhD ϕ
Stanford Cancer Institute

Andrew McDonald, MD, MS §
O'Neal Comprehensive Cancer Center at UAB

M. Rachel McDowell, MSN, ACNP-BC † $\#$
Vanderbilt-Ingram Cancer Center

Kathryn McGrath, MD £
Abramson Cancer Center at the
University of Pennsylvania

Natalie Moryl, MD £ P
Memorial Sloan Kettering Cancer Center

Suzanne Nesbit, PharmD, BCPS Σ £
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Judith A. Paice, PhD, RN £ $\#$
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Peter Phung, MD, MBA £
UCLA Jonsson Comprehensive Cancer Center

Elizabeth Prsic, MD £ †
Yale Cancer Center/Smilow Cancer Hospital

Michael W. Rabow, MD £ P
UCSF Helen Diller Family
Comprehensive Cancer Center

Elizabeth Rickerson, MD £ ϕ
Dana-Farber/Brigham and Women's
Cancer Center

Rebecca Shatsky, MD †
UC San Diego Moores Cancer Center

Jill Sindt, MD ϕ
Huntsman Cancer Institute
at the University of Utah

Michael Smith, PharmD Σ £
University of Michigan
Rogel Cancer Center

Justine Yang Bruce, MD †
University of Wisconsin
Carbone Cancer Center

NCCN

Lisa A. Gurski, PhD
Beth McCullough, RN, BS

ϕ Anesthesiology	§ Radiotherapy/Radiation oncology
¶ Complementary and alternative medicine	£ Supportive care including palliative, pain management, pastoral care, & oncology social work
‡ Hematology/Hematology oncology	\ast Discussion section writing committee
P Internal medicine	
† Medical oncology	
$\#$ Nursing	
Σ Pharmacology	
θ Psychiatry, psychology, including health behavior	

Continue

[NCCN Guidelines Panel Disclosures](#)



[NCCN Adult Cancer Pain Panel Members](#)
[Summary of the Guidelines Updates](#)

[Principles of Cancer Pain Management \(PAIN-1\)](#)
[Universal Screening, Assessment, and Management of Pain \(PAIN-2\)](#)
[Management of Pain in Opioid-Naïve Patients \(PAIN-3\)](#)
[Management of Pain in Opioid-Tolerant Patients \(PAIN-4\)](#)
[Management of Pain Crisis \(PAIN-5\)](#)
[Ongoing Care \(PAIN-6\)](#)

[Pain Intensity Rating \(PAIN-A\)](#)
[Comprehensive Pain Assessment \(PAIN-B\)](#)
[Psychosocial Support \(PAIN-C\)](#)
[Integrative Interventions \(PAIN-D\)](#)
[Non-Opioid Analgesic \(Nonsteroidal Anti-inflammatory Drugs \[NSAIDs\] and Acetaminophen\) Prescribing \(PAIN-E\)](#)
[Adjuvant Analgesics for Neuropathic Pain \(PAIN-F\)](#)
[Opioid Principles, Prescribing, Titration, Maintenance, and Safety \(PAIN-G\)](#)
[Management of Opioid Adverse Effects \(PAIN-H\)](#)
[Patient and Family/Caregiver Education \(PAIN-I\)](#)
[Procedure-Related Pain and Anxiety \(PAIN-J\)](#)
[Management Strategies for Specific Cancer Pain Syndromes \(PAIN-K\)](#)
[Specialty Consultations for Improved Pain Management \(PAIN-L\)](#)
[Interventional Strategies \(PAIN-M\)](#)
[Principles of Pharmacogenetics \(PAIN-N\)](#)

Clinical Trials: NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

Find an NCCN Member Institution:
<https://www.nccn.org/home/member-institutions>.

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2022.



Updates in Version 2.2022 of the NCCN Guidelines for Adult Cancer Pain from Version 1.2022 include:

[PAIN-G](#)

- Header revised: Opioid Principles, Prescribing, *Initiation*, Titration, Maintenance, and Safety.
- Section updated with new buprenorphine content. (PAIN-G, 8 - 11)

[PAIN-G, 17 of 18](#)

- References updated.

[PAIN-G, 18 of 18](#)

- References updated.

Updates in Version 1.2022 of the NCCN Guidelines for Adult Cancer Pain from Version 2.2021 include:

[PAIN-1](#)

Principles of Cancer Pain Management

- Pain definition revised: Pain is defined by the International Association for the Study of Pain (IASP) as an unpleasant sensory and emotional experience associated with, or resembling that associated with, actual or potential tissue damage. *IASP defines chronic cancer-related pain as chronic pain caused by the primary cancer itself or metastases (chronic cancer pain) or its treatment (chronic post-cancer treatment pain).*
- New footnote b added: Bennett MI, Kaasa S, Barke A, et al. The IASP classification of chronic pain for ICD-11: Chronic cancer-related pain. Pain 2019;160:38-44.

[PAIN-2](#)

Universal Screening

- Following If pain present, second bullet revised: Severe uncontrolled pain is a medical emergency and should be addressed promptly (See *PAIN-5*).

[PAIN-3](#)

Pain Intensity

- General Principles: Link added to Principles of Cancer Pain Management (PAIN-1) (Also page PAIN-4)
 - ▶ First bullet following General Principles added: Optimize pain management therapies to improve function and meet patient's goals of care. (Also page PAIN-4)

[PAIN-6](#)

Ongoing Care

- Seventh bullet, fifth sub-bullet revised: Consider use of a pain diary to *facilitate* communication between patient and provider.

[PAIN-D](#)

• Cognitive modalities

- ▶ First bullet revised: *Mindfulness-based stress reduction (MBSR).*
- ▶ Tenth bullet revised: *Cognitive behavior therapy (CBT), cognitive restructuring.*

• Physical modalities

- ▶ Eighth bullet revised: Acupuncture, *electro-acupuncture*, or acupressure.

- New section added: Nutritional modalities

[Continued](#)

UPDATES



Updates in Version 1.2022 of the NCCN Guidelines for Adult Cancer Pain from Version 2.2021 include:

[PAIN-E, 2 of 2](#)

• GI Toxicities

- ▶ **First bullet revised:** Patients at high risk: age >60 years, history of peptic ulcer disease or significant alcohol use (3 or more alcoholic beverages/day), major organ dysfunction including hepatic dysfunction, highdose NSAIDs given for long periods, concomitant steroid use, and cardioprotective dose of daily aspirin, *and concomitant use of selective serotonin reuptake inhibitor (SSRI) antidepressants.*

[PAIN-G, 1 of 13](#)

- **General Principles, fourth bullet revised:** The appropriate opioid dose is the lowest dose that relieves the patient's pain and maximizes his or her function throughout the dosing interval without causing unmanageable adverse effects.

[PAIN-G, 2 of 13](#)

- **Sixth bullet, first sub-bullet revised:** If pain is inadequately controlled, to allow for dose titration, the short-acting opioids could be given as often as once an *per* hour as needed (if hourly dosing is needed for more than 3 cycles, reassessment or other intervention is recommended).

[PAIN-G, 4 of 13](#)

- **Opioids and Risk Evaluation and Mitigation Strategy (REMS), fourth bullet, fourth sub-bullet revised:** Patients with a history of psychiatric disorder, including anxiety, depression, *attention deficit hyperactivity disorder (ADHD), post-traumatic stress disorder (PTSD), bipolar disorder, or schizophrenia.*

[PAIN-G, 6 of 13](#)

- **Strategies to Maintain Patient Safety and Minimize the Risk of Opioid Misuse and Abuse During Chronic Opioid Use, links updated.**
- **Risk Mitigation for All patients Receiving Opioid Analgesics, first bullet, second sub-bullet revised:** Instruct ~~patients~~/caregivers to call emergency services (911) if naloxone is administered.

[PAIN-G, 8 of 13](#)

- **Mixed-Mechanism Drugs, third bullet revised:** Tramadol and tapentadol should be used with caution or avoided in patients taking other serotonergic or monoamine oxidase inhibitor (MAOI)-like medications (eg, tricyclic antidepressants [TCAs], ~~selective serotonin reuptake inhibitors~~ [SSRIs], SNRIs) due to risk of serotonin syndrome.

[PAIN-G, 10 of 13](#)

- **Special Notes Regarding Transdermal Fentanyl, sixth bullet revised:** When converting from continuous parenteral infusion fentanyl to transdermal fentanyl... *An alternative method is to stop the basal infusion 6 hours after patch placement.* In some patients, additional dose titration of the fentanyl patch may be necessary.
- **References updated.**

[Continued](#)

UPDATES



Updates in Version 1.2022 of the NCCN Guidelines for Adult Cancer Pain from Version 2.2021 include:

[PAIN-G, 12 of 13](#)

• Cautions Regarding Oral Methadone Use

▶ Third bullet, new sub-bullet added: 5–7 days is the time to steady state, and therefore, any significant sedation, significant improvement in pain OR immediate resolution of pain prior to the 5th day of methadone may indicate the dose is too high and the patient may be at risk of oversedation or respiratory depression by day 5–7 if the dose is not immediately adjusted.

▶ Fifth bullet revised: *Electrocardiogram (ECG)* should be considered prior to initiation and with each dose titration of methadone...

• Special Notes Regarding Oral Methadone Use, first bullet revised: *Methadone is commercially available in 5-mg and 10-mg tablets and 1 mg/mL, 2 mg/mL, and 10 mg/mL oral solution.*

[PAIN-H, 1 of 3](#)

• Constipation, first bullet, first sub-bullet revised: Educate patient and family on the need for bowel movements despite minimal *food intake of food.*

[PAIN-H, 3 of 3](#)

• Respiratory Depression, seventh bullet, second sub-bullet removed: Instruct patients/caregivers to call “911” if naloxone is administered.

[PAIN-K](#)

• New bullet added: Immunotherapy-related polyarthralgias (See NCCN Guidelines for Management of Immunotherapy-Related Toxicities).

[PAIN-L](#)

• Mental health consultation, fourth bullet, sub-bullet revised: Integrative medicine practitioners can be used to deliver evidence-based treatment modalities (eg, *cognitive behavioral therapy [CBT], mindfulness-based stress reduction [MBSR], acceptance-based therapy, biofeedback, hypnosis*).

**PRINCIPLES OF CANCER PAIN MANAGEMENT****Pain Definition**

Pain is defined by the International Association for the Study of Pain (IASP) as an unpleasant sensory and emotional experience associated with, or resembling that associated with, actual or potential tissue damage.^a IASP defines chronic cancer-related pain as chronic pain caused by the primary cancer itself, or metastases (chronic cancer pain) or its treatment (chronic post-cancer treatment pain).^b

General Principles

- Optimal use of disease-specific therapies is essential to managing tumor-related pain.
- Survival is linked to symptom control and pain management, which contribute to broad quality-of-life improvement. Pain management is an essential part of oncologic management.
- Analgesic therapy is done in conjunction with management of multiple symptoms or symptom clusters. Consider the interaction of complex pharmacologic therapies and the risk for analgesic misuse.
- An interdisciplinary team is optimal; consider early referral to a palliative care provider. [\(See NCCN Guidelines for Palliative Care\)](#)
- Provide/refer for psychosocial support, including emotional and informational support and coping skills training. [\(See PAIN-C\)](#)
- Provide accessible educational material to improve pain assessment, pain management, and the safe use of opioid medications based on the patient's identified needs.^b [\(See PAIN-I\)](#)
 - ▶ Involve patients in developing treatment plans and setting meaningful, realistic expectations and measurable goals.
- Address the multidimensional impact of “suffering” on patients and caregivers in a culturally respectful manner.

Assessment

- Screen all patients for pain at each contact. [\(See PAIN-2\)](#)
- Routinely quantify and document pain intensity and quality as characterized by the patient (whenever possible). Include patient reporting of breakthrough pain, treatments used and their impact on pain, satisfaction with pain relief, pain interference, provider assessment of impact on function, and any special issues for the patient relevant to pain treatment. If necessary, get additional

^a Raja SN, Carr DB, Cohen M, et al. The revised International Association for the Study of Pain definition of pain: concepts, challenges, and compromises Pain 2020;161:1976-1982.

- information from caregiver regarding pain and impact on function.
- Perform comprehensive pain assessment if new or worsening pain is present and regularly for persisting pain. [\(See PAIN-B\)](#)
- Evaluate for risk factors for opioid abuse/misuse/diversion.

Management/Intervention

- Goals of pain management are highlighted by the “5 A’s” of outcomes:^c
 1. Analgesia (optimize analgesia)
 2. Activities (optimize activities of daily living [ADLs])
 3. Adverse effects (minimize adverse effects) [\(See PAIN-G\)](#)
 4. Aberrant drug taking (avoid aberrant drug taking) [\(See PAIN-H\)](#)
 5. Affect (relationship between pain and mood)
- Prevention of analgesic side effects, especially constipation, is of paramount importance.
- For acute, severe pain or pain crisis, consider hospital or inpatient hospice admission.
- Treat persistent cancer pain with regularly scheduled analgesics or long-acting analgesics, and breakthrough pain with supplemental doses of short-acting analgesics.
- For chronic pain in cancer survivors, [See NCCN Guidelines for Survivorship](#).

Reassessment

- Perform pain reassessment at specified intervals to ensure that analgesic therapy is providing maximum benefit with minimal adverse effects, and that the treatment plan is followed.
- Encourage patients to report ongoing pain assessments in between visits, as needed.

^b Bennett MI, Kaasa S, Barke A, et al. The IASP classification of chronic pain for ICD-11: chronic cancer-related pain. Pain 2019;160:38-44.

^c The Joint Commission. New and Revised Pain Assessment and Management Standards. 2018. <https://www.jointcommission.org/resources/patient-safety-topics/pain-management-standards-for-accredited-organizations/>

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.


[See Universal Screening \(PAIN-2\)](#)



UNIVERSAL SCREENING

ASSESSMENT

MANAGEMENT OF PAIN



^c For chronic pain in cancer survivors, see the [NCCN Guidelines for Survivorship](#).

^d Opioid-naïve patients are those not chronically receiving opioid analgesic on a daily basis and therefore have not developed significant tolerance. Opioid tolerant includes patients who are chronically receiving opioid analgesic on a daily basis. The FDA identifies tolerance as receiving at least 25 mcg/h fentanyl patch, at least 60 mg of morphine daily, at least 30 mg of oral oxycodone daily, at least 8 mg of oral hydromorphone daily, or an equianalgesic dose of another opioid for a week or longer.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PAIN INTENSITY MANAGEMENT OF PAIN IN OPIOID-NAÏVE PATIENTS^d

[See Pain Intensity Rating \(PAIN-A\)](#)

General Principles
See [Principles of Cancer Pain Management \(PAIN-1\)](#)

- Optimize pain management therapies to improve function and meet patient's goals of care
- Select the most appropriate medication based on the pain diagnosis ([See PAIN-K](#)), comorbid conditions, and potential drug interactions
- Analgesic regimen may include an opioid ([See PAIN-G](#)), acetaminophen, nonsteroidal anti-inflammatory drugs (NSAIDs) ([See PAIN-E](#)), and/or adjuvant analgesics ([See PAIN-F](#))
 - ▶ If pain is continuous, consider regularly scheduled analgesics
- Anticipate and treat analgesic adverse effects, including opioid-induced constipation ([See PAIN-H](#))
- Provide psychosocial support ([See PAIN-C](#))
- Provide patient and family/caregiver education ([See PAIN-I](#))
- Optimize integrative interventions ([See PAIN-D](#))

Reevaluate pain at each contact and as needed to meet patient-specific goals for comfort, function, and safety

[See Ongoing Care \(PAIN-6\)](#)

Mild Pain

- See General Principles above AND
- First consider non-opioids and adjuvant therapies, unless these are contraindicated due to adverse effects, potential drug interactions, or comorbid conditions ([See PAIN-G](#))

Moderate/
Severe Pain

- See General Principles above AND
- Non-opioids and adjuvant therapies as appropriate with short-acting opioids as needed ([See PAIN-G](#))
- Start and titrate short-acting opioid, every 3–4 hours as needed^{e,f} ([See PAIN-G, 7 of 13](#))
 - ▶ Oxycodone immediate release (IR) 2.5–5 mg with or without acetaminophen 325 mg
 - ▶ Hydrocodone 5 mg with acetaminophen 325 mg
 - ▶ Hydromorphone 2 mg PO
 - ▶ Morphine 5 mg (solution) or IR 7.5 mg (1/2 tablet)
- If 4 or more doses of short-acting opioid are consistently needed per day, consider addition of a long-acting opioid based on the total daily dose

- Titrate further as needed
- If pain is stable, see Ongoing Care ([PAIN-6](#))
- If pain is inadequately controlled, reevaluate working diagnosis with a comprehensive pain assessment ([See PAIN-B](#))
- Consider pain specialty and/or palliative care consultation ([See PAIN-L](#))
- Consider opioid rotation if dose-limiting adverse effects are noted

Severe Pain/
Pain Crisis

For acute, severe pain or pain crisis, consider hospital or inpatient hospice admission to achieve patient-specific goals for comfort and function ([See PAIN-5](#))

[See Footnotes on PAIN-3A](#)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



FOOTNOTES

- ^d Opioid-naïve patients are those not chronically receiving opioid analgesic on a daily basis and therefore have not developed significant tolerance. Opioid tolerant includes patients who are chronically receiving opioid analgesic on a daily basis. The FDA identifies tolerance as receiving at least 25 mcg/h fentanyl patch, at least 60 mg of morphine daily, at least 30 mg of oral oxycodone daily, at least 8 mg of oral hydromorphone daily, or an equianalgesic dose of another opioid for a week or longer.
- ^e Select, extended-release opioids may also be indicated for opioid-naïve patients in rare circumstances.
- ^f More frequent dosing may be indicated in select situations.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PAIN INTENSITY MANAGEMENT OF PAIN IN OPIOID-TOLERANT PATIENTS^d

[See Pain Intensity Rating \(PAIN-A\)](#)

General Principles
[See Principles of Cancer Pain Management \(PAIN-1\)](#)

- Optimize pain management therapies to improve function and meet patient's goals of care
- Select the most appropriate medication based on the pain diagnosis ([See PAIN-K](#)), comorbid conditions, and potential drug interactions
- Analgesic regimen may include an opioid ([See PAIN-G](#)), acetaminophen, NSAIDs ([See PAIN-E](#)), and/or adjuvant analgesics ([See PAIN-F](#))
 - ▶ If pain is continuous, consider regularly scheduled analgesics
- Anticipate and treat analgesic adverse effects, including opioid-induced constipation ([See PAIN-H](#))
- Provide psychosocial support ([See PAIN-C](#))
- Provide patient and family/caregiver education ([See PAIN-I](#))
- Optimize integrative interventions ([See PAIN-D](#))

Reevaluate pain at each contact and as needed to meet patient-specific goals for comfort, function, and safety

[See Ongoing Care \(PAIN-6\)](#)

Mild Pain →

- See General Principles above AND
- Non-opioids and adjuvant therapies, unless these are contraindicated due to adverse effects or potential drug interactions ([See PAIN-G](#))
- Re-evaluate need for opioids and reduce if appropriate ([See PAIN-G, 3 of 13](#))

Moderate/ Severe Pain →

- See General Principles above AND
- Non-opioids and adjuvant therapies as appropriate with short-acting opioids as needed ([See PAIN-G](#))
- Titrate short-acting opioid (may require dose increase of 30%–50% or more) ([See PAIN-5](#)) →
- If 4 or more doses of short-acting opioid are consistently needed per day, consider addition or increase in dose of a long-acting opioid based on the total daily dose

- Titrate further as needed
- If pain is stable, see Ongoing Care ([PAIN-6](#))
- If pain is inadequately controlled, reevaluate working diagnosis with a comprehensive pain assessment ([See PAIN-B](#))
- Consider pain specialty and/or palliative care consultation ([See PAIN-L](#))
- Consider opioid rotation if dose-limiting adverse effects are noted

Severe Pain/ Pain Crisis →

For acute, severe pain or pain crisis, consider hospital or inpatient hospice admission to achieve patient-specific goals for comfort and function ([See PAIN-5](#))

^d Opioid-naïve patients are those not chronically receiving opioid analgesic on a daily basis and therefore have not developed significant tolerance. Opioid tolerant includes patients who are chronically receiving opioid analgesic on a daily basis. The FDA identifies tolerance as receiving at least 25 mcg/h fentanyl patch, at least 60 mg of morphine daily, at least 30 mg of oral oxycodone daily, at least 8 mg of oral hydromorphone daily, or an equianalgesic dose of another opioid for a week or longer.


Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

MANAGEMENT OF PAIN CRISIS

Monitor for acute and chronic adverse effects. [See Management of Opioid Adverse Effects \(PAIN-H\)](#)

INITIAL DOSE [\(See PAIN-G\)](#)⁹

SUBSEQUENT DOSE⁹



^d Opioid-naïve patients are those not chronically receiving opioid analgesic on a daily basis and therefore have not developed significant tolerance. Opioid tolerant includes patients who are chronically receiving opioid analgesic on a daily basis. The FDA identifies tolerance as receiving at least 25 mcg/h fentanyl patch, at least 60 mg of morphine daily, at least 30 mg of oral oxycodone daily, at least 8 mg of oral hydromorphone daily, or an equianalgesic dose of another opioid for a week or longer.

⁹ Dose and titrate with caution in patients with risk factors such as decreased renal/hepatic function, chronic lung disease, upper airway compromise, sleep apnea, and poor performance status.

^h Not including transmucosal fentanyl dose.


Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



ONGOING CARE

- If applicable, convert from parenteral to oral/transdermal opioids (if feasible) including extended-release or long-acting agent with rescue doses (Conversion details, [See PAIN-G](#))
 - ▶ Simplify analgesic regimen for improved patient compliance, if feasible.
- Have regular follow-up schedule to monitor pain therapy outcomes
 - ▶ Assess pain during each outpatient contact or at least each day for inpatients or more frequently based on:
 - ◊ Patient's condition, including analgesic therapy adverse effects
 - ◊ Institutional standards
 - ◊ Regulatory requirements
- Monitor for the use of analgesics as prescribed, especially in patients with risk factors for or history of substance abuse/diversion or cognitive dysfunction
- Provide written follow-up pain plan, including prescribed medications ([See PAIN-I](#))
- Ensure continuity of care during transition between sites of care
 - ▶ Collaborate with patient's pharmacist and insurance company if needed
 - ▶ Clarify which clinician will be prescribing patient's ongoing analgesics
- Address system barriers, and recruit assistance from social services as needed
 - ▶ Analgesic cost/pharmacy benefit coverage
 - ▶ Availability of analgesics
 - ▶ Local laws/regulations
- Instruct the patient on the importance of: ([See PAIN-I](#))
 - ▶ Following documented pain plan
 - ▶ Scheduling and keeping outpatient appointments
 - ▶ Contacting clinician if pain worsens or adverse effects are inadequately controlled, including availability of after-hours assistance to facilitate titration of analgesic
 - ▶ Safe handling, storage, and disposal of analgesics
 - ▶ Consider use of a pain diary to facilitate communication between patient and provider
- Reevaluate patient-centered goals of care in the context of current disease and available therapies
- Maintain communication and consider referral to pain/palliative care specialist and relevant providers, especially during transition between sites of care. ([See NCCN Guidelines for Palliative Care](#))

GOALS OF TREATMENT



Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PAIN INTENSITY RATING

- Pain intensity rating scales can be used as part of universal screening and comprehensive pain assessment. At minimum, patients should be asked about “current” pain, as well as “worst” pain, “average” pain, and “least” pain in the past 24 hours. For each pain intensity rating, use one of the scales below.
- For comprehensive assessment, also include “worst pain in past week,” “pain at rest,” and “pain with movement.” [See Comprehensive Pain Assessment \(PAIN-B\)](#) for more details.

Table 1: Numerical Rating Scale

• Verbal: “What number describes your pain from 0 (no pain) to 10 (worst pain you can imagine)?”
• Written: “Circle the number that describes your pain.”

0	1	2	3	4	5	6	7	8	9	10
No pain					Worst pain you can imagine					

Categorical scale: “What word best describes your pain?”

None (0)	Mild (1–3)	Moderate (4–6)	Severe (7–10)
----------	------------	----------------	---------------

Table 2: The Faces Pain Rating Scale - Revised^{1,2}

Instructions:
“These faces show how much something can hurt. This face (point to the left-most face) shows no pain. Each face shows more and more pain (point to each face from left to right) up to this one (point to the right-most face)—it shows very much pain. Point to the face that shows how much you hurt (right now).”

¹ Hicks CL, von Baeyer CL, Spafford P, et al. The Faces Pain Scale - Revised: Toward a common metric in pediatric pain measurement. *Pain* 2001;93:173-183.

² Ware LJ, Epps CD, Herr K, Packard A. Evaluation of the Revised Faces Pain Scale, Verbal Descriptor Scale, Numeric Rating Scale, and Iowa Pain Thermometer in older minority adults. *Pain Manag Nurs* 2006;7:117-125.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PAIN INTENSITY RATING****Pain Assessment in the Nonverbal Patient**³

- The inability of patients to verbally communicate pain intensity because of cognitive or physiologic issues is a major barrier relating to pain assessment and management. Therefore, the American Society for Pain Management Nursing (www.aspmn.org) has developed a position statement and clinical practice recommendations clinicians may find useful in caring for such patients.
- In the absence of self-report, observation of behavior is a valid approach to pain assessment with the understanding that behaviors may also indicate other sources of distress, such as emotional stress or delirium, which may complicate assessment ([See NCCN Guidelines for Distress Management](#)). Potential causes and the context of the behavior must be considered when making pain treatment decisions.
- A multi-faceted approach is recommended that combines direct observation, family/caregiver input, and evaluation of response to pain medicines or nonpharmacologic interventions.
- For patients with advanced dementia, a comprehensive review of currently published tools, including those available at https://prc.coh.org/pain_assessment_new.asp, is recommended.
- For patients who are intubated and/or are unconscious, pain assessment tools have been tested in specific situations and include, but are not limited to:
 - ▶ Behavioral Pain Scale (BPS) tested in adults in intensive care: <http://www.ncbi.nlm.nih.gov/pubmed/11801819>⁴
 - ▶ Critical-Care Pain Observation Tool (CPOT) tested in adults in intensive care: <http://www.ncbi.nlm.nih.gov/pubmed/17575489>⁵
- Clinicians are encouraged to monitor current research regarding new developments in strategies and tools for assessing pain in patients who have difficulty with self-reporting.

Cultural and Linguistic Assessment^{6,7}

- Health care providers should be aware of impact of cultural and linguistic diversity during universal screening and comprehensive pain assessment and respond with trained interpreters and culturally and linguistically appropriate educational materials.

³ Herr K, Coyne P, Key T, et al. Pain assessment in the nonverbal patient: Position statement with clinical practice recommendations. Pain Manag Nurs 2006;7:44-52.

⁴ Payen JF, Bru O, Bosson JL, et al. Assessing pain in critically ill sedated patients by using a behavioral pain scale. Crit Care Med 2001;29:2258-2263.

⁵ Gélinas C, Johnston C. Pain assessment in the critically ill ventilated adult: validation of the Critical-Care Pain Observation Tool and physiologic indicators. Clin J Pain 2007;23:497-505.

⁶ Al-Atiyat HNM. Cultural diversity and cancer pain. J Hosp Palliat Nurs 2009;11:154-164.

⁷ Ezenwa MO, Ameringer S, Ward SE, Serlin RC. Racial and ethnic disparities in pain management in the United States. J Nurs Scholarsh 2006;38:225-233.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**COMPREHENSIVE PAIN ASSESSMENT**

- Patient's self report of pain is the standard of care. If the patient is unable to verbally report pain, an alternative method to obtain pain rating and response should be utilized. ([See PAIN-A, 2 of 2](#)).
- The goal of comprehensive pain assessment is to find the cause of the pain and identify optimal therapies. Individualized pain treatment is based on the etiology and characteristics of pain, the patient's clinical condition, and patient-centered goals of care.
- The etiology and pathophysiology of the pain should be investigated, including medical history (including psychosocial factors), physical exam, laboratory tests, and imaging studies.
 - ▶ Etiology factors may include direct involvement of cancer itself, cancer therapy (chemotherapy, RT, surgery) or procedures, and coincidental or acute or chronic noncancer pain (eg, arthritis).
 - ▶ Pathophysiology factors may include nociceptive, neuropathic, visceral, affective, behavioral, and cognitive components.
- **Pain experience**
 - ▶ Location, referral pattern, radiation of pain(s)
 - ▶ Intensity [See Pain Intensity Rating \(PAIN-A\)](#)
 - ◇ Last 24 hours worst and least pain and pain now
 - ◇ At rest and with movement
 - ▶ Interference with activities
[See Impact of Pain Measurement \(PAIN-B, 3 of 3\)](#)
 - ◇ General activity, mood, walking ability, work ability, relationship with others, sleep, appetite, and enjoyment of life
 - ▶ Timing: onset, duration, course, persistent, or intermittent
 - ▶ Description or quality
 - ◇ Aching, stabbing, throbbing, or pressure often associated with somatic pain in skin, muscle, and bone
 - ◇ Gnawing, cramping, aching, or sharp pain often associated with visceral pain in organs or viscera
 - ◇ Burning, tingling, shooting, or electric/shocking pain often associated with neuropathic pain caused by nerve damage
 - ▶ Aggravating and alleviating factors
 - ▶ Other current symptoms; symptom clusters
 - ▶ Current pain management plan, both pharmacologic and non-pharmacologic. If medications are used, determine:
 - ◇ What medication(s), prescription and/or over the counter (OTC)?
 - ◇ Dose, route of administration, frequency?
 - ◇ Current prescriber?
- **Pain experience continued**
 - ▶ Response to current therapy
 - ◇ Pain relief
 - ◇ Patient adherence to medication plan
 - ◇ Medication adverse effects such as constipation, sedation, cognitive slowing, nausea, and others
 - ▶ Breakthrough pain is episodic pain not controlled with existing pain regimen; see breakthrough pain on [PAIN-G, 2 of 13](#).
 - ▶ Prior pain therapies
 - ◇ Reason for use, length of use, response, reasons for discontinuing, and adverse effects encountered
 - ▶ Special issues relating to pain
 - ◇ Meaning and consequences of pain for patient and family/caregiver
 - ◇ Patient and family/caregiver knowledge and beliefs surrounding pain and pain medications
 - ◇ Cultural beliefs toward pain, pain expression, and treatment
 - ◇ Spiritual, religious considerations, and existential suffering
 - ◇ Patient goals and expectations regarding pain management
 - ◇ Assess for use of integrative therapies ([See PAIN-D](#))
 - ◇ Screen for potential adverse interactions or effects
 - ◇ Assess risk of opioid abuse/misuse/diversion
- List of potential risk factors for misuse/abuse ([See PAIN-G, 4 of 13](#))

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Return to Universal Screening \(PAIN-2\)](#)

PAIN-B
1 OF 3



COMPREHENSIVE PAIN ASSESSMENT

- **Psychosocial Support** ([See PAIN-H](#)) ([See NCCN Guidelines for Palliative Care](#))
 - ▶ **Patient distress** ([See NCCN Guidelines for Distress Management](#))
 - ▶ **Family and other support; assess impact and burden on caregiver and recommend resources as appropriate**
 - ▶ **Psychiatric history including current or prior patient, family/caregiver, or household history of substance abuse**
 - ▶ **Risk factors for aberrant use or diversion of pain medication** ([See PAIN-G, 4 of 13](#))
 - ◇ **Patient, environmental, and social factors as identified by a detailed patient evaluation¹ and/or screening tools at initiation of care (eg, SOAPP®-R², ORT³) and monitoring of ongoing analgesic use (eg, COMM).⁴ (Specific screening tools have not been validated in the setting of cancer care).⁵** ([See PAIN-G, 6 of 13](#))
 - ▶ **Risk factors for undertreatment of pain**
 - ◇ **Geriatric, minority, or female patients; communication barriers; history of substance abuse; neuropathic pain; cultural factors**
- **Medical history**
 - ▶ **Oncologic treatment including current and prior chemotherapy, hormonal therapy, RT, and surgery**
 - ▶ **Other significant illnesses, conditions**
 - ▶ **Pre-existing chronic pain**
- **Clinical assessment, physical examination, and laboratory and imaging studies to evaluate for disease progression**

¹ Moore TM, Jones T, Browder JH, Daffron S, Passik SD. A comparison of common screening methods for predicting aberrant drug-related behavior among patients receiving opioids for chronic pain management. *Pain Medicine* 2009;10:1426-1433.

² Butler SF, Fernandez K, Benoit C, et al. Validation of the revised Screener and Opioid Assessment for Patients with Pain (SOAPP-R). *J Pain* 2008;9:360-372.

³ Webster LR and Webster RM. Predicting aberrant behaviors in opioid-treated patients: Preliminary validation of the Opioid Risk Tool. *Pain Med* 2005;6:432-442.

⁴ Meltzer EC, Rybin D, Saitz R, et al. Identifying prescription opioid use disorder in primary care: diagnostic characteristics of the current opioid misuse measure (COMM). *Pain* 2011;152:397-402.

⁵ Angheliescu DL, Ehrentraut JH, Faughnan LG, et al. Opioid misuse and abuse: risk assessment and management in patients with cancer pain. *J Natl Compr Canc Netw* 2013;11:1023-1031.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Return to Universal Screening \(PAIN-2\)](#)



IMPACT OF PAIN MEASUREMENT^{6,7}

Mark the number that describes how much, in the past [week/24 hours], pain has interfered with your:

1. General Activity 0 1 2 3 4 5 6 7 8 9 10 Does not Interfere Completely Interferes
2. Mood 0 1 2 3 4 5 6 7 8 9 10 Does not Interfere Completely Interferes
3. Walking Ability 0 1 2 3 4 5 6 7 8 9 10 Does not Interfere Completely Interferes
4. Normal Work (includes both work outside the home and housework) 0 1 2 3 4 5 6 7 8 9 10 Does not Interfere Completely Interferes
5. Relations with other people 0 1 2 3 4 5 6 7 8 9 10 Does not Interfere Completely Interferes
6. Sleep 0 1 2 3 4 5 6 7 8 9 10 Does not Interfere Completely Interferes
7. Enjoyment of life 0 1 2 3 4 5 6 7 8 9 10 Does not Interfere Completely Interferes

⁶ Used with permission from Cleeland CS, Nakamura Y, Mendoza TR, et al. Dimensions of the impact of cancer pain in a four country sample: New information from multidimensional scaling. Pain 1996;67:267-273.

⁷ For the complete Brief Pain Inventory assessment tool, [see mdanderson.org/bpi](http://see.mdanderson.org/bpi).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PSYCHOSOCIAL SUPPORT****General**

- Due to the complexity of cancer-related pain and associated symptoms, health care providers should anticipate patients' and families' need for support and education in management strategies.
- Assessing each patient's need for psychosocial support is an essential component of a comprehensive pain assessment. ([See PAIN-B](#)).

Support

- Inform patient and family/caregiver that emotional reactions to pain are normal and are evaluated and treated as part of pain treatment.
- Provide emotional support to patient and family/caregiver that acknowledges that the pain is a problem to be addressed.
- Assist in accessing treatment as needed.
- State that you will work together with the patient and family/caregiver as part of the team to address the pain problem.
- Describe the mutually agreed upon plan of care to be taken and when results can be expected.
- Express your commitment to being available to help with pain management.
- Inform patient and family/caregiver that there is always something else that can be done to try to adequately manage pain and other noxious symptoms.
- Assess impact upon family and significant others; provide education and support as indicated.
- Verbally repeat your concern and the plan of action to be taken.
- Consider referral to spiritual care provider. ([See NCCN Guidelines for Distress Management](#))

Skills Training

- Teach coping skills (to be used in conjunction with and not in lieu of appropriate analgesia) to provide pain relief, enhance a sense of personal control, and refocus energy on optimizing quality of life.
 - ▶ Consider referral to a licensed mental health professional who is trained in any of the following domains: cognitive behavioral therapy (CBT), hypnosis, biofeedback, and mindfulness-based stress reduction (MBSR).
 - ▶ Coping skills for acute pain include breathing exercises and distraction techniques.
 - ▶ Coping skills for chronic pain (not pain emergency) include all of the above, plus relaxation techniques, guided imagery, graded task assignments, hypnosis to maximize function, cognitive restructuring, and behavioral activation.
 - ▶ Provide training on how to encourage assertiveness to maximize comfort.
- Educate patient and family/caregiver that in pain management a team effort is necessary to comprehensively assess and treat the impact of pain. Members of the team may include: oncologist, nurse, palliative care clinician, physiatrist, neurologist, psychologist, social worker, psychiatrist, physical therapist, and spiritual counselor. ([See Patient and Family/Caregiver Education \(PAIN-I\)](#))

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



INTEGRATIVE INTERVENTIONS

Consider integrative interventions in conjunction with pharmacologic interventions as needed. Integrative interventions may be especially important in vulnerable populations (eg, frail, elderly) in whom standard pharmacologic interventions may be less tolerated or based on patient preference. The utility of integrative interventions underscores the necessity for pain management to be carried out with a team approach that contains a wide range of treatment options. ([See PAIN-L](#))

Pain likely to be relieved or function improved with cognitive, physical, or interventional modalities:

- **Cognitive modalities**

- ▶ Mindfulness-based stress reduction (MBSR)
- ▶ Imagery
- ▶ Hypnosis
- ▶ Biofeedback
- ▶ Acceptance-based training
- ▶ Distraction training
- ▶ Relaxation training
- ▶ Active coping training
- ▶ Graded task assignments, setting goals, pacing, and prioritizing
- ▶ Cognitive behavioral therapy (CBT), cognitive restructuring
- ▶ Behavioral activation

- **Nutritional modalities**

- ▶ Nutrition consult
- ▶ Dietary recommendations
- ▶ Assess and educate on herbal, botanical, and dietary supplements

- **Spiritual care** ([See NCCN Guidelines for Distress Management](#))

- **Physical modalities**

- ▶ Bed, bath, and walking supports
- ▶ Positioning instruction
- ▶ Instruction in therapeutic and conditioning exercise
- ▶ Energy conservation, pacing of activities
- ▶ Massage
- ▶ Heat and/or ice
- ▶ Transcutaneous electrical nerve stimulation (TENS)
- ▶ Acupuncture, electro-acupuncture, or acupressure
- ▶ Ultrasonic stimulation

- [See Interventional Strategies \(PAIN-M\)](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**NON-OPIOID ANALGESIC (NONSTEROIDAL ANTI-INFLAMMATORY DRUGS [NSAIDS] AND ACETAMINOPHEN) PRESCRIBING****Acetaminophen**

- Acetaminophen, 650 mg every 4 hours or 1 g every 6 hours (daily maximum 4 g/day) in adult patients with normal liver function. For chronic administration, consider limiting the maximum daily dose to 3 g/day or less due to concerns for hepatic toxicity.
- Due to concerns with liver toxicity, acetaminophen should be used with caution or not used at all with combination opioid-acetaminophen products to prevent excess acetaminophen dosing.
- See the FDA website (www.fda.gov) for the latest information on acetaminophen adverse effects and dosing.
- Consider OTC medications as additional sources of acetaminophen.

NSAIDs

- Use NSAIDs with caution, especially for chronic use, as many oncology patients may be at high risk for renal, GI (ie, upper GI surgery, RT), or cardiac toxicities, thrombocytopenia, or bleeding disorder.
 - ▶ NSAID use should be coordinated with other oncologic therapies.
 - ▶ The FDA warns that NSAID use increases the risk of heart attack or stroke. <http://www.fda.gov/Drugs/DrugSafety/ucm451800.htm>
- Note that the potential adverse effects of chemotherapy (especially angiogenesis inhibitors), such as hematologic (ie, thrombocytopenia, coagulopathy), renal, hepatic, and cardiovascular toxicities, can be increased by the concomitant prescription of NSAIDs.
- For some patients opioid analgesics may be a safe and effective alternative analgesic to NSAIDs.
- Use any NSAID that the patient has found to be effective and well tolerated in the past; otherwise, consider ibuprofen to the maximal dose. Shortest duration, lowest effective dose of NSAIDs should be used while a safer, long-term pain management strategy is being developed.
 - ▶ Ibuprofen, 400 mg four times daily (daily maximum = 3200 mg); or naproxen 220–500 mg 2–3 times daily (daily maximum of 1500 mg). If needed, consider short-term use of ketorolac, 15–30 mg IV every 6 hours for a maximum of 5 days.
 - ▶ Compounds that do not inhibit platelet aggregation:
 - ◊ Nonacetylated salicylate (eg, salsalate), 2–3 g/day in two or three divided doses; magnesium salicylate, OTC
 - ◊ Celecoxib (selective COX-2 inhibitor) up to 200 mg/twice daily.
- Consider topical NSAID - diclofenac gel 1% 4 times/day; or diclofenac patch 180 mg, 1–2 patches/day

[See NSAIDs
and Toxicities
\(PAIN-E, 2 of 2\)](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**NON-OPIOID ANALGESIC (NONSTEROIDAL ANTI-INFLAMMATORY DRUGS [NSAIDS] AND ACETAMINOPHEN) PRESCRIBING****NSAIDs and Toxicities****• Monitoring for NSAID toxicities**

- ▶ Baseline blood pressure, blood urea nitrogen (BUN), creatinine, liver function studies (alkaline phosphatase, lactate dehydrogenase [LDH], serum glutamic-oxaloacetic transaminase [SGOT], serum glutamic-pyruvic transaminase [SGPT]), and complete blood count (CBC). Repeat these evaluations as clinically indicated.

• Further NSAID considerations

- ▶ If two NSAIDs are tried in succession without efficacy, use another approach to analgesia.
- ▶ If NSAIDs are effective but treatment is limited by toxicities that are not deemed serious, consider trial of another NSAID.
- ▶ When systemic administration is not feasible, consider topical NSAID preparations in place of oral NSAIDs.
- ▶ Toxicity of anti-cancer treatment may increase the risk profile of anti-inflammatory treatment.

Cardiac Toxicities

- Patients at high risk: history of cardiovascular disease or at risk for cardiovascular disease or complications.¹
- The use of concomitant NSAID with prophylactic aspirin may reduce the effectiveness of aspirin. Therefore, it is recommended to either avoid use or take separately to avoid this possibility.
- Treatment: discontinue NSAID if congestive heart failure or hypertension develops or worsens. All NSAIDs have been associated with cardiac toxicities.

Hematologic Toxicities

- NSAIDs taken with prescribed anticoagulants, such as warfarin or heparin, may significantly increase the risk of bleeding complications.
- Avoid the use of oral NSAIDs in the setting of prophylactic or therapeutic anticoagulation. Celecoxib or topical NSAIDs such as diclofenac gel or patch may be useful in this population.

Renal Toxicities

- Patients at high risk: age >60 years, compromised fluid status, multiple myeloma, diabetes, interstitial nephritis, papillary necrosis, and concomitant administration of other nephrotoxic drugs (including cyclosporine, cisplatin) and renally excreted chemotherapy.
- Treatment: reevaluate NSAID use if renal function deteriorates or if hypertension develops or worsens.
- All NSAIDs with systemic administration have been associated with renal toxicities.

GI Toxicities

- Patients at high risk: age >60 years, history of peptic ulcer disease or significant alcohol use (3 or more alcoholic beverages/day), major organ dysfunction including hepatic dysfunction, high-dose NSAIDs given for long periods, concomitant steroid use, cardioprotective dose of daily aspirin, and concomitant use of selective serotonin reuptake inhibitor (SSRI) antidepressants.
- Treatment:
 - ▶ If patient develops gastric upset or nausea, consider discontinuing NSAID or changing to selective COX-2 inhibitor. COX-2 inhibitors are associated with lower incidence of GI adverse effects and do not inhibit platelet aggregation; however, they have not been demonstrated to have reduced renal adverse effects.
 - ▶ As prophylaxis for NSAID peptic ulceration, consider adding misoprostol or proton pump inhibitors. If patient develops GI peptic ulcer or GI hemorrhage, discontinue NSAID.
 - ▶ Discontinue NSAID if liver function studies increase 1.5 times the upper limit of normal.

¹ Antman EM, Bennett JS, Daugherty A, et al. Use of nonsteroidal antiinflammatory drugs: an update for clinicians. A scientific statement from the American Heart Association. *Circulation* 2007;115:1634-1642.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



ADJUVANT ANALGESICS FOR NEUROPATHIC PAIN (ANTIDEPRESSANTS, ANTICONVULSANTS, AND TOPICAL AGENTS)

Principles of Adjuvant Analgesic Use

- Antidepressants and anticonvulsants are first-line adjuvant analgesics for the treatment of cancer-related neuropathic pain.
- These drugs can also be used in combination with opioids, for patients whose pain is otherwise inadequately controlled.
- The use of adjuvant analgesics in the cancer population is often based on guidelines or experience derived from data for the treatment of pain not caused by cancer (non-malignant pain).
- Effective use is predicated on an assessment that clarifies the nature of the pain as most adjuvant analgesics are more likely to be effective in management of neuropathic pain.
- As with opioids, response to adjuvant analgesics may vary according to the etiology of neuropathic pain and the individual patient. Failure to control pain with one agent in a particular class does not mean the entire class of medications will not work.
- Drug selection may be influenced by other symptoms and comorbidities. For example, a sedating drug may be useful in a patient in whom insomnia is a problem.
- Patient education should emphasize the trial and error nature of the treatment so patients do not get discouraged.
- Doses should be increased until the analgesic effect is achieved, adverse effects become unmanageable, or the conventional maximal dose is reached.
- For information on cannabinoids and medical marijuana/cannabis, [see Discussion](#).

[See Examples of Adjuvant Analgesics
Use for Neuropathic Pain \(PAIN-F, 2 of 2\)](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**ADJUVANT ANALGESICS FOR NEUROPATHIC PAIN**
(ANTIDEPRESSANTS, ANTICONVULSANTS, TOPICAL AGENTS, AND CORTICOSTEROIDS)**Examples of Adjuvant Analgesics Use**

- Extrapolated from non-cancer neuropathic pain management
- Both antidepressants and anticonvulsants are frequently used as an adjuvant analgesic in combination with an opioid to treat neuropathic components of pain.

Antidepressants: Analgesic effectiveness is not dependent on its antidepressant activity. Effective analgesic dose 1) may be lower than that required to treat depression; and 2) the onset of analgesic relief may occur earlier than anti-depressive effects.

- Frequently used as an adjuvant analgesic in combination with an opioid for the neuropathic component of the pain.
- Check for drug interactions with special regard to serotonergic medications due to risk for serotonin syndrome.
- TCAs (eg, amitriptyline, imipramine, nortriptyline, desipramine)
 - TCAs should be used with caution in patients with conduction abnormalities, including QTc prolongation, or ischemic heart disease
 - Start with low dose and increase every 5–7 days if tolerated (eg, nortriptyline and desipramine starting dose 10–25 mg nightly with increase to 50–150 mg nightly). The tertiary amines (ie, amitriptyline, imipramine) may be more efficacious but secondary amines (ie, nortriptyline, desipramine) are better tolerated. Anticholinergic adverse effects such as sedation, dryness of mouth, and urinary hesitancy are more likely to occur with amitriptyline and imipramine.
- Serotonin-norepinephrine reuptake inhibitors (SNRIs) (dose adjustment is required for those with renal and/or hepatic insufficiency):
 - Duloxetine- Starting dose 20–30 mg daily; after 1 week, increase to 60 mg daily as needed/tolerated (doses up to 120 mg per day may be useful for mood disorders) (there are data supporting use of duloxetine for chemotherapy-induced peripheral neuropathy)
 - Venlafaxine- Starting dose 37.5 mg daily, increase every 4 days as needed, up to 75–225 mg daily (dose must be at least 150 mg per day to achieve SNRI effects/analgesia)

Anticonvulsants: Frequently used as an adjuvant analgesic in combination with an opioid for the neuropathic component of the pain.

- Anticonvulsants examples:
 - Gabapentin- Starting dose 100–300 mg nightly, increase to 900–3600 mg daily in divided doses 2–3 times a day. Dose increments of 50%–100% may occur as often as every 3 days. Slower titration is needed for the elderly or medically frail. Dose adjustment is required for those with renal insufficiency.
 - Pregabalin- Starting dose 75 mg twice a day, with increasing dose increments of 50%–100% every 3 days to a maximum daily dose of 600 mg. Slower titration is needed for the elderly or medically frail. Dose adjustment is required for those with renal insufficiency. Pregabalin is more efficiently absorbed through the GI tract than gabapentin.
 - Consider other anticonvulsant agents, many of which have been shown to have efficacy in non-cancer neuropathic pain.

Topical Agents: Act locally and may be used as an adjuvant analgesic in combination with an opioid, antidepressant, and/or an anticonvulsant.

- Topical agent examples:
 - Lidocaine patch- 5% - Apply daily to the painful site; minimal systemic absorption. Continuous application to a given area may increase likelihood of cutaneous damage over time.

Corticosteroids: Typically dexamethasone (due to less mineralocorticoid effect). Long half-life of these drugs allows for once-daily dosing, preferably in the morning due to their stimulating effect and to prevent nighttime insomnia. Useful in the acute management of a pain crisis when neural structures or bones are involved. Long-term adverse effects are significant.

Note: Some SSRI and SNRI antidepressants may inhibit the conversion of tamoxifen to its active metabolite, thereby decreasing the effectiveness of tamoxifen. [See Discussion](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY****General Principles**

- Periodically review prescription drug monitoring program (PDMP) databases.
- Consider documentation of opioid and controlled substance agreement.
- Dose and titrate with caution in patients with risk factors such as decreased renal/hepatic function, chronic lung disease, upper airway compromise, sleep apnea, and poor performance status.
- The appropriate opioid dose is the lowest dose that relieves the patient's pain and maximizes function throughout the dosing interval without causing unmanageable adverse effects.
- Generally, oral route is most common; however, other routes (ie, IV, subcutaneous, rectal, transdermal, transmucosal) can be considered as indicated to maximize patient comfort. For intrathecal route administration, [see PAIN-M](#).
- Calculate dosage increase based upon total opioid dose (around the clock/scheduled and as needed) taken in the previous 24 hours and increase both around-the-clock and as-needed dose as required. The rapidity of dose escalation should be related to the severity of the symptoms, expected analgesic onset and duration, and ability to monitor during dose titration.
 - ▶ When initiating opioid therapy or making significant dose adjustments, close follow-up should be considered.
- According to FDA guidelines, when higher doses of analgesic are needed, switch from preparations of opioid combined with other medications [such as aspirin or acetaminophen] to a pure opioid preparation to allow optimized titration of both agents. [\(See PAIN-E\)](#)
- Steady state drug levels will be achieved when a stable drug dose has been routinely administered for a period equal to 5 times the drug elimination half life.
- Consider opioid rotation if pain is inadequately controlled and further dose titration is limited by adverse effects. Other indications for switching to a different opioid include: out-of-pocket costs, limitations based on insurance formularies, or change in a patient's condition (eg, dysphagia, NPO [nothing by mouth] status, initiation of tube feeding, renal and/or hepatic function).
- For breakthrough pain, [see PAIN-G, 2 of 18](#).
- For opioid dose reduction, [see PAIN-G, 3 of 18](#).
- Patient evaluations should include assessment of risk factors for aberrant use of pain medications. [\(See PAIN-G, 6 of 18\)](#)
- Educate the patients and caregivers about safe use, storage, and disposal of opioids. [\(See PAIN-I\)](#)
- Use caution when combining opioid medications with other medications that have a sedating effect (eg, benzodiazepines). <http://www.fda.gov/downloads/drugs/drugsafety/ucm518672.pdf>
- Consider pain or palliative care consult. [See Management of Pain in Opioid-Tolerant Patients \(PAIN-4\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

References

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY****Principles of Maintenance Opioid Therapy**

- For continuous pain, it is appropriate to give pain medication on a regular schedule with supplemental doses for breakthrough pain.
- Add extended-release or long-acting formulation to provide background analgesia for control of chronic persistent pain controlled on stable doses of short-acting opioids.
 - ▶ Initial range for converting to long-acting opioid would be 50% to 100% of the daily requirement, depending on expected pain natural history.
 - ▶ If using different forms of long-acting and short-acting opioids, particular care must be taken with conversions and appropriate dosing.
- When using methadone as a long-acting opioid, a short-acting opioid should also be provided for breakthrough pain.
- Increase dose of regularly scheduled opioid if patient persistently needs doses of as-needed opioids or when dose of around-the-clock opioid fails to relieve pain at peak effect or at end of dose.
- Breakthrough pain (pain that fails to be controlled or “breaks through” a regimen of regularly scheduled opioid) may require additional doses of opioid for pain not relieved by regular schedule of long-acting (eg, extended-release) opioid. Breakthrough pain may be further evaluated into the following categories, which have direct impact on treatment:
 - ▶ Incident pain: Pain associated with or incident to specific activities or events, potentially managed with short-acting opioid given in anticipation of those events (eg, physical therapy, exercise, or routine procedures that may induce pain)
 - ▶ End-of-dose failure pain: Pain recurring towards the end of dosing interval for regularly scheduled opioid, potentially managed by increasing the dose or frequency of regularly scheduled opioid
 - ▶ Uncontrolled persistent pain: Pain routinely uncontrolled by existing regularly scheduled opioid, potentially managed by adjusting dose of regularly scheduled opioid
- Allow rescue doses of short-acting opioids (10% to 20% of the 24-hour total of long-acting or regularly scheduled oral opioid dose) up to every 3–4 hours as needed. Titrate rescue dose as needed.
 - ▶ If pain is inadequately controlled, to allow for dose titration, the short-acting opioids could be given as often as once per hour as needed (if hourly dosing is needed for more than 3 cycles, reassessment or other intervention is recommended).
- Consider rapidly acting transmucosal fentanyl (various formulations and delivery systems are available) in opioid-tolerant patients for brief episodes of incident pain not relieved by traditional immediate-release opioids and not attributed to inadequate dosing of around-the-clock opioid (transmucosal fentanyl requires REMS certification [see [PAIN-G 3 of 18](#)]).
 - ▶ Data do not support a specific transmucosal fentanyl dose equianalgesic to other opioids or between different transmucosal formulations. Always initiate transmucosal fentanyl with lowest dose in chosen formulation and titrate to effect. (See specific transmucosal prescribing information for appropriate dosing intervals.)
- Continue to monitor patients for opioid adverse effects and patients/family for abnormal patterns of opioid use that may suggest aberrant drug use and/or diversion. ([See PAIN-G, 6 of 18](#))
- Consider potential drug interactions. ([See Table 1 on MS-30](#))

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)



OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY

Principles of Opioid Dose Reduction

- Consider opioid dose reduction by 10% to 20% when possible; situations that may warrant dose reduction include:
 - ▶ Patient never or rarely needs breakthrough analgesic
 - ▶ Completion of acute pain event or response to cancer-directed therapies
 - ▶ Improvement of pain control through use of non-opioid pain management therapies
- If patient is experiencing unmanageable opioid-related adverse effects, [see PAIN-H](#). If pain is 3 (mild), consider downward dose titration by approximately 10% to 25% and reevaluate. Close follow-up is required to make sure that the pain does not escalate, and that the patient does not develop symptoms of withdrawal.
 - ▶ If patient has rapid clinical deterioration (eg, marked sedation due to sepsis), temporary opioid dose reduction by 50% to 75% may be necessary.
- If pain is worsened with increasing dose, consider opioid-induced hyperalgesia; opioid dose reduction or rotation with attention to other pain therapies may be indicated.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY****Opioids and Risk Evaluation and Mitigation Strategy (REMS)**

- Opioids are the principal analgesics for moderate to severe pain, yet opioids pose risks to patients and society. In 2018, 67,367 drug overdose deaths occurred in the United States, including 46,802 deaths involving opioids. Drug poisoning still remains the number one cause of injury-related deaths.¹ Most people who overdose on prescription opioids not prescribed to them have been given (not bought or stolen) opioids from friends or family. [See CDC Drug Overdose Death Data \(January 2020\)](#).
- Responding to the “public health crisis of addiction, misuse, abuse, overdose, and death,” the FDA established REMS programs for all potent opioid products. [See Opioid Drugs and Risk Evaluation and Mitigation Strategies \(REMS\)](#). Provider and patient education are the principal recommendations of proposed opioid REMS programs. Highlights include:
 - ◊ Patient’s therapeutic response to opioid therapy should be regularly evaluated as to patient treatment goals of therapy.
 - ◊ Prescriber should routinely evaluate each patient for risk factors associated with opioid misuse/abuse/diversion.
 - ◊ Prescriber should educate each patient on safe use, storage, and disposal of opioid. [\(See PAIN-I\)](#)
 - ◊ Prescriber should routinely monitor patients for opioid misuse or abuse. Different screening tools have been described for this purpose but have yet to be evaluated in cancer-related pain.² If signs of aberrant opioid use are observed, urine drug testing and limiting or restricting use accordingly to minimize risk of diversion is strongly encouraged.
 - ◊ Make use of state PDMPs. The National Association of State Controlled Substances Authorities [\(https://www.nascsa.org\)](https://www.nascsa.org) maintains a database of state PDMP contacts.
- REMS programs are currently in place for all opioid analgesics. [\(https://opioidanalgesicrems.com\)](https://opioidanalgesicrems.com)
 - ▶ Registration is required in order to prescribe transmucosal fentanyl products. [\(https://www.tirfremssaccess.com\)](https://www.tirfremssaccess.com)
- It is important for prescribers to be aware of the range of opioid use patterns to detect any potential aberrant behaviors. [\(See PAIN-G, 6 of 18\)](#)
 - ▶ Potential risk factors for misuse/abuse of prescribed analgesics include:
 - ◊ Patients with a history of prescription, illicit drug, or alcohol dependence/substance abuse
 - ◊ Patients who have a history of binge drinking or peers who binge drink
 - ◊ Patients who have a family history of substance abuse
 - ◊ Patients with a history of psychiatric disorder, including anxiety, depression, attention deficit hyperactivity disorder (ADHD), post-traumatic stress disorder (PTSD), bipolar disorder, or schizophrenia
 - ◊ Patients who have a history of sexual abuse victimization may be at increased risk for prescribed medication misuse/abuse
 - ◊ Young age (<45 years)
 - ◊ Patients with a history of legal problems or incarceration
- Patients receiving treatment for addiction should be encouraged to continue with therapy and pain management should be carried out in coordination with an addiction specialist.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)

OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY

Table 1. Glossary of Terms Related to Opioid Misuse²

Abuse	A maladaptive pattern of a prescription opioid use leading to clinically significant impairment and/or distress
Addiction	The aberrant use of a substance characterized by <ul style="list-style-type: none"> • loss of control, craving • compulsive use and preoccupation • continued use despite harm
Chemical coping³	Misuse of medication in a non-prescribed way to cope with the various stressful events associated with the diagnosis and management of cancer
Diversion	The transfer of a prescribed medication from the person for whom it was prescribed to another person
Misuse	The inappropriate use of a prescription drug, whether intentional or unintentional, and regardless of motivation
Opioid use disorder	A problematic pattern of opioid use leading to clinically significant impairment or distress⁴
Physical dependence	Pharmacologic property of some drugs, defined solely by the occurrence of an abstinence syndrome after abrupt dose reduction, discontinuation of dosing, or administration of an antagonist drug
Pseudoaddiction	Distress and perceived drug-seeking behaviors that occur in the context of unrelieved pain. These behaviors subside when analgesia is achieved
Tolerance	Diminution of one or more drug effects (either favorable or adverse effects) caused by exposure to the drug; may be pharmacologic or associative (related to learning)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY****Strategies to Maintain Patient Safety and Minimize the Risk of Opioid Misuse and Abuse During Chronic Opioid Use²**

- Use caution when combining opioid medications with other medications that have a sedating effect (eg, benzodiazepines).
<http://www.fda.gov/downloads/drugs/drugsafety/ucm518672.pdf>
- **Risk assessment** prior to and during treatment is recommended, although current assessment tools have not been validated in the setting of cancer care and clinical judgment should be exercised.
 - ▶ [The Screener and Opioid Assessment for Patients with Pain-Revised \(SOAPP®-R\)](#)
 - ▶ [The Opioid Risk Tool \(ORT\)](#)
 - ▶ [Current Opioid Misuse Measure \(COMM\)](#)
 - ▶ Comprehensive psychological evaluation can be helpful in assessing risk for substance use disorders.
- **Educate** regarding the potential risks and benefits of opioid therapy; educate regarding not sharing opioids with family members or friends.
 - ▶ Discuss the purpose of the assessment and reassure that responses will not prevent receiving appropriate treatment.
 - ▶ Provide guidance and education about the potential for diversion and misuse of opioids and the addictive potential associated with prescription opioids.
- **Educate regarding safe manipulation, storage, and disposal of controlled substances.** These interventions contribute to maintaining a safe community and minimize opioid misuse and abuse in the community. ([See PAIN-I](#))
 - ▶ Encourage use of community take-back programs for disposal of unneeded controlled substances where available; otherwise, FDA regulations recommend flushing unneeded opioids down the toilet: https://www.fda.gov/drugs/safe-disposal-medicines/disposal-unused-medicines-what-you-should-know#Flush_List

Risk Mitigation for All Patients Receiving Opioid Analgesics

- Consider prescribing naloxone for administration by caregivers in the event of respiratory depression and sedation.
 - ▶ Ensure education of caregivers in the proper indications and usage of naloxone. <https://store.samhsa.gov/system/files/sma18-4742.pdf>
 - ▶ Instruct caregivers to call emergency services (911) if naloxone is administered.
- Pain medication diaries are recommended for patients to document the dose and/or number of tablets and the date and time taken.
- Pill counts may be used at outpatient visits or by home health/hospice to assist in correct use of medication.
- Urine drug testing at baseline and during treatment should be strongly considered to help document opioid analgesic adherence, detect illegal drug use, and identify opioid diversion. Urine drug testing via immunoassay (eg, point of care) is susceptible to false positives/negatives. For unexpected results prescribers should request confirmatory testing and/or input from a certified laboratory professional or toxicologist.⁵

High-risk patients who exhibit one or more opioid misuse and abuse risk factors may benefit from additional education and support services. Behavioral and cognitive-behavioral interventions may increase a patient's ability to implement problem-solving strategies and reduce the impact of modifiable risk factors.

- Increase frequency of outpatient visits weekly, if possible, and/or reduce quantity of drug prescribed per prescription.
- Consider earlier referral to interventional pain specialist to maximize non-opioid options for pain control.
- Consider referral to interdisciplinary team including an addiction specialist.
- Counsel high-risk patients that continuation of opioid therapy is contingent upon appropriate, safe use of prescribed analgesics.
- Consider utilizing programmable electronic medication dispensers.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)



OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY

Table 2. Oral and Parenteral Opioid Equivalences and Relative Potency of Drugs as Compared with Morphine Based on Single-Dose Studies

Opioid Agonists	Parenteral Dose	Oral Dose	Factor (IV to PO)	Duration of Action ^j
Morphine ^{a,b}	10 mg	30 mg	3	3–4 h
Hydromorphone ^a	1.5 mg	7.5 mg	2.5–5	2–3 h
Fentanyl ^c	0.1 mg	–	–	–
Methadone ^{d,e}	–	–	–	–
Oxycodone	–	15–20 mg	–	3–5 h
Hydrocodone ^{a,f}	–	30–45 mg	–	3–5 h
Oxymorphone ^a	1 mg	10 mg	10	3–6 h
Codeine ^{a,g}	–	200 mg	–	3–4 h
Tramadol ^h	100 mg	300 mg	3	6 h
Tapentadol ⁱ	–	75–100 mg	–	–

NOT RECOMMENDED
Meperidine^k
Mixed agonist-antagonists^l
(pentazocine, nalbuphine, butorphanol)

[See Miscellaneous Analgesics \(PAIN-G, 8 of 18\)](#)

^a Codeine, morphine, hydromorphone, hydrocodone, and oxymorphone should be used with caution in patients with fluctuating renal function due to potential accumulation of renally cleared metabolites—monitor for neurologic adverse effects.

^b Conversion factor listed for chronic dosing.

^c In single-dose administration, 10 mg IV morphine is equivalent to approximately 100 mcg IV fentanyl but with chronic fentanyl administration, the ratio of 10 mg IV morphine is equivalent to approximately 250 mcg IV fentanyl. For transdermal fentanyl conversions, ([See PAIN-G, 13 of 18](#)).

^d Long half-life: observe for drug accumulation and adverse effects, especially over first 4–5 days. In some individuals, steady state may not be reached for several days to 2 weeks. Methadone is typically dosed every 8–12 h.

^e The oral conversion ratio of methadone varies. PRACTITIONERS ARE ADVISED TO CONSULT WITH A PAIN OR PALLIATIVE CARE SPECIALIST IF THEY ARE UNFAMILIAR WITH METHADONE PRESCRIBING. ([See Special Notes Regarding Oral Methadone, PAIN-G, 15 of 18](#)).

^f Equivalence data not substantiated. Clinical experience suggests use as a mild, initial-use opioid but effective dose may vary. Immediate-release hydrocodone is only available commercially combined with acetaminophen (325 mg/tablet) or ibuprofen (200 mg/tablet). The FDA has limited the amount of acetaminophen in all prescription drug products to no more than 325 mg per dosage unit. Dosage must be monitored for safe limits of acetylsalicylic acid (ASA) or acetaminophen.

^g Codeine has no analgesic effect unless it is metabolized into morphine by hepatic enzyme CYP2D6 and then to its active metabolite morphine-6-glucuronide by phase II metabolic pathways. Individuals with low CYP2D6 activity may receive no analgesic effect from codeine, but rapid metabolizers may experience toxicity from higher morphine production. Dosage must be monitored for safe limits as it may be available in combination with ASA or acetaminophen. Dose listed refers only to opioid portion.

^h The manufacturer recommends a maximum single dose of tramadol not to exceed 100 mg, with a maximum daily dose of 400 mg for IR formulations (300 mg/day in older adults, 200 mg/day for renal impairment) or 300 mg/day for ER formulations.

ⁱ The maximum daily dose for tapentadol ER is 500 mg, or 600 mg IR (lower doses are recommended for moderate hepatic impairment; avoid with severe impairment).

^j Shorter time generally refers to parenterally administered opioids (except for controlled-release products, which have some variability); longer time generally applies to oral dosing.

^k Not recommended for cancer pain management because of CNS toxic metabolite - normeperidine.

^l Mixed agonists-antagonists have limited usefulness in cancer pain; however, they can be used to treat opioid-induced pruritus. They should NOT be used in combination with opioid agonist drugs. Converting from an agonist to an agonist-antagonist could precipitate a withdrawal crisis in the opioid-dependent patient.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY****Mixed-Mechanism Drugs:**

- Tramadol is a weak mu-opioid agonist with some norepinephrine and serotonin reuptake inhibition used for mild to moderate pain. A maximum daily dose of 400 mg (100 mg four times daily) is recommended for adults with normal hepatic and renal function, and lower daily doses are recommended for older adults (≥75 years) and those with hepatic and/or renal dysfunction, to reduce the risk of seizures. Even at a maximum dose of 100 mg four times daily, tramadol is less potent than other opioid analgesics such as morphine.
- Tapentadol⁶ is a mu-opioid analgesic with norepinephrine reuptake inhibition for treatment of moderate to severe pain. Typical doses would start at 50 to 100 mg PO every 4 hours as needed, with a maximal daily dose of 500 mg per day (if using the extended release) or 600 mg per day (if using the immediate release only) due to lack of published data regarding higher doses. Some comparative data suggest tapentadol may have a lower incidence of GI adverse effects than oxycodone.
- Tramadol and tapentadol should be used with caution or avoided in patients taking other serotonergic or monoamine oxidase inhibitor (MAOI)-like medications (eg, tricyclic antidepressants [TCAs], SSRIs, SNRIs) due to risk of serotonin syndrome.

Non-Opioid Analgesic (given in collaboration with a pain/palliative care specialist):

- Ketamine⁷ is a noncompetitive NMDA receptor antagonist that blocks glutamate. Low (subanesthetic) doses produce analgesia and modulate central sensitization, hyperalgesia, and opioid tolerance. There are only limited data regarding the use of ketamine as an adjuvant to opioids for management of cancer pain.
- Intravenous lidocaine infusion may be a useful therapy for refractory pain.⁸

Buprenorphine:

- Potential advantages of buprenorphine:
 - Lower abuse potential than other potent opioids.⁹
 - Causes less analgesic tolerance and less constipation when compared to other mu-receptor agonists with no effects on the sphincter of Oddi.¹⁰
 - Has a ceiling effect for respiratory depression.¹¹
 - Can affect QTc interval, although not to the same degree as methadone.¹²
 - May cause less cognitive impairment and has been safely used in older adults.¹³
 - No dose adjustment required for patients with renal failure or renal insufficiency.¹⁴
 - Less drug dependence and milder withdrawal symptoms.¹⁵

[Continued](#)[References](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY****Buprenorphine (continued):****• Buprenorphine for cancer pain:**

- ▶ Buprenorphine is increasingly recognized as an effective analgesic with an improved therapeutic index relative to certain potent opioids, however, it has not been extensively studied in cancer pain. Its use in cancer pain is extrapolated from data on its effectiveness in non-malignant chronic pain.¹⁶
- ▶ Buprenorphine transdermal has been suggested to be safe and effective in patients with cancer pain and can be started in opioid-naive patients as a long-acting opioid.^{17,18}
 - ◊ Due to its long-duration of effect, it is best used in patients with stable and predictable opioid requirements.
 - ◊ Immediate-release opioid formulations may be used to treat breakthrough pain.

• Buprenorphine for chronic pain:

- ▶ Buprenorphine is a Schedule III opioid analgesic, considered a partial agonist with very high binding affinity for the mu-opioid receptor, an antagonist with high binding affinity for the delta- and kappa-opioid receptors, and an agonist with low binding affinity for the opioid receptor-like 1 receptor. It has a potential for better tolerability as well as safety advantages compared with full μ -opioid receptor agonists. Can also be used with supplemental dosing.¹⁹
- ▶ Buprenorphine use has been associated with better pain control without precipitating opioid withdrawal in patients with pain and on long term opioid therapy.²⁰

• Buprenorphine for opioid use disorder (OUD):

- ▶ It is essential to continue addressing and treating OUD while treating cancer pain.^{21,22}
- ▶ Majority of cancer hospitals do not offer treatment for OUD.²³
 - ◊ Patients with OUD may be at higher risk for poorly controlled cancer-associated pain. Further, patients with OUD may face challenges accessing clinical support and psychosocial services.²¹
 - ◊ Clinicians treating cancer pain are encouraged to collaborate and coordinate care with local experts in addiction medicine/ buprenorphine prescribers already involved in the patient's care.
 - If patient with OUD is admitted on buprenorphine, continue buprenorphine unless clinically contraindicated.
 - If patient with OUD comes to outpatient clinic on buprenorphine, ask buprenorphine prescriber to continue prescribing buprenorphine for OUD and to continue mental health/counseling/support.
 - If a local buprenorphine prescriber is not available, consult with your institution's pain, palliative care or mental health clinicians certified to prescribe buprenorphine for OUD.

[Continued](#)[References](#)**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY

Regulatory requirements for use of buprenorphine:

- Special training or DEA waiver is not required to prescribe buprenorphine (transdermal or buccal film) for pain.
 - ▶ A consultation with pain or palliative care specialist is recommended.²⁴
 - ▶ DEA waiver is not required for off-label use of buprenorphine and naloxone combination products that are approved for OUD. Adding pain as the indication when prescribing is required.
- DEA waiver is required to use buprenorphine as treatment for OUD.²⁵
 - ▶ No additional training is required by the DEA to obtain the waiver for prescribing up to 30 patients.²⁶
- The FDA strongly encourages healthcare providers complete a REMS-compliant education program when prescribing these products.

Adding an opioid (full mu-agonist) to treat pain in a patient receiving buprenorphine for OUD:

- Oncology Practice
 - ▶ Data and consensus statements are lacking.
 - ▶ In select patients with cancer pain, buprenorphine-naloxone alone may be an effective analgesic. In general, the formulations used for OUD may be used interchangeably.²⁷
 - ▶ Short-acting full opioid agonist may be added if needed for breakthrough pain after initiating buprenorphine.
- Surgical Practice
 - ▶ Patients on buprenorphine perioperatively used significantly less opioids than those whose buprenorphine was stopped.²⁸
 - ▶ Goals of perioperative pain management in patients with OUD are to:²⁹
 - ◊ Minimize the risk of OUD relapse.
 - ◊ Prioritize perioperative communication with the buprenorphine prescriber.
 - ◊ Continue the patient's home buprenorphine dose perioperatively.
 - ▶ Consult with institutional anesthesiologist, pain or palliative care providers if the patient is on buprenorphine products for OUD.
- Medical Practice
 - ▶ Continuing buprenorphine when admitted to the hospital reduces opioid requirements during the hospital stay.³⁰
 - ▶ Stopping buprenorphine for OUD during a hospitalization may lead to failure to reinstate at discharge with potential risk of OUD relapse.³⁰
 - ◊ In patients who stopped therapy, 50% relapse into OUD.³¹
 - ▶ Buprenorphine reduces morbidity and mortality in patients with OUD.³²

Pitfalls of adding buprenorphine to a full opioid agonist:

- Abrupt initiation of buprenorphine in an opioid tolerant patient receiving full agonist opioid may precipitate acute opioid withdrawal.³³ ([See page PAIN-G 11 of 18](#))

[Continued](#)

[References](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY

Buprenorphine (continued):

- Reduce the risk of opioid withdrawal by decreasing the dose of the current opioid to no more than 30 mg/day oral morphine mg equivalents (MME) (immediate-release only) before starting buprenorphine.
- Discontinue all around-the-clock and long-acting opioids when initiating buprenorphine.

Buprenorphine Buccal Film for chronic pain:³⁴

- Titrate individually in increments of up to 150 mcg every 12 hours no more frequently than every 4 days to a dose that provides adequate analgesia and minimizes adverse reactions.
- Film strengths of 600 mcg, 750 mcg, and 900 mcg are only for use following titration from lower doses. Do not exceed 900 mcg due to risk of QTc interval prolongation.

Transdermal Buprenorphine Patch for chronic pain:³⁵

- Start with 5 mcg/h patch transdermal; change every 7 days (can be started in an opioid naive patient).
- Individually titrate by 5 mcg/h to a dose that provides adequate analgesia and minimizes adverse reactions up to the maximum dose of 20 mcg/h; minimal titration interval is 72 hours.
- Transdermal patches of 7.5, 10, 15, and 20 mcg/hour are only for use in patients who are opioid-tolerant. Do not exceed 20 mcg/h due to risk of QTc prolongation.

Table 3. Dose Conversion Guidelines for Daily Oral Morphine Equivalents to Buprenorphine

- Reduce opioid dose to maximum 30 mg/day oral morphine equivalent before initiating buprenorphine at low dose and then proceeding with gradual dose titration.
- Other micro dosing initiation protocols have been described.^{36,37} Consider consultation with pain management specialist or opioid addiction specialist familiar with buprenorphine initiation.

Daily Oral Morphine Equivalents (MME) Before starting Buprenorphine Buccal Film/Patch	Buprenorphine Buccal Film	Transdermal Buprenorphine Patch
<30 mg/day including opioid naive	75 mcg daily or every 12 hours	5 mcg/h every 7 days
30-80 mg/day	150 mcg every 12 hours	Taper around-the-clock opioids for up to 7 days to no more than 30 MME/day; then, initiate TD buprenorphine 10 mcg/h at next dosing interval; may use short-acting analgesics as needed until analgesic efficacy is attained
81-89 mg/day		Consider alternative analgesic
90-160 mg	300 mcg every 12 hours	
>160 mg	Consider alternative analgesic	

[References](#)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY****Convert or Rotate From One Opioid to Another Opioid**

1. Determine the amount of current opioid(s) taken in a 24-hour period.
2. Calculate the equianalgesic dose of the new opioid. [See Table 2 on PAIN-G, 7 of 18.](#)
3. If pain was effectively controlled, and the patient is opioid tolerant, reduce the dose by 25%–50% to allow for incomplete cross-tolerance between different opioids. During the first 24 hours, titrate as needed to analgesic effect.
4. If previous dose was ineffective, may begin with 100% of equianalgesic dose.
5. Lastly, for oral opioids divide the total daily dose of new opioid needed by the number of doses per day to determine the individual dose (eg, if using short-acting opioid: 6 doses for regular PO morphine every 4 hours; if using long-acting opioid: 2 doses for extended-release morphine every 12 hours). In addition, consider as needed doses for breakthrough pain of 10% to 20% of the total daily dose.
6. Data do not support a specific transmucosal fentanyl dose equianalgesic to other opioids or between different transmucosal formulations. See package insert of specific transmucosal formulations for appropriate dosing information.
<https://www.tifremsaccess.com/TirfUI/rems/home.action>
7. Consider the impact of impaired liver and renal function (if present) on metabolism and clearance of the new opioid. [See Table 2 on PAIN-G, 7 of 18.](#)

Case Example of Converting IV Morphine to IV Hydromorphone

A patient is taking IV morphine at 8 mg/h and needs to be converted to IV hydromorphone.

1. Determine the total amount of current IV morphine in a 24-hour period for this patient
(8 mg/h x 24 hours) = 192 mg/day
2. From Table 2 on [PAIN-G, 7 of 18](#), calculate the equianalgesic dose of IV hydromorphone
10 mg IV morphine = 1.5 mg IV hydromorphone; therefore,
192 mg/day IV morphine = 28.8 mg/day IV hydromorphone = 1.2 mg/h IV hydromorphone
3. If patient was effectively controlled with IV morphine (192 mg/day), reduce the dose of hydromorphone by 25%–50%.
(28.8 mg/day reduced by 25%) = 21.6 mg/day IV hydromorphone = 0.9 mg/h IV hydromorphone
(28.8 mg/day reduced by 50%) = 14.4 mg/day IV hydromorphone = 0.6 mg/h IV hydromorphone
If dose of IV morphine was ineffective in controlling pain, may begin with 100% of equianalgesic hydromorphone dose:
(28.8 mg/day IV hydromorphone = 1.2 mg/h IV hydromorphone)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY****Convert or Rotate From Another Opioid to Transdermal Fentanyl**

1. Determine the 24-h analgesic requirement of morphine.
2. For conversion from oral morphine to transdermal fentanyl, consider ratio of 200 mg/day oral morphine = 100 mcg/h fentanyl patch.
[See Table 2 PAIN-G, 7 of 18](#) for converting other opioids to morphine equivalent with subsequent conversion to transdermal fentanyl.³⁸
3. Clinical data are unavailable to recommend specific ratio to convert from fentanyl patch to oral morphine. (Common clinical practice is to use a similar conversion ratio as when switching from oral morphine to transdermal fentanyl. Titrate with caution.)

NOTE: Due to patient variability the doses suggested by this conversion are approximate and clinical judgment must be used to titrate to the desired response.

Case Example of Converting Oral Morphine to Transdermal Fentanyl Patch

A patient is taking 30 mg of sustained-release oral morphine every 12 hours and needs to be converted to transdermal fentanyl patch.

1. Calculate the total amount of current oral morphine in a 24-hour period.
(oral morphine 30 mg x 2 = 60 mg/day oral morphine)
2. Using the conversion ratio of 200 mg/day oral morphine = 100 mcg/h fentanyl patch:
60 mg/day oral morphine is ≈ 30 mcg/h transdermal fentanyl patch.
Round down (for incomplete cross tolerance) to the closest equivalent patch, in this case 25 mcg/h.
Fentanyl patch is available in 12, 25, 50, 75, and 100 mcg/h; therefore, begin with 25 mcg/h patch.

Case Example of Converting Oral Oxymorphone to Transdermal Fentanyl Patch

A patient is taking 10 mg of sustained-release oral oxymorphone every 12 hours and needs to be converted to transdermal fentanyl patch.

1. Calculate the total amount of current oral oxymorphone in a 24-hour period
(oral oxymorphone 10 mg x 2 = 20 mg/day oral oxymorphone)
2. From Table 2 on [PAIN-G, 7 of 18](#), convert to the equianalgesic dose of oral morphine:
10 mg oral oxymorphone = 30 mg oral morphine; therefore,
(20 mg/day oral oxymorphone x 3) = 60 mg total daily dose oral morphine
3. Using the conversion of 2 mg/day oral morphine: 1 mcg/h transdermal fentanyl:
60 mg/day oral morphine is ≈ 30 mcg/h transdermal fentanyl patch.
Round down (for incomplete cross tolerance) to the closest equivalent patch, in this case 25 mcg/h.
Fentanyl patch is available in 12, 25, 50, 75, and 100 mcg/h; therefore, begin with the 25 mcg/h patch.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY****Special Notes Regarding Transdermal Fentanyl:**

- Pain should be relatively well-controlled on an opioid prior to initiating the fentanyl patch.
 - Use fentanyl patch only in patients tolerant to opioid therapy.
 - Patches are NOT recommended for unstable pain requiring frequent dose changes or dose titration.
- Fever, topical application of heat, or extreme exertion may accelerate transdermal fentanyl absorption and are contraindications for transdermal fentanyl. Avoid exposing the fentanyl transdermal system application site and surrounding area to direct external heat sources. Temperature-dependent increases in fentanyl release from the system may result in overdose and death.
- Transdermal fentanyl patch should not be punctured or cut.
- An as-needed dose of morphine or other short-acting opioid should be prescribed and will be needed, particularly during the first 8 to 24 hours.
- Once the levels have reached a steady state after at least 2 to 3 days, increase the patch dosage based on the average amount of stable daily opioid required. Continue breakthrough medication once the patch dose is stabilized.
- When converting from continuous parenteral infusion fentanyl to transdermal fentanyl, a straight 1:1 ratio³⁹ is appropriate, (ie, the number of mcg of parenteral fentanyl per hour should be approximately equal to the number of mcg of transdermal fentanyl per hour). Decrease basal rate of infusion by 50% at 6 hours after placing fentanyl patch, then stop the basal infusion 12 hours after patch placement. An alternative method is to stop the basal infusion 6 hours after patch placement.⁴⁰ In some patients, additional dose titration of the fentanyl patch may be necessary.
- The fentanyl patch analgesic duration is usually 72 hours, but patients experiencing end-of-dose failure may require fentanyl patch replacement every 48 hours.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY**

Please note: PRACTITIONERS ARE ADVISED TO CONSULT WITH A PAIN OR PALLIATIVE CARE SPECIALIST IF THEY ARE UNFAMILIAR WITH METHADONE PRESCRIBING or if individual patient considerations necessitate very rapid switching to or from methadone. Due to the unique nature of methadone with a long and variable half-life (and variability within a patient over time and variability between patients), caution should be used and frequent and careful evaluation should be performed.

Cautions Regarding Oral Methadone Use:

- The conversion ratio varies with the amount of morphine (or other opioid) a patient has been using chronically. The higher the dose of morphine, the more potent methadone is.
- To a significantly greater extent than with other opioids, methadone has been associated with many drug-drug interactions. The potential for such interactions must be investigated in each patient before initiating methadone.
- Methadone should not be titrated more frequently than every 5 to 7 days or longer, usually by 5 mg/dose or less. If more rapid titration is desired, consult with a pain or palliative care specialist.
 - ▶ 5–7 days is the time to steady state, and therefore, any significant sedation, significant improvement in pain OR immediate resolution of pain prior to the 5th day of methadone may indicate the dose is too high and the patient may be at risk of oversedation or respiratory depression by day 5–7 if the dose is not immediately adjusted.
- Methadone should be reserved for the management of chronic, not acute, pain.
- Electrocardiogram (ECG) should be considered prior to initiation and with each dose titration of methadone and should be performed prior to initiation of methadone in patients who have risk factors for increased QTc, including medications that may lengthen QTc (including some chemotherapies and biologic agents).
 - ▶ If QTc is abnormally prolonged, consider adjusting other factors, including other medications, that may impact QTc.
 - ▶ Methadone should not be used with QTc >500.
 - ▶ With QTc 450–500, consider alternate opioids and/or adjusting other factors that could prolong QTc.
 - ▶ Consider ECG when doses exceed 30–40 mg/day and again with dose of 100 mg/day. Obtain follow-up ECGs in patients with risk factors for prolonged QTc after initiation of methadone.⁴¹
[http://www.jpain.org/article/S1526-5900\(14\)00522-7/fulltext](http://www.jpain.org/article/S1526-5900(14)00522-7/fulltext)
- ▶ For patients at the end of life, ECG may not be indicated based on prognosis, goals of care, and risk/benefit ratios.

Indications for Oral Methadone Use:

- Methadone may be a viable option for pain relief in patients experiencing hyperalgesia or unrelieved pain with current opioid use.
- Consider using methadone when a long-acting opioid that can be crushed or given in a liquid solution is needed.
- Methadone may be an option when opioid rotation is indicated.
- Methadone may have superior tolerability compared to other opioids in patients experiencing chronic pain.
- Methadone is a cost-effective alternative to other opioids.
- Oral methadone has very good bio-availability.
- Methadone may be preferred over other opioids for pain control in certain circumstances (eg, in patients with a history of substance use disorder, neuropathic pain related to malignancy, renal insufficiency due to the absence of renally cleared active metabolites)

Special Notes Regarding Oral Methadone Use:

- Methadone is commercially available in 5-mg and 10-mg tablets and 1 mg/mL, 2 mg/mL, and 10 mg/mL oral solution
- Methadone tablets can be given sublingually, and may be crushed to facilitate oral, transmucosal, or enteral administration without affecting absorption.
- It may be necessary to educate patients and families about analgesic utility of methadone. Some may only be familiar with methadone use for maintenance of addiction and be unaware of its utility as a potent opioid analgesic.
- Methadone is typically given on a regular schedule with additional doses of a short-acting opioid, as needed.

[See Convert from Oral Morphine to Oral Methadone \(PAIN-G 16 of 18\) References](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY**

Please note: The conversion ratios in Table 2 should NOT be used when converting FROM methadone to other opioids. Methadone conversion can be complex and must be individualized for each patient. Assistance from a practitioner familiar with prescribing methadone or a pain/palliative care specialist is recommended.

Convert from Oral Morphine to Oral Methadone⁴²

1. Calculate the total daily oral morphine dose (or morphine-equivalent dose) the patient is using.
2. Based on the oral morphine dose, use Table 3 to determine the appropriate dose conversion ratio and calculate the oral methadone dose. These ratios take into account the potential for incomplete cross-tolerance and are based on expert consensus.
3. Divide the total daily oral methadone dose into 2–4 daily doses.

Table 4. Dose Conversion Guidelines for Total 24-hour Oral Morphine to Oral Methadone⁴³

ORAL MORPHINE	DOSE CONVERSION GUIDELINES
<60 mg	2–7.5 mg methadone per day
60–199 mg	10:1 (and patient <65 years of age)
≥200 mg	20:1 (and/or patient >65 years of age) Caution: not to exceed an initial dose of 45 mg/day

Case Example Converting Oral Morphine to Oral Methadone:

A 50-year-old patient is taking oral morphine at 30 mg every 4 hours around the clock for 3–5 days or longer, prior to conversion to methadone. (Please note that methadone should be reserved for the management of chronic, not acute, pain.)

1. Calculate the total amount of current oral morphine in a 24-hour period for this patient: (30 mg x 6) = 180 mg/day
2. From Table 3 above, calculate equianalgesic dose of oral methadone. (This conversion includes appropriate dose reduction for cross tolerance.)
For 180 mg/day of oral morphine: oral methadone, the dose conversion ratio is 10:1. (180 mg/day morphine ÷ 10) = 18 mg/day oral methadone, which is ≈ 15 mg/day oral methadone.
3. Divide the total daily oral methadone dose into 3 daily doses.
(reduced dose of 15 mg/day oral methadone ÷ 3 daily doses) = 5 mg oral methadone every 8 hours.
4. Consider continuing breakthrough dosing of short-acting opioid.

[See Special Notes Regarding Oral Methadone \(PAIN-G, 15 of 18\)](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

References

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY**
REFERENCES

- 1 Dart RC, Surratt HL, Cicero TJ, et al. Trends in opioid analgesic abuse and mortality in the United States. *N Engl J Med* 2015 Jan;372:241-248.
- 2 Anghelescu DL, Ehrentraut JH, Faughnan LG. Opioid misuse and abuse: Risk assessment and management in patients with cancer pain. *J Natl Compr Canc Netw* 2013;11:1023-1031.
- 3 Kwon JH, Tanco K, Hui D, et al. Chemical coping versus pseudoaddiction in patients with cancer pain. *Palliat Support Care* 2014;12:413-417.
- 4 American Psychiatric Association (2013). *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition*. Washington, DC. American Psychiatric Association, page 541.
- 5 Arthur JA. Urine drug testing in cancer pain management. *Oncologist* 2020;25:99-104.
- 6 Hartrick CT, Rodriguez Hernandez JR. Tapentadol for pain: a treatment evaluation. *Expert Opin Pharmacother* 2012;13:283-286.
- 7 Bell RF, Eccleston C, Kalso EA. Ketamine as an adjuvant to opioids for cancer pain. *Cochrane Database of Systematic Reviews* 2017;6:CD003351.
- 8 Ferrini R, Paice JA. How to initiate and monitor infusion lidocaine for severe and/or neuropathic pain. *J Support Oncol* 2004;2:90-94.
- 9 Kumar R, Viswanath O, Saadabadi A. Buprenorphine. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing; August 6, 2021.
- 10 Khanna IK, Pillarisetti S. Buprenorphine - an attractive opioid with underutilized potential in treatment of chronic pain. *J Pain Res*. 2015;8:859-870. Published 2015 Dec 4.
- 11 Dahan A. Opioid-induced respiratory effects: new data on buprenorphine. *Palliat Med*. 2006;20 Suppl 1:s3-s8.
- 12 Kao DP, Haigney MC, Mehler PS, Krantz MJ. Arrhythmia associated with buprenorphine and methadone reported to the Food and Drug Administration. *Addiction*. 2015;110(9):1468-1475.
- 13 Vadivelu N, Hines RL. Management of chronic pain in the elderly: focus on transdermal buprenorphine. *Clin Interv Aging*. 2008;3(3):421-430.
- 14 Böger RH. Renal impairment: a challenge for opioid treatment? The role of buprenorphine. *Palliat Med*. 2006;20 Suppl 1:s17-s23.
- 15 Ling W. Buprenorphine implant for opioid addiction. *Pain Manag*. 2012;2(4):345-350.
- 16 Davis MP, Pasternak G, Behm B. Treating Chronic Pain: An overview of Clinical Studies Centered on the Buprenorphine Options. *Drugs*. 2018;78(12):1211-1228.
- 17 Pergolizzi JV Jr, Mercadante S, Echaburu AV, et al. Euromed Communications meeting. The role of transdermal buprenorphine in the treatment of cancer pain: an expert panel consensus. *Curr Med Res Opin* 2009;25:1517-1528.
- 18 Ahn JS, Lin J, Ogawa S, et al. Transdermal buprenorphine and fentanyl patches in cancer pain: a network systematic review. *J Pain Res*. 2017;10:1963-1972.
- 19 Gudín J, Fudin J. A Narrative Pharmacological Review of Buprenorphine: A Unique Opioid for the Treatment of Chronic Pain. *Pain Ther*. 2020;9(1):41-54.
- 20 Powell VD, Rosenberg JM, Yaganti A, et al. Evaluation of Buprenorphine Rotation in Patients Receiving Long-term Opioids for Chronic Pain: A Systematic Review. *JAMA Netw Open*. 2021;4(9):e2124152.
- 21 Hser YI, Evans E, Grella C, Ling W, Anglin D. Long-term course of opioid addiction. *Harv Rev Psychiatry*. 2015;23(2):76-89.
- 22 Moryl N, Malhotra VT. A Case for Palliative Care and Addiction Specialists Collaboration and Joint Research. *JAMA Netw Open*. 2021;4(12):e2143436. Published 2021 Dec 1.
- 23 Niazi SK, Spaulding A, Brennan E, et al. Mental Health and Chemical Dependency Services at US Cancer Centers. *J Natl Compr Canc Netw*. 2021;19(7):829-838. Published 2021 Mar 4.
- 24 Heit HA, Covington E, Good PM. Dear DEA. *Pain Med*. 2004;5(3):303-308.
- 25 <https://www.samhsa.gov/>
- 26 <https://www.samhsa.gov/medication-assisted-treatment/become-buprenorphine-waivered-practitioner>
- 27 Moryl N, Filkins A, Griffio Y, et al. Successful use of buprenorphine-naloxone medication-assisted program to treat concurrent pain and opioid addiction after cancer therapy. *J Opioid Manag*. 2020;16(2):111-118.
- 28 Attaar A, Curran M, Meyenburg L, Bottner R, Johnston C, Roberts Mason K. Perioperative pain management and outcomes in patients who -discontinued or continued pre-existing buprenorphine therapy. *J Opioid Manag*. 2021;17(7):33-41.
- 29 Engle AL, Winans ARM, Demma L, Cleary J. The divided dose approach to perioperative buprenorphine -management in patients with opioid use disorder. *J Opioid Manag*. 2021;17(7):101-107.
- 30 Houchard G, Kullgren J, Sapphire M, Porter K, Abel S. Hospital Opioid Requirements Following Continuation Versus Discontinuation of Buprenorphine for Addiction - A Retrospective Cohort Study. *J Pain Palliat Care Pharmacother*. 2019;33(3-4):98-106.

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**[Continue](#)

**OPIOID PRINCIPLES, PRESCRIBING, INITIATION, TITRATION, MAINTENANCE, AND SAFETY**
REFERENCES

- ³¹ Bentzley BS, Barth KS, Back SE, Book SW. Discontinuation of buprenorphine maintenance therapy: perspectives and outcomes. *J Subst Abuse Treat*. 2015;52:48-57.
- ³² Hawk K, Hoppe J, Ketcham E, et al. Consensus Recommendations on the Treatment of Opioid Use Disorder in the Emergency Department. *Ann Emerg Med*. 2021;78(3):434-442.
- ³³ Prescribing information for buprenorphine available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2014/021306s015s019lbl.pdf. Accessed March 22, 2022.
- ³⁴ Prescribing information for buprenorphine buccal film available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2015/207932s000lbl.pdf. Accessed March 18, 2022.
- ³⁵ Prescribing information for buprenorphine transdermal system available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/021306s035lbl.pdf. Accessed March 18, 2022.
- ³⁶ Spreen LA, Dittmar EN, Quirk KC, Smith MA. Buprenorphine initiation strategies for opioid use disorder and pain management: A systematic review. *Pharmacotherapy*. 2022;42(5):411-427.
- ³⁷ Ghosh S, Klair S, Tanguay R, et al. A Review of Novel Methods to Support the Transition From Methadone and Other Full Agonist Opioids to Buprenorphine/Naloxone Sublingual in Both Community and Acute Care Settings. *Canadian J Addict*. 2019;10(4):41-50.
- ³⁸ Breitbart W, Chandler S, Egel B, et al. An alternative algorithm for dosing transdermal fentanyl for cancer-related pain. *Oncology (Williston Park)* 2000;14:695-702.
- ³⁹ Kornick CA, Santiago-Palma J, Khojainova N, et al. A safe and effective method for converting patients from intravenous to transdermal fentanyl. *Cancer* 2001;92:3056-3061.
- ⁴⁰ Samala RV, Bloise R, Davis MP. Efficacy and safety of a six-hour continuous overlap method for converting intravenous to transdermal fentanyl in cancer pain. *J Pain Symptom Manage* 2014;48:132-136.
- ⁴¹ Chou R, Cruciani RA, Fiellin DA, et al. Methadone safety: A clinical practice guideline from the American Pain Society and College on Problems of Drug Dependence, in collaboration with the Heart Rhythm Society. *J Pain* 2014;15:321-337.
- ⁴² Manfredi PL, Houde RW. Prescribing methadone, a unique analgesic. *J Support Oncol* 2003;1:216-220.
- ⁴³ McPherson ML, Walker KA, Davis MP, et al. Safe and appropriate use of methadone in hospice and palliative care: expert consensus white paper. *J Pain Symptom Manage* 2019;57:635-645.

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**MANAGEMENT OF OPIOID ADVERSE EFFECTS****Principles of Managing Opioid Adverse Effects**

- Adverse effects to opioids are common, should be anticipated, and should be managed aggressively.
- Opioid adverse effects generally improve over time, except with constipation. Maximize non-opioid and nonpharmacologic interventions to limit opioid dose and treat adverse effects. If adverse effects persist, consider opioid rotation.
- Patient and family/caregiver education is essential for successful anticipation and management of pain and opioid adverse effects.
- Information from patient and family/caregiver about adverse effects is essential for appropriate opioid dose adjustment and treatment.
- Recognize that pain is rarely treated in isolation in cancer and adverse effects also may be from other treatments or cancer itself.
- Chronic opioid therapy may depress HPA axis and cause hypogonadism in males¹ and females.
- Multisystem assessment is necessary.

Constipation

- Preventive measures
 - ▶ Educate patient and family on the need for bowel movements despite minimal food intake.
 - ▶ Set goals of treatment and explain to patient and family (eg, soft stool, ease of defecation, bowel movement every 2 days or less, adjusted per individual bowel habits).
 - ▶ Patients taking daily opioids almost always require agents for management of constipation.
 - ▶ Prophylactic medications
 - ◇ Stimulant laxative (eg, senna, 2 tablets daily; maximum 8 tablets per day of senna)
 - ◇ Polyethylene glycol 17 gm = 1 heaping tablespoon in 8 oz water PO 1–2 times daily
 - ◇ Increase dose of laxative when increasing dose of opioids.
 - ▶ Maintain adequate fluid intake.
 - ▶ While maintaining adequate dietary fiber intake is recommended, supplemental medicinal fiber such as psyllium is unlikely to control opioid-induced constipation and may worsen constipation.
 - ▶ Docusate does not provide benefit.
 - ▶ Exercise, if feasible.
- If constipation develops
 - ▶ Assess for cause and severity of constipation, including impact of other contributing medications.
 - ▶ Rule out obstruction.
 - ▶ Titrate laxatives as needed with goal of one non-forced bowel movement every 1 to 2 days.
 - ▶ Consider adjuvant analgesic to allow reduction of the opioid dose.
- If constipation persists
 - ▶ Reassess for the cause and severity of constipation, rule out bowel obstruction/impaction and hypercalcemia, and evaluate for impact of other medications potentially associated with constipation.
 - ▶ Consider adding another agent, such as magnesium hydroxide 30–60 mL daily; bisacodyl 5–15 mg PO daily, 10-mg suppository PR daily; lactulose 30–60 mL daily; sorbitol 30 mL every 2 hours x 3, then as needed; magnesium citrate 8 oz PO daily; or polyethylene glycol (17 g/8 oz water PO two times daily).
 - ▶ Oral sodium phosphate should only be used with extreme caution in patients with acute renal insufficiency.
 - ▶ Sodium phosphate, saline, or tap water enema should be used sparingly with awareness of possible electrolyte abnormalities.
 - ▶ The use of rectal suppositories and/or enemas are contraindicated in neutropenic or thrombocytopenic patients.
 - ▶ When response to laxative therapy has not been sufficient for opioid-induced constipation, consider peripherally acting mu-opioid receptor antagonists (PAMORAs) such as methylnaltrexone, naloxegol, or naldemedine (FDA approved for opioid-induced constipation).
 - ◇ Other agents including lubiprostone (FDA approved for opioid-induced constipation) could also be considered.
 - ◇ These agents should not be used in patients with known or suspected mechanical bowel obstruction, recent bowel surgery, transmural bowel metastases, or other processes affecting integrity of GI lumen due to potential increased risk of perforation.
 - ▶ For intractable chronic constipation, consider opioid rotation to transdermal fentanyl or methadone.
 - ▶ Consider neuraxial analgesics, neuroablative techniques, or other interventions to decrease pain, alleviate constipation, and/or reduce opioid dose.

¹ Rajagopal A, Vassilopoulou-Sellin R, Palmer JL, et al. Symptomatic hypogonadism in male survivors of cancer with chronic exposure to opioids. *Cancer* 2004;15:100:851-858.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MANAGEMENT OF OPIOID ADVERSE EFFECTS****Nausea (See NCCN Guidelines for Antiemesis)****Preventive measures**

- ▶ Ensure that patient is having bowel movements consistently.
- ▶ For patients with a prior history of opioid-induced nausea, prophylactic treatment with antiemetic agents (see below) is highly recommended.
- If nausea develops:
 - ▶ Assess for other causes of nausea (eg, central nervous system [CNS] pathology, chemotherapy, RT, hypercalcemia, bowel obstruction).
 - ▶ Consider prochlorperazine, 10 mg PO every 6 hours as needed; or metoclopramide, 10–15 mg PO 4 times daily as needed; or haloperidol, 0.5–1 mg PO every 6–8 hours as needed. Chronic use of any of these agents may be associated with development of tardive dyskinesia, especially in frail, elderly patients.
 - ▶ Alternative agents to consider include:
 - ◊ Serotonin antagonists (lower risk of CNS adverse effects; however, use with caution as constipation is an adverse effect)
 - Ondansetron, 4–8 mg PO 3 times daily oral tablet or orally disintegrating tablet
 - Granisetron, 2 mg PO daily
 - ◊ Olanzapine 2.5–5 mg QHS
 - ◊ Scopolamine (especially indicated for motion-related nausea; may cause or worsen constipation)
 - ◊ Dexamethasone
 - ◊ FDA-approved cannabinoids, such as dronabinol
 - ▶ If nausea persists despite as-needed regimen, administer antiemetics around the clock for 1 week, then change as needed. There may be utility in using multiple agents from different classes to maximize effect.
- Opioid-induced nausea may resolve with continued exposure; if nausea persists for more than 1 week:
 - ▶ Reassess cause and severity of nausea.
 - ▶ Consider opioid rotation.
- If nausea persists after a trial of several opioids and above measures:
 - ▶ Reassess cause and severity of nausea.
 - ▶ Consider neuraxial analgesics, neuroablative techniques, and other interventions to potentially reduce opioid dose.

Pruritus

- If pruritus develops:
 - ▶ If pruritus is associated with rash, hives, or shortness of breath, consider true allergy and reconsider selection of opioid therapy.
 - ▶ Consider antihistamines such as cetirizine, 5–10 mg PO once daily; diphenhydramine, 25–50 mg PO or IV every 6 hours; promethazine, 12.5–25 mg PO every 6 hours; or hydroxyzine, 25–50 mg every 6 hours PO or IM.
 - ▶ Assess for other causes (eg, other medications).
- If pruritus persists:
 - ▶ Consider changing to another opioid if symptomatic management has failed.
 - ▶ Consider adding to analgesic regimen: small doses of mixed agonist-antagonist, nalbuphine, 0.5–1 mg IV every 6 hours as needed.
 - ▶ Consider continuous infusion of naloxone, 0.25 mcg/kg/h and titrate up to 1 mcg/kg/h for relief of pruritus without decreasing effectiveness of the analgesic.
 - ▶ Consider ondansetron, 4–8 mg PO every 8 hours, as needed.

Delirium

- Assess for other causes of delirium (eg, infection, hypercalcemia, CNS, metastases, other psychoactive medications).
- If other possible causes of delirium are excluded, consider lowering the dose of the current opioid or consider changing the opioid.
- Consider nonopioid analgesic to allow reduction of the opioid dose.
- If delirious behavior necessitates medical intervention, consider:
 - ▶ Initial titration with haloperidol, 0.5–2 mg PO or IV every 4–6 hours
 - ▶ Olanzapine, 2.5–5 mg PO or sublingual every 6–8 hours
 - ▶ Risperidone, 0.25–0.5 mg 1–2 times per day
 - ▶ Consider quetiapine, especially in patients with Parkinson's syndrome.
 - ▶ Consider initially dosing on an as-needed basis.
 - ▶ With prolonged administration of these agents, it may be necessary to decrease dose due to long elimination half-life.
 - ▶ Antipsychotics may prolong QTc interval and ECG monitoring should be considered.
- For further information about delirium, [see NCCN Guidelines for Palliative Care](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MANAGEMENT OF OPIOID ADVERSE EFFECTS****Motor and Cognitive Impairment**

- Studies have shown that stable doses of opioids (>2 weeks) are not likely to interfere with psychomotor and cognitive function, but these functions should be monitored during analgesic administration and titration.
- Exercise caution regarding concurrent therapies and additive sedative effects, especially with benzodiazepines and gabapentin.
- Consider evaluation for driving impairment, often done through occupational therapy.

Respiratory Depression

- Sedation often precedes respiratory depression; therefore, progressive sedation should be noted and adjustments in care should be made.
- Respiratory rate <10 breaths per minute may be an early sign of respiratory depression.
- Patients with limited cardiopulmonary reserve are more susceptible.
- Hypercarbia occurs before hypoxia.
- For concerns about respiratory depression:
 - ▶ Reduce opioid dose
 - ▶ Increase interval of opioid administration
 - ▶ Assess for transdermal preparations (eg, a forgotten fentanyl patch)
 - ▶ Monitor closely
- If respiratory depression or opioid-induced sedation occur, and patient is medically stable, consider providing noninvasive respiratory support and hold additional doses of opioid until respiratory status improves.
- If patient is unstable or response is inadequate, consider naloxone administration but use reversing agents cautiously.
 - ▶ Dilute one ampule of naloxone (0.4 mg/1 mL) into 9 mL of normal saline for a total volume of 10 mL. Give 1–2 mL (0.04–0.08 mg) every 30–60 seconds until improvement in symptoms is noted.
 - ▶ Be prepared to repeat this process (the half-life of opioids is generally longer than that of the naloxone [plasma half-life is 30–80 minutes]).
 - ▶ If the patient is not responsive within 10 minutes and total naloxone dose of 1 mg, consider another reason for the change in neurologic status.
- If reversing an opioid with a long half-life or sustained-release preparation, consider naloxone infusion.
- Closely monitor for the recurrence of pain as opioid is metabolized during reversal, which may require a cautious administration of an additional opioid.

- Slowed respiration is expected in patients at end of life receiving comfort measures only. Naloxone administration may be inconsistent with goals of care, and, if so, its use should be discouraged.

Sedation

- Sedation may be assessed using a tool such as the Pasero Opioid-induced Sedation Scale (POSS): <https://pubmed.ncbi.nlm.nih.gov/19500754/>
- It is critical to recognize the difference between cancer-related fatigue and opioid-induced sedation. (See [NCCN Guidelines for Cancer-Related Fatigue](#))
- If significant or unexpected sedation develops:
 - ▶ Rule out other causes of sedation (eg, CNS pathology, other sedating medications, hypercalcemia, dehydration, infection, hypoxia)
 - ▶ If sedation is due to opioids, consider a lower dose of opioid given more frequently to decrease peak concentrations
 - ▶ Decrease the dose of opioid if pain control can be maintained at a lower dose
 - ▶ Consider opioid rotation
 - ▶ Consider nonopioid analgesic to allow reduction of the opioid dose
 - ▶ Consider the addition of caffeine, 100–200 mg PO every 6 h; or methylphenidate, 5–10 mg 1–3 times per day; or dextroamphetamine, 5–10 mg PO 1–3 times per day; or modafinil, 100–200 mg per day; or armodafinil 150–200 mg per day.
 - ◊ When using CNS stimulants for sedation, limit dosing to morning and early afternoon to avoid insomnia at night.
- If sedation persists despite several changes of opioids and the above measures:
 - ▶ Reassess cause and severity of sedation.
 - ▶ Consider neuraxial analgesics, neuroablative techniques, and other interventions to potentially reduce opioid dose.
- If the patient has had marked sleep deprivation related to poor pain control, adjustments of analgesics to improve pain control may result in “catch up” sleep lasting 2–3 days. Therefore, extreme fatigue can result in somnolence that may be difficult to differentiate from opioid-induced sedation. If related to fatigue, patients generally can be fully aroused, although this may require some effort.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PATIENT AND FAMILY/CAREGIVER EDUCATION**

To assess for patient and family/caregiver educational needs regarding pain treatment, the health care team should:

- Provide educational materials.
- Assess for literacy to ensure understanding of education.
- Assess for meaning and consequences of pain for patient and family/caregiver.
- Assess patient and family expectations for pain management, knowledge of pain, and pain treatment.^{1,2}
- Assess for meaning and understanding of the use and risks of opioid analgesics.

Messages to be conveyed to patient and family/caregiver regarding management of pain:

- Relief of pain is medically important and there is no medical benefit to suffering with pain.
- Pain can usually be well-controlled with pain medications. For persistent pain, taking an analgesic on a regular schedule will improve pain control.
- Patients with pain often have other symptoms (eg, constipation, nausea, fatigue, insomnia, depression) that need to be controlled; management of these other symptoms may facilitate control of pain.

Messages to be conveyed to patient and family/caregiver regarding opioid analgesics:

- Morphine and morphine-like medications are principal medications used to relieve severe pain.
 - If you take these medications now, they will still work later.
 - If these medications do not work, many other options are available.
 - Opioid analgesics should only be used to treat pain and not to assist with sleep, anxiety, or other mood issues.
- When working closely with health care providers these medications can be used to safely and adequately provide cancer pain relief and avoid untoward side effects.
 - For potential risk factors for misuse/abuse, [see \(PAIN-G, 4 of 13\)](#) and [see \(PAIN-G, 6 of 13\)](#) for information on naloxone.
 - Patients with a history of prescription, illicit drug, or alcohol dependence/substance abuse are at increased risk.
 - Patients with a history of opioid use/abuse may also have increased tolerance, which may require higher doses for optimal pain control ([see PAIN-L](#)).
- These medications are controlled substances and must be used with caution:
 - These medications should not be mixed with alcohol or illicit substances.
 - Potent analgesics should be taken only as prescribed and by the person for whom the medication is prescribed; advise patients not to self increase dosage or frequency; advise patients to contact health care provider if the pain management regimen is not controlling their pain.
 - Analgesics must be in a secured location, preferably in a locked box and not in a medicine cabinet.
 - Unused or unneeded medications (especially opioid analgesics) must be properly disposed of:
 - ◊ [Per the FDA](#), unless a take-back drug program is immediately available, the recommendation is to flush excess opioids down the sink or toilet.
 - ◊ Read the product-specific disposal information included with the extended-release/long-acting opioid product.
 - Provide information pertaining to local regulations regarding the operation of machinery or motor vehicles while taking potentially sedating medication and advise patient and family/caregiver accordingly and provide appropriate medical counseling.

¹ Stewart M, Brown JB, Donner A, et al. The impact of patient-centered care on outcomes. J Fam Pract 2000;49:797-804.

² Syrjala KL, Abrams JR, Polissar NL, et al. Patient training in cancer pain management using integrated print and video materials: a multisite randomized controlled trial. Pain 2008;135:175-186.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PATIENT AND FAMILY/CAREGIVER EDUCATION****Communication with the health care provider is critical for the patient and family/caregiver to assist in meeting goals of care.**

- Be certain that patient/family know how to contact physician/hospital.
- Explain that health care providers cannot discern the patient's pain level, and that describing pain is not viewed as “complaining,” but rather is an essential source of information to enable the health care provider to adjust treatment.
- Explain that health care providers want to know about any problems the patient believes the pain medications may be causing, as there are probably ways to alleviate these issues.
- Tell the patient to let the health care providers know about difficulty obtaining medication or concerns about taking medication. Explain that providers have dealt with such issues before and that they can help.
- Expect optimal management for pain and adverse effects. Inform the patient of the right to expect pain management as part of overall care.

The following must be reviewed with each patient and family/caregiver and provided in writing on a dated form:

- A list of each medication prescribed, a description of what each medication is for, and instructions on how and when to take each one
 - Plan for obtaining prescription refills, especially for potent opioids, because schedule II narcotics cannot be ordered by telephone.
- A list of potential adverse effects of these medications and what to do if they occur
 - List may be provided by clinician and/or pharmacy
- A list of all medications to be discontinued
- A list of telephone numbers to reach an appropriate health care provider and specific instructions to call regarding:
 - Any problems obtaining the prescriptions or taking the medication
 - New pain, change in pain, or pain not relieved with medication
 - Nausea and vomiting that prevents eating for 1 day
 - Problems with bowel movements, including no bowel movements for 3 days
 - Difficulty arousing the patient from sleep easily during the daytime
 - Confusion
- A plan for follow-up visits and/or phone calls, including availability of after-hours assistance
- A plan for proper storage and disposal ([See PAIN-I, 1 of 2](#))

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PROCEDURE-RELATED PAIN AND ANXIETY**

- Anticipate and offer analgesic (topical, local, and/or systemic) and anxiolytic therapy for procedures that are frequently accompanied by pain and/or anxiety. (See [PAIN-G, 2 of 13](#) for incident pain/breakthrough pain)
- Make every effort to create a calm, comfortable procedural environment.
- Events that are expected to cause discomfort to the patient such as diagnostic and therapeutic procedures (eg, wound care, IV, arterial line, central line, injection, manipulation, bone marrow aspiration, lumbar puncture, skin biopsy, bone marrow biopsy, radiation procedure), as well as transportation/change in position for patients with incident pain, merit pretreatment with an analgesic intervention.
- Providing information regarding the analgesic techniques described below prior to the procedure is ideal as it allows the patient and family/caregiver the time they may need to assimilate the information, ask questions, and learn self-management techniques to reduce anticipatory anxiety.
- Intervention may be multimodal and potentially include one or more of the following as appropriate.
 - ▶ **Analgesics**
 - ◊ Supplemental doses of analgesics should be given in anticipation of procedure-related pain (60 min prior for oral or 15 min for IV).
 - ◊ If procedure or transportation precludes continuation of IV PCA, give the prescribed IV bolus dose 10 minutes before procedure/transport and consider administering a single subcutaneous dose equivalent to 2-hour basal infusion rate.
 - ◊ Additional analgesics and/or local anesthetics should be available for further titration as needed.
 - ▶ **Anxiolytics**
 - ◊ Anxiolytics should be given preemptively when feasible. Examples include midazolam if experienced with its administration and provided onsite, or oral lorazepam or alprazolam. Oral anxiolytics should be administered at least 30 minutes before a procedure, up to an hour before.
 - ◊ Patients should be cautioned to avoid driving or operating machinery if taking an anxiolytic prior to a procedure.
 - ◊ Use caution when combining anxiolytic medications with other medications that have a sedating effect (eg, opioids).
<http://www.fda.gov/downloads/drugs/drugsafety/ucm518672.pdf>
 - ▶ **Local anesthetics such as:**
 - ◊ Topical local anesthetic creams (containing lidocaine, prilocaine, or tetracaine) applied to intact skin with sufficient time for effectiveness as per package insert.
 - ◊ Subcutaneous administration of lidocaine with a 27-gauge needle.
 - ▶ Administration of sedatives/analgesics/general anesthesia by trained personnel.
 - ▶ Integrative and nonpharmacologic interventions for relief of pain and/or anxiety ([See PAIN-D](#)).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MANAGEMENT STRATEGIES FOR SPECIFIC CANCER PAIN SYNDROMES**

Moderate to severe cancer pain is treated with opioids as indicated ([PAIN-3](#) and [PAIN-4](#)); these interventions are meant to complement opioid management. Adjuvant analgesics are used depending on the pain diagnosis, comorbidities, and potential for drug interactions. Integrative interventions should also be optimized. ([See PAIN-D](#))

- Pain from oral mucositis:
 - ▶ Gabapentin orally or in liquid preparation
 - ▶ Local anesthetic formulations/oral care protocols
 - ▶ For more information, including for GI mucositis, see
 - ◊ <https://www.ons.org/pep/mucositis>
 - ◊ [MASCC Guidelines](#)
 - ◊ [ESMO Guidelines](#)
- Bone pain without oncologic emergency:
 - ▶ NSAIDs, acetaminophen, or steroids^a
[See Non-Opioid Analgesic \(Nonsteroidal Anti-Inflammatory Drugs \[NSAIDs\] and Acetaminophen\) Prescribing \(PAIN-E\)](#)
 - ▶ Consider bone-modifying agents (eg, bisphosphonates, denosumab).
 - ▶ Diffuse bone pain: Consider hormonal therapy or chemotherapy, corticosteroids,^a and/or systemic administration of radioisotopes.
 - ▶ Local bone pain:
 - ◊ Consider local RT, nerve block (eg, rib pain), vertebral augmentation, or percutaneous ablation techniques.
 - ◊ Assess for impending fracture with plain radiographs.
 - ▶ Consider physical medicine evaluation.
[See Specialty Consultations for Improved Pain Management \(PAIN-L\)](#)
 - ▶ Consider orthopedic consultation for stabilization, if feasible.

Consider referral to a pain specialist or interventional therapist for interventional pain therapies including percutaneous ablation techniques for bone lesions. [See Interventional Strategies \(PAIN-M\)](#)
- Nerve pain
 - ▶ Nerve compression or inflammation:
 - ◊ Trial of corticosteroids^a
 - ▶ Neuropathic pain:
 - ◊ Trial of antidepressant (SNRI or TCA), [see PAIN-F](#) and/or
 - ◊ Trial of anticonvulsant, [see PAIN-F](#) and/or
 - ◊ Consider trial of topical agent, [see PAIN-F](#)
 - ◊ For refractory pain, consider referral to a pain specialist and/or the use of interventional strategies.
[See Interventional Strategies \(PAIN-M\)](#)
- Painful lesions that are likely to respond to antineoplastic therapies:
 - ▶ Consider trial of radiation, hormones, or chemotherapy.
- For severe refractory pain in the imminently dying, consider palliative sedation ([See NCCN Guidelines for Palliative Care](#)).
- Immunotherapy-related polyarthralgias ([See NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#)).

^a Due to potential impact on immunotherapies or other treatments, the use of steroids should be coordinated with the oncology care team.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**SPECIALTY CONSULTATIONS FOR IMPROVED PAIN MANAGEMENT****Major indication for referral:**

- Pain likely to be relieved or function improved through consultation delivered by a specialty service provider as suggested below. Note that the specific provider of these services may vary in different treatment settings.

Pain and palliative care specialty consultation**See NCCN Guidelines for Palliative Care****See PAIN-M**

- Consider interventional strategies (See PAIN-M)
- Management of symptoms refractory to initial treatment
- Management of sleep disturbances
- Diagnosis and treatment of underlying condition
- Consider oral or IV ketamine for pain resistant to other analgesics
- Consider methadone for pain resistant to other opioids
- Consider palliative sedation for intractable pain
- Adjustment of drugs and doses beyond the expertise of the primary team/oncologist
- Management of complicated psychosocial issues, including aberrant drug behavior
- Clarity of goals of care, especially regarding pain and medication side effects

Physical/occupational therapy, rehabilitation/mobility, integrative medicine consultation

- Physical modalities
 - Bed, bath, and walking supports
 - Positioning instruction
 - Energy conservation, pacing of activities
 - Lymphedema management
 - Massage
 - Heat and/or ice
 - TENS
 - Acupuncture or acupressure
 - Ultrasonic stimulation

Social worker consultation

- Caregiver burden and support needs
- Recommend use of community care resources

Addiction specialist consultation

- Management of aberrant drug behavior

Mental health consultation**See NCCN Guidelines for Distress Management**

- Assessment
 - Diagnostic interview: assess for depression, anxiety, psychiatric disease, and substance abuse disorder
 - Ongoing evaluation for misuse/abuse/diversion and other defined problems
- Pharmacologic management and psychotherapy
- Adaptive coping skills
 - Imagery
 - Distraction
 - Relaxation training
 - Active coping
 - Graded task assignments, setting goals, pacing, and prioritizing
- Evidence-based treatment modalities
 - Integrative medicine practitioners can be used to deliver evidence-based treatment modalities (eg, cognitive behavioral therapy [CBT], mindfulness-based stress reduction [MBSR], acceptance-based therapy, biofeedback, hypnosis)
- Education
 - Communicate regarding need to accomplish pain relief but avoid misuse/diversion
 - Provide psycho-education
 - Discuss psychosocial factors that impact pain experience and perception
- Assist in establishing treatment agreements, limit setting, single provider/pharmacy as needed

Spiritual care consultation

- Determine importance to patient and family/caregiver and current availability of support
- Manage spiritual, existential concerns

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



INTERVENTIONAL STRATEGIES

Interventional consultation¹

• **Major indications for referral:**

- ▶ Pain likely to be relieved with nerve block (eg, pancreas/upper abdomen with celiac plexus block, lower abdomen with superior hypogastric plexus block, intercostal nerve)
- ▶ Failure to achieve adequate analgesia and/or the presence of intolerable adverse effects (may be handled with intraspinal agents, blocks, spinal cord stimulation, or destructive neurosurgical procedures)
- ▶ Desire to avoid or limit systemic opioid administration

• **Commonly used interventional procedures:**

▶ **Regional infusions (requires infusion pump)**

- ◊ **Epidural:** easy to place, requires the use of an externalized catheter/pump; for infusions of opioids, local anesthetics, and clonidine; useful for acute postoperative pain; use beyond several days to a few weeks is limited by concerns for catheter displacement and infection
- ◊ **Intrathecal:** easy to internalize to implanted pump; for infusions of opioids, local anesthetics, clonidine, and ziconotide; implanted infusion pumps may be costly, refills require technical expertise
- ◊ **Regional plexus:** for infusions of local anesthetics, to anesthetize single extremity; use beyond several days to a few weeks is limited by concerns for catheter displacement and infection

▶ **Percutaneous vertebral augmentation and/or cementoplasty for bone lesions**

▶ **Neurodestructive procedures for well-localized pain syndromes (spinal analgesics are used more frequently)**

- ◊ **Head and neck:** peripheral neurolysis generally associated with sensory and/or motor deficit
- ◊ **Upper extremity:** brachial plexus neurolysis
- ◊ **Dorsal root entry zone (DREZ) lesioning**
- ◊ **Thoracic wall:** epidural or intrathecal, intercostal, or dorsal root ganglion neurolysis
- ◊ **Upper abdominal pain (visceral):** celiac plexus block, thoracic splanchnicectomy
- ◊ **Pelvic pain:** superior hypogastric plexus block
- ◊ **Rectal/perineal pain:** intrathecal neurolysis, midline myelotomy, superior hypogastric plexus block, or ganglion impar block
- ◊ **Unilateral pain syndromes:** cordotomy
- ◊ **Consider intrathecal L/S phenol block**

▶ **Neurostimulation procedures for cancer-related symptoms**

(ie, peripheral neuropathy, neuralgias, complex regional pain syndrome)

▶ **Percutaneous ablation techniques for bone lesions**

- ◊ **Specific therapies for bone pain are outside the scope of this guideline. Other resources (eg, [Filippiadis 2019](#)) may be referred to for more information**

If interventional approaches are appropriate:

- Evaluate which pain site can be relieved
- Verify that interventional technique will provide sufficient benefit
- ▶ If interventional treatment is undertaken and is successful, patient may require significant reduction in systemic opioid

If interventional approaches are not appropriate²:

- Reassess therapeutic plan

¹ Patient prognosis should be communicated to interventional pain colleagues as an important consideration when selecting interventional pain therapies.

² Infection, coagulopathy, very short life expectancy, distorted anatomy, patient unwillingness, medications that increase risk for bleeding (eg, anti-angiogenesis agents such as bevacizumab), or technical expertise is not available.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF PHARMACOGENETICS**

- Patients may respond differently to the same drug given at the same dose for the same indication often as a result of inherent differences in drug disposition due to genetic alterations that impact drug metabolism. These differences may lead to little or no analgesic response or significant adverse effects.
- Genetic factors can influence the analgesic response to opioids via pharmacokinetic (metabolic enzymes, ie, CYP P450) or pharmacodynamic (receptors and signal transduction) pathways.¹
- Pharmacogenomic testing may be considered prior to initiation or during analgesic pharmacologic treatment when concerns of toxicity or lack of analgesic response are demonstrated or suspected.
- Many commonly prescribed analgesics are metabolized via P450 (CYP) such as CYP2D6, CYP2C19, or CYP2C9.
- Opioid-mediated analgesia can be influenced by the Catechol-O-methyltransferase (*COMT*) gene and the μ -opioid receptor (*OPRM1*) A118G single-nucleotide polymorphism; however, the clinical importance of these are unclear.
- FDA-approved pharmacogenetic tests for *CYP2D6*, *CYP2C19* and *CYP2C9* are currently available; however, insurance reimbursement and availability of approved laboratories may be limited.
- Consider consulting a clinical pharmacist or clinical pharmacogenomics specialist to aid in drug selection and dose adjustments based on the interpretation and evaluation of pharmacogenomic test results.
 - ▶ **CYP2D6: Codeine, Tramadol¹**
 - ◊ Avoid codeine and tramadol in patients who are known CYP2D6 ultrarapid metabolizers (UM) due to the risk of increased toxicity. If a patient is determined to be a CYP2D6 UM, rotate to another opioid (morphine, oxycodone, or hydromorphone) and/or consider non-opioid analgesic alternatives.
 - ◊ Avoid codeine and tramadol in patients who are known poor metabolizers (PM) due to the lack of analgesic effect. If a patient is determined to be CYP2D6 PM, rotate to another opioid (morphine, oxycodone, or hydromorphone). Tramadol is not recommended as an alternative to codeine.
 - ◊ Monitor codeine and tramadol use in patients who are intermediate metabolizers (IM) for less than optimal response and offer an alternative analgesic if warranted.
 - ▶ **CYP2C19 & CYP2D6: Amitriptyline, Doxepin^{2,3}**
 - ◊ CYP2C19 PM and UM consider alternatives to doxepin and amitriptyline such as nortriptyline or desipramine
 - ◊ CYP2D6 UM consider alternatives to amitriptyline. CYP2D6 PM consider lower starting doses of amitriptyline or a 50% dose reduction.
 - ▶ **CYP2C9: Celecoxib, Meloxicam, Ibuprofen⁴**
 - ◊ CYP2C9 PM, consider alternatives to celecoxib or ibuprofen, or initiate therapy with 25%–50% of the lowest recommended starting dose (ie, 50%–75% dose reduction), and careful dose titration to effect.
 - ◊ CYP2C9 IM or PM consider alternatives to meloxicam.

¹ Crews KR, et al. Clin Pharmacol Ther 2021;110:888-896.² Hicks JK, et al. Clin Pharmacol Ther 2013;93:402-408.³ Hicks JK, et al. Clin Pharmacol Ther 2017;102:37-44.⁴ Theken KN, et al. Clin Pharmacol Ther 2020;108:191-200.**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



NCCN Categories of Evidence and Consensus

Category 1	Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 2.2022 Adult Cancer Pain

Discussion

This discussion corresponds to the NCCN Guidelines for Adult Cancer Pain. Last updated June 03, 2021

Table of Contents

OverviewMS-2

Literature Search Criteria and Guidelines Update Methodology.MS-3

Pathophysiologic Classification of Cancer Pain SyndromesMS-3

Comprehensive Pain Assessment.....MS-3

 Selecting Tools for Assessing Pain.....MS-4

 Assessing PainMS-4

Management of Adult Cancer PainMS-5

 Management of Pain Related to Oncologic EmergencyMS-6

 Management of Pain Not Related to Oncologic EmergencyMS-6

 Management of Pain CrisisMS-7

 Management of Procedure-Related Pain and AnxietyMS-8

 Subsequent Management of Cancer PainMS-9

 Ongoing CareMS-9

 Pain in Cancer Survivors.....MS-10

Pharmacologic Interventions for Cancer Pain ManagementMS-10

 Non-Opioid Analgesics.....MS-10

 Adjuvant Analgesics.....MS-11

 Opioids and Miscellaneous Analgesics.....MS-13

Opioid Prescription, Titration, and Maintenance MS-19

Opioid Adverse Effects MS-20

Opioids and Risk Evaluation and Mitigation Strategy MS-24

Management Strategies for Specific Cancer Pain Syndromes MS-25

Pharmacogenetic Considerations MS-27

Specialty Consultations MS-28

Non-Pharmacologic Interventions for Cancer Pain Management
..... MS-28

 Integrative Interventions MS-28

 Interventional Strategies..... MS-29

Summary..... MS-30

Recommended Readings MS-30

Table 1..... MS-31

Table 2..... MS-32

References MS-33



NCCN Guidelines Version 2.2022

Adult Cancer Pain

Overview

Pain is one of the most common symptoms associated with cancer. Pain is defined by the International Association for the Study of Pain as an unpleasant sensory and emotional experience associated with, or resembling that associated with, actual or potential tissue damage.¹ Cancer pain or cancer-related pain distinguishes pain experienced by patients with cancer from that experienced by patients without malignancies. A meta-analysis revealed that pain was reported in 59% of patients undergoing cancer treatment, in 64% of patients with advanced disease, and in 33% of patients after curative treatment.² In addition, this is one of the symptoms patients fear most. Unrelieved pain denies patients comfort and greatly affects their activities, motivation, interactions with family and friends, and overall quality of life.³ There is mounting evidence in oncology that quality of life and survival are linked to early and effective palliative care, including pain management.⁴⁻⁹ Although improvements have been observed, undertreatment of pain remains an issue in a significant subset of patients with cancer and this issue may be exacerbated by the inappropriate application of recommendations against the use of opioids to patients with cancer in the setting of the United States opioid epidemic.^{10,11}

Goals of pain management are to optimize pain treatment outcomes in 5 dimensions, frequently referred to as the “5 A’s” of pain management outcomes (the “4 A’s” originally proposed by Passik and Weinreb¹² were later amended to include “Affect”):

- Analgesia: optimize analgesia (pain relief)
- Activities: optimize activities of daily living (psychosocial functioning)
- Adverse effects: minimize adverse events
- Aberrant drug taking: avoid aberrant drug taking (addiction-related outcomes)
- Affect: relationship between pain and mood

The importance of relieving pain and the availability of effective therapies make it imperative that health care providers be adept at cancer pain assessment and treatment.¹³⁻¹⁵ This requires familiarity with the pathogenesis of cancer pain, pain assessment techniques, and common barriers to the delivery of appropriate analgesia. Providers should be familiar with pertinent pharmacologic, anesthetic, neurosurgical, and behavioral interventions for treating cancer pain, as well as complementary approaches such as physical/occupational therapy. Early referral to a palliative care provider may also be helpful in managing cancer pain, as well as other symptoms related to cancer and its treatment.

The most widely accepted algorithm for the treatment of cancer pain was developed by the WHO.^{16,17} It suggests that patients with pain be started on acetaminophen or a nonsteroidal anti-inflammatory drug (NSAID). If this is not sufficient, therapy should be escalated to a “weak opioid,” such as codeine, and subsequently to a “strong opioid,” such as morphine. Although this algorithm has served as an excellent teaching tool, the management of cancer pain is considerably more complex than this three-tiered “cancer pain ladder” suggests.

These NCCN Guidelines for Adult Cancer Pain are unique in several important ways. The NCCN Guidelines identify central principles for assessing and managing cancer pain in adults. First, they list general principles of pain management, followed by guiding principles for assessment, management/intervention, and reassessment. The NCCN Guidelines acknowledge the range of complex decisions faced in the management of these patients. As a result, they provide dosing guidelines for opioids, non-opioid analgesics, and adjuvant analgesics. They also provide specific suggestions for titration and rotation of opioids, escalation of opioid dosage, management of opioid adverse effects, and when and



NCCN Guidelines Version 2.2022

Adult Cancer Pain

how to proceed to other techniques/interventions for the management of cancer pain.

Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines for Adult Cancer Pain, an electronic search of the PubMed database was performed to obtain key literature in adult cancer pain, using the following search terms: “cancer pain” (title/abstract) OR “oncologic pain” (title/abstract) OR “cancer-related pain.” The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Practice Guideline, Randomized Controlled Trial, Meta-analysis, Multi-center Study, Observational Study, Systematic Reviews, and Validation Studies.

The potential relevance of the PubMed search results was examined. The data from key PubMed articles selected by the panel for review during the Guidelines update meeting as well as articles from additional sources deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting abstracts). Recommendations for which high-level evidence is lacking are based on the panel’s review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Pathophysiologic Classification of Cancer Pain Syndromes

Different types of pain occur in patients with cancer. A number of attempts have been made to classify pain according to different criteria. Pain classification includes differentiating between pain associated with tumor, pain associated with treatment, and pain unrelated to either. Acute and chronic pain should also be distinguished from each other when deciding which therapy to use. Therapeutic strategy depends on the pain pathophysiology, which is determined by patient examination and evaluation. There are two predominant mechanisms of pain pathophysiology: nociceptive and neuropathic.^{18,19}

Nociceptive pain is the result of injury to somatic and visceral structures and the resulting activation of nociceptors. Nociceptors are present in skin, viscera, muscle, and connective tissue. Nociceptive pain can further be divided into somatic pain and visceral pain.²⁰ Pain described as sharp, well localized, throbbing, and pressure-like is likely to be somatic nociceptive pain. It occurs often after surgical procedures or from bone metastasis. Visceral nociceptive pain is often described as more diffuse, aching, and cramping. It is secondary to compression, infiltration, or distension of abdominal or thoracic viscera.

Neuropathic pain results from injury to the peripheral or central nervous system (CNS). This type of pain might be described as burning, sharp, or shooting. Examples of neuropathic pain include pain due to spinal stenosis or diabetic neuropathy, or as an adverse effect of chemotherapy (eg, vincristine), radiation therapy, or following surgical injury to the nerves.

Comprehensive Pain Assessment

A comprehensive evaluation is essential to ensure proper pain management. Failure to adequately assess pain frequently leads to poor pain management. It is therefore important to find the cause of the pain



NCCN Guidelines Version 2.2022

Adult Cancer Pain

and identify optimal therapies. This algorithm begins with the premise that all patients with cancer should be screened for pain and quality of life during the initial evaluation, at each subsequent contact, and whenever new therapy is initiated. If pain is present on a screening evaluation, the pain intensity must be quantified by the patient (whenever possible). Since pain is inherently subjective, patients' self-reporting of pain is the current standard of care for assessment.

Selecting Tools for Assessing Pain

Various methods and tools exist to assess pain severity. Intensity of pain should be quantified using a numerical rating scale (ie, 0–10), visual analog scale, categorical scale, or pictorial scale (eg, The Faces Pain Rating Scale).^{21–24} Although pain is commonly assessed using numerical or categorical ratings, some patients may experience difficulty with these scales. The Faces Pain Rating Scale may be successful with patients who have difficulty with other scales, for example, children, the elderly, and patients with language or cultural differences or other communication barriers. If the patient is unable to verbally report pain, an alternative method to obtain pain rating and pain assessment must be utilized. In addition to pain intensity, the patient should be asked to describe the characteristics of his/her pain (ie, aching, burning).

The Brief Pain Inventory (BPI) assesses pain severity in patients with cancer in two important dimensions: intensity of pain and interference of pain with a patient's life.^{22,25,26} Studies suggest that pain may interfere with daily functions to a different extent in patients with cancer versus those with chronic noncancer pain.²⁷ As such, pain interference (ie, a measure of the impact of pain on daily functions) is of particular importance when assessing pain in patients with cancer. The BPI quantifies these measures using a 0 to 10 numerical scale. Based on these numerical ratings, cut-points have been established to categorize pain severity as mild, moderate, or severe for the purpose of treatment planning.^{22,25,26}

Specialized assessment tools have been developed for specific cancer pain syndromes. For example, the Oral Mucositis Assessment Scale (OMAS) developed by the Mucositis Study Group²⁸ and the Patient-Reported Oral Mucositis Symptom (PROMS) scale^{29,30} have been validated in patients with cancer who experienced mucositis as a result of chemotherapy, radiotherapy, or bone marrow transplantation. Several tools have been developed that include assessment for neuropathic pain, including the Neuropathic Pain Scale, Leeds Assessment of Neuropathic Symptoms and Signs (LANSS), and the Neuropathic Pain Questionnaire, among others.^{31,32} A validated clinical assessment tool for bone-related pain and its effect on quality of life is the Functional Assessment of Cancer Therapy – Bone Pain (FACT-BP) Quality of Life Measurement in Patients with Bone Pain.³³ The FACT-BP was developed specifically to assess cancer-related bone pain and reflected clinical change as evidenced by differences in performance status.

Assessment of both pain intensity and impact of pain on daily functions should be considered when establishing patient-specific goals for comfort and function.

An additional assessment tool that has undergone psychometric evaluation is the PROMIS pain interference (PROMIS-PI) bank; early validation studies suggest the potential utility of this approach to pain assessment as an alternative to standard-of-care assessment methods based on the BPI.³⁴ Additional studies are needed to assess the application of the PROMIS-PI for assessing cancer pain severity.

Assessing Pain

If the patient has no pain, re-screening should be performed at each subsequent visit or as requested. Identifying the presence of pain through repeated screening is essential to allow implementation of effective pain management.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

If the Pain Rating Scale score is above 0, a comprehensive pain assessment is initiated. The comprehensive pain assessment should focus on the type and quality of pain; pain history (eg, onset, duration, course); pain intensity (ie, pain experienced at rest; with movement); location; referral pattern; radiation of pain; impact of pain (ie, interference with activities such as work, sleep, and interpersonal interactions); the associated factors that exacerbate or relieve the pain; current pain management plan; patient's pain experience and response to current therapy; prior pain therapies; breakthrough or episodic pain inadequately managed with existing pain regimen; important psychosocial factors (eg, patient distress, family/caregiver and other support, psychiatric history, risk factors for undertreatment of pain¹¹); and other special issues relating to pain (eg, meaning of pain for patient and family/caregiver; cultural beliefs toward pain, pain expression, and treatment; spiritual or religious considerations and existential suffering).^{35,36} Finally, the patient's goals and expectations of pain management should be discussed, including level of comfort and function, with family/caregivers included.

In addition, a thorough physical examination and review of appropriate laboratory and imaging studies are essential for a comprehensive pain assessment. This evaluation should enable caregivers to determine if the pain is related to an underlying cause that requires specific therapy. For example, it is inappropriate to provide only opioids to a patient suffering with pain from impending spinal cord compression. Without glucocorticoids and local radiation therapy, the pain is unlikely to be well-managed, and the patient will remain at high risk for spinal cord injury.

The NCCN Panel recommends monitoring risk factors for aberrant use or diversion of pain medication, which might be identified at initiation of care using tools such as SOAPP-R (Screener and Opioid Assessment for Patients with Pain-Revised) or ORT (Opioid Risk Tool). Although specific screening tools have not been validated in the setting of cancer care, their

validated efficacy for evaluating risk in patients with non-malignant pain supports their use in this setting.³⁷ The SOAPP was developed to predict which patients, being considered for long-term opioid therapy, may exhibit aberrant medications behaviors in the future.³⁸ SOAPP-R is a revised version of the SOAPP.³⁹ Similar to the SOAPP-R, the ORT assesses the risk of aberrant behaviors when patients are prescribed opioid medication for chronic pain with a high degree of sensitivity and specificity for determining which individuals are at risk for opioid abuse.⁴⁰ SOAPP-R and ORT discriminate between high-risk and low-risk patients.⁴¹ A high-risk score on the SOAPP-R or ORT correlates with an increased likelihood of drug abuse.⁴² Randomly administered urine drug screens and periodic review of prescription drug monitoring programs (PDMPs, also known as PMPs) can also be used to monitor for aberrant use or diversion of pain medications. A study of 552 patients who were receiving opioids for cancer pain and underwent random urine drug screening found that 24% had one or more abnormal results and that random drug screening detected abnormalities earlier compared to a historical cohort that underwent targeted urine drug screening.⁴³

The endpoint of comprehensive pain assessment is to diagnose the etiology and pathophysiology (somatic, visceral, or neuropathic) of the pain. Treatment must be individualized based on clinical circumstances and patient wishes, with the goal of maximizing function and quality of life.

Management of Adult Cancer Pain

For management of cancer-related pain in adults, the algorithm distinguishes three levels of pain intensity determined by a numerical or pictorial rating scale employed as part of the comprehensive pain assessment. The three levels of pain intensity referred to in the algorithm are mild pain, moderate pain, and severe pain.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

The NCCN Panel recommends that providers consider all pain management interventions in the context of patient-specific goals for comfort and function, as well as safety. Individualized pain treatment should also take into account the etiology and characteristics of pain and the patient's clinical condition. Patients presenting with an acute, severe pain or pain crisis may be candidates for hospital admission to achieve patient-specific goals for comfort and function. It is important to separate pain related to an oncologic emergency from pain not related to an oncologic emergency.

In addition, the algorithm distinguishes pain management approaches in patients not chronically taking opioids (opioid naïve) from patients who have previously or are chronically taking opioids for cancer pain (opioid tolerant). It also distinguishes circumstances related to anticipated procedure-related pain and anxiety.

Opioid-tolerant patients are those chronically taking opioids for pain, defined by the U.S. Food and Drug Administration (FDA) as “patients who are taking at least 60 mg oral morphine per day, 25 mcg transdermal fentanyl per hour, 30 mg oral oxycodone per day, 8 mg oral hydromorphone per day, 25 mg oral oxymorphone per day, or an equianalgesic dose of another opioid for one week or longer.”^{44,45}

Therefore, patients who do not meet the above criteria of opioid tolerance, based on not having had exposure to opioid doses at least as much as those listed above for a week or more, are considered to be opioid naïve.

Management of Pain Related to Oncologic Emergency

An oncologic emergency is defined as a life-threatening event directly or indirectly related to a patient's cancer or cancer treatment. Pain related to an oncologic emergency includes pain due to bone fracture or impending fracture of weight-bearing bone; epidural or leptomeningeal metastases seen in patients with advanced cancers; pain related to infection; or

obstructed or perforated viscus. Pain associated with oncologic emergency should be treated directly while concurrently proceeding with the treatment of the underlying condition.

Management of Pain Not Related to Oncologic Emergency

For all patients experiencing pain, care providers should offer psychosocial support and begin educational activities. Psychosocial support is needed to ensure that patients encountering common barriers to appropriate pain management (eg, fear of addiction or side effects, inability to obtain opioids) or needing assistance in managing additional problems (eg, depression, rapidly declining functional status) receive appropriate aid. The patient and the family/caregiver must be educated regarding pain management and related issues.^{46,47} Patients should be reevaluated at each contact and as needed to meet their goals for comfort and function.

Although pharmacologic analgesics, including non-opioids (such as NSAIDs or acetaminophen), opioids, and adjuvant analgesics (such as antidepressants, anticonvulsants, topical agents, and corticosteroids) are the cornerstone of cancer pain management, they are not always adequate and are associated with adverse effects. Optimal use of nonpharmacologic integrative interventions (physical, cognitive modalities, and spiritual) may serve as valuable additions to pharmacologic interventions.

When deciding upon the most appropriate medication, the patient's pain diagnosis, comorbid conditions, and potential drug interactions should be considered. Addition of adjuvant analgesics for specific pain syndromes should be considered for all groups of patients. Adjuvant analgesics may be used as the main analgesics (especially for neuropathic pain), or to enhance the effects of opioid- or non-opioid (eg, NSAIDs, acetaminophen) analgesics.⁴⁸



NCCN Guidelines Version 2.2022

Adult Cancer Pain

For opioid-naïve patients (as defined above) experiencing mild pain intensity, treatment with nonopioid analgesics such as NSAIDs or acetaminophen as well as adjuvant analgesics should be considered prior to opioid analgesics unless they are contraindicated due to adverse effects or potential drug interactions. Opioid-naïve patients experiencing moderate pain should receive non-opioid and adjuvant therapies, as appropriate, with titration of short-acting opioids as needed (see section below on *Opioid Prescription, Titration, and Maintenance*). Short-acting formulations have the advantage of rapid onset of analgesic effect. The route of administration of opioid is decided (oral vs. intravenous [IV]) based on what is best suited to the patient's ongoing analgesic needs. If four or more doses of a short-acting opioid are consistently needed throughout the day, the addition of a long-acting opioid may be considered.

Opioid-tolerant patients (as defined above) who are experiencing mild pain should continue to receive non-opioid and adjuvant therapies, as appropriate. The need for opioid analgesics should be reevaluated and gradual dose reduction may be initiated, if indicated. Opioid-tolerant patients who are experiencing moderate pain should continue non-opioid and adjuvant therapies, as appropriate, with short-acting opioids, as needed. Short-acting opioids may be titrated by increasing the daily dose by 30% to 50%, or more, until pain relief is achieved. If four or more doses of a short-acting opioid are consistently needed throughout the day, the addition or increase in dose of a long-acting opioid may be considered.

In cases of acute, severe pain or pain crisis, hospital or inpatient hospice admission may be considered to achieve patient-specific goals for comfort and function (see section below on *Management of Pain Crisis*).

The use of opioid analgesics is potentially associated with substantial adverse effects. The management of common opioid-induced adverse effects should be started simultaneously with initiation of opioid therapy.

Opioid-induced bowel dysfunction should be anticipated and treated prophylactically with a stimulating laxative to increase bowel motility, as indicated.⁴⁹

Patients with chronic persistent pain managed by stable doses of short-acting opioids should be provided with round-the-clock extended-release (ER) or long-acting (LA) formulation opioids with provision of a “rescue dose” to manage breakthrough or transient exacerbations of pain. The rescue dose is usually equivalent to 10% to 20% of the total opioid daily consumption, and may be given every hour as needed during severe exacerbations of pain. Opioids with a rapid onset and short duration are preferred as rescue doses. The repeated need for numerous rescue doses per day may indicate the necessity to adjust the baseline treatment.

Management of Pain Crisis

In patients who are experiencing severe pain (or uncontrolled pain when goals of pain management and function are not met), in order to achieve adequate analgesia, an initial dose of short-acting opioid should be determined and administered. For opioid-naïve patients, this dose should be 5 to 15 mg oral or 2 to 5 mg IV morphine sulfate or equivalent. A subcutaneous route of administration can be substituted for IV; however, the time to peak effect is generally longer (~30 minutes). For opioid-tolerant patients, a rescue dose equivalent to 10% to 20% of the total opioid taken in the previous 24 hours should be given in supplement to the patient's LA (chronic) opioid dose. However, a retrospective cohort study of 216 opioid-tolerant patients with cancer who presented to the emergency department with acute pain found that while 77.4% of those taking less than 200 oral morphine equivalent received adequate rescue doses, only 3.2% of those taking greater than 400 oral morphine equivalent received adequate breakthrough medication.⁵⁰ Continuation of a patient's previous opioid could be considered or upward titration to accommodate dose requirements could be warranted.^{51,52} In patients with



NCCN Guidelines Version 2.2022

Adult Cancer Pain

risk factors such as decreased renal or hepatic function, chronic lung disease, upper airway compromise, sleep apnea, or poor performance status, the initial dosing and upward titration of opioid analgesia should be approached with caution.

Efficacy and adverse effects should be assessed approximately every 60 minutes for orally administered opioids and every 15 minutes for IV/subcutaneous opioids to determine a subsequent dose. Upon assessment, if the pain score remains unchanged or is increased, further increase in opioid rescue dose by 50% to 100% is recommended. If the pain is reduced but still inadequately controlled, the same opioid dose is repeated and reassessment is performed at 60 minutes for orally administered opioids and every 15 minutes for opioids administered by IV/subcutaneously. If pain score remains unchanged upon reassessment after 2 to 3 cycles of the opioid, in patients with moderate to severe pain, changing the route of administration from oral to IV/subcutaneous or alternate management strategies should be considered. If the pain score decreases to a level where it is adequately controlled, the current effective dose can be continued “as needed” over an initial 24 hours before proceeding to subsequent management strategies.

Management of Procedure-Related Pain and Anxiety

Procedure-related pain represents an acute short-lived experience that may be accompanied by a great deal of anxiety. Procedures reported as painful include bone marrow aspirations; wound care; lumbar puncture; skin and bone marrow biopsies; and injections into or manipulations of an IV line, arterial line, or central line. Much of the data available on procedure-related pain come from studies on pediatric patients with cancer, which are then extrapolated to adults.

Interventions to manage procedure-related pain should take into account the type of procedure, the anticipated level of pain, and other individual

characteristics of the patient, such as age and physical condition. The interventions may be multimodal and may include pharmacologic and/or nonpharmacologic approaches. Supplemental doses of analgesics should be given in anticipation of procedure-related pain; topical, local, and/or systemic formulations can be considered. Anxiolytics, such as midazolam, lorazepam, or alprazolam, are drugs used for the treatment of anxiety and its related psychological and physical symptoms. Anxiolytics should be given between 30 and 60 minutes before a procedure to manage procedure-related anxiety when feasible. Patients should be cautioned to avoid driving or operating machinery when taking an anxiolytic.

Local anesthetics can be used to manage procedure-related pain with sufficient time for effectiveness as per package inserts. Examples of local anesthetics include lidocaine, prilocaine, and bupivacaine. Physical approaches such as cutaneous warming, laser or jet injection, and ultrasound (US) may accelerate the onset of cutaneous anesthesia. Sedatives may also be used. However, deep sedation and general anesthesia must be carried out only by trained professionals. In addition, use of nonpharmacologic interventions may be valuable in managing procedure-related pain and anxiety. The major goal of nonpharmacologic interventions that include physical and cognitive modalities is to promote a sense of control, thus increasing hope and reducing helplessness that many patients with pain from cancer experience. Creating a calm, comfortable procedural environment can help achieve this.

Patients usually tolerate procedures better when they know what to expect. Therefore, patients and family members/caregivers should receive written instructions for managing pain. Pre-procedure patient education that includes procedure details and pain management strategies is essential to allow time for assimilating the information, asking questions, and learning self-management techniques to reduce anticipatory anxiety.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

Subsequent Management of Cancer Pain

The subsequent treatment is based on the patient's continued pain rating score as well as its function and evidence of appropriate use of prior treatments. Approaches for all pain intensity levels must include psychosocial support and education for patients and their families/caregivers. For all levels of pain requiring ongoing use of an opioid, opioid doses should be administered on a routine schedule with rescue doses as needed. Constipation should be routinely evaluated and managed.

If pain at any time is severe, not improved, or increased, the working diagnosis must be re-evaluated and comprehensive pain assessment must be carried out. For patients unable to tolerate dose escalation of their current opioid due to adverse effects, an alternate opioid must be considered. Addition of adjuvant analgesics should be re-evaluated to either enhance the analgesic effect of the opioids or in some cases to counter the adverse effects associated with the opioids.⁴⁹ Optimal use of nonpharmacologic integrative interventions (physical, cognitive modalities, and spiritual) may serve as valuable additions to pharmacologic interventions. Given the multifaceted nature of cancer pain, additional interventions for specific cancer pain syndromes and specialty consultation must be considered to provide adequate analgesia. If the patient is experiencing pain of moderate intensity, with inadequate pain relief on the ongoing opioid regimen, the titration of the opioid may be continued or increased. In addition, as with patients experiencing severe pain, addition of adjuvant analgesics; additional interventions for specific cancer pain syndromes; and specialty consultation must be considered.

For patients experiencing mild pain, if they have adequate analgesia but intolerable or unmanageable adverse effects, the analgesic dose may be reduced by 10% to 25% of the current opioid dose. Addition of adjuvant

analgesics may be considered. The need for opioid analgesics should be frequently reassessed and the dose reduced, if appropriate.

Ongoing Care

Although pain intensity ratings may be obtained frequently during analgesic titration, formal pain reevaluation is required at each contact to ensure that pain management therapies are successfully meeting patient-specific goals for comfort, function, and safety.

If an acceptable level of comfort and function has been achieved for the patient, and 24-hour opioid requirement is stable, the NCCN Panel recommends converting to an ER oral medication (if feasible) or other ER formulation (ie, transdermal fentanyl). The subsequent treatment is based on the patient's continued pain rating score. Rescue doses of the short-acting formulation of the same LA opioid may be provided during maintenance therapy for the management of pain in patients with cancer not relieved by ER opioids.

A regular follow-up schedule should be initiated to monitor outcomes of analgesic therapy, including adverse effects. Pain should be assessed during each outpatient contact or at least each day for inpatients depending on patient conditions, institutional standards, and regulatory requirements.

System-related barriers exist that include cost of analgesics and a lack of access to/availability of analgesics, particularly in low-income neighborhoods or for those who are economically disadvantaged. Studies have documented the inequalities that persist since those with financial burdens or minorities have less access to pain treatment.^{36,53} The NCCN Panel recommends addressing these system barriers, including recruiting assistance from social services as needed.⁵⁴⁻⁵⁷



NCCN Guidelines Version 2.2022

Adult Cancer Pain

The patients must be provided with a written follow-up pain plan, including prescribed medications. It is important to ensure that the patient has adequate access to prescribed medications, particularly as recent research shows that patients with cancer and cancer survivors are having increasing difficulty in accessing their opioid prescriptions in recent years.⁵⁸ Therefore, the patient should be encouraged to maintain communication and coordination of care with relevant providers, especially during transitions between sites of care. Collaboration with the patient's pharmacist and insurance company is helpful in achieving this. It should be clarified with the patient which clinician will be prescribing his/her ongoing pain care and confirmed that the patient/caregiver(s) know how to contact the providers and hospital. The use of a pain diary may help to facilitate communication between the patient and their providers.⁵⁹

Equally important is monitoring for the use of analgesics as prescribed, especially in patients with risk factors for or history of substance abuse, diversion, or cognitive dysfunction. Particular attention should be paid to early recognition of ineffective analgesia despite rapid escalation of opioid analgesics, which may indicate opioid misuse or abuse. Patients and the family/caregiver should be informed that opioids should only be used to treat pain and are not intended for the treatment of sleep, anxiety, or other mood issues. However, if working closely with health care providers, opioid medications can be used to safely and effectively relieve cancer-related pain.

If an acceptable level of comfort and function has not been achieved for the patients, universal screening and assessment must be carried out and additional strategies for pain relief must be considered. Other sources of distress (eg, psychological, social, spiritual) should also be recognized, documented, and treated as these may contribute to poorly controlled physical pain. See the [NCCN Guidelines for Distress Management](#) for more information.

Pain in Cancer Survivors

Chronic pain in cancer survivors may have a unique etiology and symptomatology compared with pain experienced by patients with cancer. Up to a third of post-treatment cancer survivors experience chronic pain, which can cause psychological distress and impact quality of life.^{60,61} In 2016, ASCO issued a guideline on chronic pain management in adult cancer survivors.⁶² For more information on pain in cancer survivors as well as other survivor-related issues, please see the [NCCN Guidelines for Survivorship](#).

Pharmacologic Interventions for Cancer Pain Management

Optimal management of cancer pain is often accomplished by using a combination of pharmacologic and non-pharmacologic therapies. Pharmacologic therapies may include nonopioid analgesics (such as acetaminophen or an NSAID), adjuvant analgesics (antidepressants, anticonvulsants, topical agents, and corticosteroids), and/or opioid analgesics.

Non-Opioid Analgesics

Acetaminophen

Acetaminophen has analgesic and antipyretic, but not anti-inflammatory properties.⁶³ Recent attention has been drawn towards the relative limited efficacy and significant adverse effects of acetaminophen, particularly hepatic toxicity and possibly renal impairment.^{64,65} Concerns are compounded by the inclusion of acetaminophen in a variety of prescription opioid preparations (eg, in combination with hydrocodone or codeine), as well as in a wide selection of over-the-counter products. Due to concerns about liver toxicity, the NCCN Panel Members advise that acetaminophen should be used with caution or not used at all with combination opioid-acetaminophen products to prevent excess acetaminophen dosing.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

The FDA recommends that patients be advised to limit daily acetaminophen intake to a maximum of 4 grams, and imposes a limit of 325 mg of acetaminophen per tablet, capsule, or other dosage unit in prescription products to reduce the risk of severe liver injury from acetaminophen overdosing, an adverse event that can lead to liver failure and death.⁶⁶ The FDA has issued a boxed warning to communicate the risk of severe liver injury associated with acetaminophen to health care professionals. In addition, the companies are required to add a new warning about the risk of allergic reactions, including anaphylaxis, to the label of all prescription acetaminophen-containing products.⁶⁶ Due to concerns of hepatic toxicity, the NCCN Panel suggests that providers consider limiting chronic administration of acetaminophen to 3 grams or less per day.

NSAIDs

NSAIDs produce analgesia by blocking the biosynthesis of prostaglandins, inflammatory mediators that initiate, cause, intensify, or maintain pain. History of peptic ulcer disease or gastrointestinal bleeding, advanced age (>60 years old), male gender, and concurrent corticosteroid or anticoagulant therapy should be considered before NSAID administration to prevent upper gastrointestinal tract bleeding and perforation. The risk of gastrointestinal bleeding is increased in patients with untreated *H. pylori* infection and with chronic, rather than short-term, use of NSAIDs. As prophylaxis for NSAID peptic ulceration, consider adding misoprostol or proton pump inhibitors. Well-tolerated proton pump inhibitors are recommended to reduce gastrointestinal adverse effects induced by NSAIDs. The FDA cautions that the concomitant use of an NSAID with aspirin may reduce the cardioprotective efficacy of aspirin,⁶⁷ and concomitant use of an NSAID and low-dose (or cardioprotective) aspirin may increase the risk of gastrointestinal bleeding.^{68,69} The NCCN Panel recommends avoiding concurrent use or administering these agents separately.

NSAIDs should be prescribed with caution in patients older than 60 years of age or in those having compromised fluid status, renal insufficiency, concomitant administration of other nephrotoxic drugs, and renally excreted chemotherapy in order to prevent renal toxicities. NSAID use should be coordinated with other oncologic therapies. While there is a paucity of high-quality evidence supporting the role of NSAIDs in analgesia of cancer pain,^{70,71} the addition of NSAIDs to opioids has the potential benefit of reducing the opioid dose when sedation, cognitive function, or other CNS effects of opioid analgesic therapy become burdensome.

In patients at high risk for cardiac toxicities such as those with a history of cardiovascular disease or at risk for cardiovascular disease or complications, NSAIDs should be discontinued if congestive heart failure or hypertension develop or worsen. The FDA has issued a warning that NSAID use may increase the risk of heart attack or stroke.⁷² This risk is present even with short-term use of NSAIDs and increases with higher doses.⁷³ NSAIDs taken with prescribed anticoagulants, such as warfarin or heparin, may significantly increase the risk of bleeding complications. Topical NSAIDs such as diclofenac gel or patch may be useful in this population.

The NSAID and acetaminophen prescribing guidelines are listed in the algorithm under *Non-Opioid Analgesic (NSAIDs and Acetaminophen) Prescribing*.

Adjuvant Analgesics

The term adjuvant analgesics refers to medications that are coadministered to enhance opioid analgesia and possibly reduce adverse effects of opioids by allowing the use of lower doses of opioids. These drugs can be helpful for patients whose pain is only partially responsive to opioids. Clinically, adjuvant analgesics consist of a diverse range of drug



NCCN Guidelines Version 2.2022

Adult Cancer Pain

classes, including anticonvulsants⁷⁴ (eg, gabapentin, pregabalin), antidepressants (eg, selective serotonin reuptake inhibitors [SSRIs], serotonin–norepinephrine reuptake inhibitors [SNRIs], tricyclic antidepressants [TCAs]), corticosteroids, and local anesthetics/topical agents (eg, topical lidocaine patch). Adjuvant analgesics are commonly used to help manage bone pain, neuropathic pain, and visceral pain and, if desired or indicated, to reduce the opioid requirement. They are particularly important in treating neuropathic pain (see *Management Strategies for Specific Cancer Pain Syndromes, Neuropathic Pain*).^{75,76}

Physicians should check for drug interactions when prescribing antidepressants, paying particular attention to serotonergic medications (eg, SSRIs) due to risk of serotonin syndrome. Several antidepressants are known inhibitors of hepatic drug metabolism via inhibition of cytochrome P450 enzymes, especially CYP2D6. Tamoxifen is an estrogen receptor blocker commonly used in patients with hormone receptor-positive breast cancer. Tamoxifen undergoes extensive hepatic metabolism, and inhibition of CYP2D6 decreases production of tamoxifen active metabolites, potentially limiting tamoxifen efficacy. While some clinical studies indicate increased risk of breast cancer recurrence in tamoxifen-treated patients with breast cancer also treated with SSRI antidepressants versus those receiving tamoxifen alone,⁷⁷ other studies have not shown this effect.^{78,79} If concomitant use of an SSRI is required in a patient receiving tamoxifen, use of a mild CYP2D6 inhibitor (ie, sertraline, citalopram, venlafaxine, escitalopram) may be preferred over a moderate-to-potent inhibitor ie, (paroxetine, fluoxetine, fluvoxamine, bupropion, duloxetine).⁷⁷

The most commonly employed anticonvulsant drugs for the treatment of cancer pain are gabapentin and pregabalin.⁸⁰ They have been studied primarily in noncancer neuropathy syndromes,⁸¹ although there are data supporting their use for treatment of cancer pain in conjunction with

opioids.^{82,83} Gabapentin has been reported to reduce mucositis pain in patients receiving concomitant radiotherapy and chemotherapy.⁸⁴ When compared in a prospective, randomized, open-label trial, pregabalin relieved neuropathic cancer-related pain more effectively than transdermal fentanyl.⁸⁵

Corticosteroids have long been used to relieve neuropathic pain syndromes and have also been effective for treating bone pain due to their anti-inflammatory effects as well as relieving malignant intestinal obstruction.^{48,86} A 2015 Cochrane review summarized the existing data for corticosteroid use in cancer pain.⁸⁷

Cannabinoids and Medical Marijuana/Cannabis

In the context of shifting legality, many patients with cancer are using cannabinoids or medical marijuana for treatment of cancer- or cancer treatment-related symptoms.^{88,89} To date, the FDA has approved three cannabinoids: dronabinol, nabilone, and cannabidiol (CBD).⁹⁰ Dronabinol and nabilone (both tetrahydrocannabinol [THC] or THC mimics) have been approved to treat refractory nausea and vomiting associated with cancer treatment; dronabinol has also been approved to treat anorexia and weight loss related to AIDS. CBD has been approved to treat seizures associated with rare forms of severe epilepsy. While medical marijuana has been legalized in many states, it has not been FDA-approved for any indication.⁹⁰ Furthermore, the U.S. Drug Enforcement Administration (DEA) classifies marijuana as a Schedule I substance, meaning that it has a high potential for abuse, no currently accepted medical use in treatment in the United States, and a lack of accepted safety for use under medical supervision.⁹¹ Regardless, use of medical marijuana is common among patients with cancer, with some recent studies reporting that as many as 24% to 40% of patients with cancer in the United States use marijuana.^{92,93} Therefore, providers should assess for cannabinoid/medical



NCCN Guidelines Version 2.2022

Adult Cancer Pain

marijuana use and provide education on state and federal regulations, as appropriate.

Data supporting the use of cannabinoids as adjuvant analgesics for treatment of cancer pain are extremely limited and the results from what little data exist are somewhat conflicting. While two randomized, placebo-controlled trials have shown that nabiximols (cannabis extract that contains both THC and CBD – it is not approved for use in the United States) significantly reduced cancer-related pain compared to placebo in patients with inadequate analgesia despite chronic opioid administration,^{94,95} THC extract alone did not show a significant benefit compared to placebo,⁹⁴ and another randomized study reported no significant benefit of nabiximols compared to placebo for treatment of chemotherapy-induced neuropathic pain.⁹⁶ In these studies, the most commonly reported adverse events associated with nabiximols were somnolence, fatigue, dizziness, confusion, nausea, dry mouth, and hypotension, although these were noted to be dose-dependent and generally manageable.⁹⁴⁻⁹⁶ Systematic reviews and meta-analyses of randomized controlled trials (RCTs) have also reported mixed results with some reporting no benefit of cannabinoids for cancer pain,^{97,98} and another reporting that some cannabinoids were able to reduce certain types of cancer pain (notably neuropathic pain) depending on route of administration.⁹⁹

The route of administration can also affect the safety profile of medical marijuana. A recent observational study in a state with legalized marijuana reported that while edible cannabis products accounted for only 0.32% of sales between 2014 and 2016, they accounted for 10.7% of emergency department visits during that time period.¹⁰⁰ The adverse effects that prompted the emergency department visits also differed by route of exposure, with cannabinoid hyperemesis syndrome more common for inhaled cannabis and acute psychiatric symptoms, intoxication, and

cardiovascular symptoms more common for edible cannabis.¹⁰⁰ The authors propose that the delayed onset of effect associated with the edible route may lead users to repeat the dose, potentially resulting in delayed higher plasma concentrations.

Opioids and Miscellaneous Analgesics

While starting therapy, attempts should be made to determine the underlying pain mechanism and diagnose the pain syndrome. Optimal analgesic selection will depend on the patient's pain intensity, any current analgesic therapy, and concomitant medical illness(es). An individual approach should be used to determine opioid starting dose, frequency, and titration in order to achieve a balance between pain relief and medication adverse effects.

Pure agonists (such as morphine, oxycodone, oxymorphone, and fentanyl) are the most commonly used medications in the management of cancer pain. The short half-life opioid agonists (morphine, hydromorphone, fentanyl, and oxycodone) are preferred, because they can be more easily titrated than the long half-life opioids (methadone and levorphanol).¹⁰¹ A randomized trial compared the efficacy of low-dose morphine, a “strong” opioid agonist, to “weak opioids” (ie, codeine, codeine plus acetaminophen, tramadol) for treating moderate-intensity cancer pain. Among the 240 patients with cancer enrolled in the trial, low-dose morphine had a significantly higher response rate and earlier onset of response compared with weak opioids. Opioid-related adverse effects were comparable across the two treatment groups, and overall well-being/symptom burden was rated as significantly better in the low-dose morphine arm.¹⁰²

Morphine, hydromorphone, hydrocodone, oxymorphone, and codeine should be used with caution in patients with fluctuating renal function due



NCCN Guidelines Version 2.2022

Adult Cancer Pain

to potential accumulation of renally cleared metabolites that may cause neurologic toxicity.¹⁰³⁻¹⁰⁵

Morphine

Morphine is a mu-opioid receptor agonist and weak kappa receptor agonist. Morphine is available in a wide range of formulations and routes, including oral, parenteral, and rectal delivery.¹⁰⁶ In a patient who has not been exposed to opioids in the past, morphine is generally considered the standard starting drug of choice.^{107,108} Oral administration is the preferred route. An initial oral dose of 5 to 15 mg of oral short-acting morphine sulfate or equivalent is recommended for opioid-naïve patients. Patients presenting with severe pain needing urgent relief should be treated with parenteral opioids, usually administered by the IV route or the subcutaneous (SC) route. If given parenterally, the equivalent dose is one-third of the oral dose.¹⁰⁹ An initial dose of 2 to 5 mg of IV morphine sulfate or equivalent is recommended for opioid-naïve patients. Morphine-6-glucuronide, an active metabolite of morphine, contributes to analgesia and may worsen adverse effects as it accumulates in patients with renal insufficiency.^{110,111}

Fentanyl

Fentanyl is a highly lipid-soluble mu-opioid receptor agonist that can be administered by the parenteral, spinal, transdermal, transmucosal, buccal, and intranasal routes.^{112,113} Transdermal fentanyl is not indicated for rapid opioid titration and should be recommended only after pain is adequately managed by other opioids in opioid-tolerant patients.¹¹⁴ It is usually the treatment of choice for patients who are unable to swallow, patients with poor tolerance to morphine, and patients with poor compliance. Findings from a Cochrane Database review support the efficacy of transdermal fentanyl for relieving moderate to severe cancer pain and suggest a reduction in opioid-related constipation compared with oral morphine regimens.¹¹⁵ Another meta-analysis of RCTs reported similar results,

showing similar effectiveness of cancer pain management between transdermal fentanyl and oral morphine, but lower rates of constipation, nausea, vomiting, drowsiness, and urinary retention with transdermal fentanyl.¹¹⁶ Conversion from IV fentanyl continuous infusion basal rate via patient-controlled analgesia to transdermal fentanyl can be accomplished effectively using a 1:1 conversion ratio.¹¹⁷ Transmucosal fentanyl may be considered in opioid-tolerant patients for brief episodes of incident pain not attributed to inadequate dosing of an around-the-clock opioid. Data do not support a specific transmucosal fentanyl dose equianalgesic to other opioids or between different transmucosal formulations. There are data showing that transmucosal immediate-release (IR) fentanyl is effective in treatment of breakthrough pain in patients with cancer.¹¹⁸⁻¹²⁰

Hydrocodone

Hydrocodone is a mu- and delta-opioid receptor agonist that may be approximately equipotent with oral morphine; however, its equivalence data are not substantiated.¹¹² Clinical experience suggests use as a mild, initial use opioid, but effective dose may vary. Hydrocodone is available in IR formulations mixed with acetaminophen or ibuprofen. Hydrocodone ER preparations (without added non-opioid analgesics) are available.

Codeine

Codeine is a weak mu- and delta-opioid receptor agonist with little direct analgesic effect; it is a prodrug that is hepatically metabolized to codeine-6-glucuronide, norcodeine, morphine, morphine-3-glucuronide, morphine-6-glucuronide, and normorphine.^{112,121} This process is largely through the action of the cytochrome P450 enzyme, CYP2D6. It is important to note that CYP2D6 exhibits polymorphism between various ethnic groups and between individuals. A significant portion of individuals who are poor metabolizers would obtain reduced or no analgesic effects from codeine administration.¹²² Conversely, rapid metabolizers may experience toxicity after codeine administration from more rapid morphine production.¹²²



NCCN Guidelines Version 2.2022

Adult Cancer Pain

Hydromorphone

Hydromorphone is primarily a mu-opioid receptor agonist and weak delta-opioid receptor agonist that has properties similar to morphine and is available in oral tablets, liquids, suppositories, and parenteral formulations.^{112,123} There is some evidence suggesting that the metabolite of hydromorphone may lead to opioid neurotoxicity, including myoclonus, hyperalgesia, and seizures.¹²⁴ This metabolite may be more neurotoxic than the morphine metabolite.¹²⁵ In a prospective, open-label trial of 879 patients with cancer, hydromorphone effectively reduced pain that was inadequately controlled by other analgesics.¹²⁶ Additionally, RCTs have demonstrated the clinical noninferiority of once-daily hydromorphone ER compared with twice-daily oxycodone controlled-release¹²⁷ and four-times-daily hydromorphone IR compared with four-times-daily oxycodone IR¹²⁸ for relieving moderate to severe cancer pain. A Cochrane review found evidence that hydromorphone provides similar effect on pain management as reported for oxycodone or morphine.¹²⁹

Oxycodone and Oxymorphone

Oxycodone is an opioid with agonist activity at the mu-, delta-, and kappa-opioid receptors and is available in IR and ER formulations.¹³⁰⁻¹³² Oxycodone is also available in combination with acetaminophen; therefore, the acetaminophen dose must be monitored for safe limits to avoid potential hepatic toxicity. Recent Cochrane reviews and other meta-analyses found overall evidence that oxycodone provided similar analgesic and adverse effects to morphine, concluding that these agents could be interchangeable in the front-line treatment setting for cancer-related pain.¹³³⁻¹³⁵ Studies of oxycodone/naloxone formulations showed effective analgesia with reduced opioid-induced constipation for long-term use in cancer-related pain.^{136,137}

Oxymorphone is an opioid agonist that acts primarily at the mu-opioid receptor. It is available in an IR formulation.

Methadone

Methadone is a mu-opioid receptor agonist and an antagonist at N-methyl-D-aspartate (NMDA) receptors; it is commercially available in multiple strength oral tablets or in an oral or IV solution.¹¹² Individual variations in methadone pharmacokinetics (long half-life ranging from 8 to more than 120 hours) make its usage complex in patients with cancer.¹³⁸ Due to its long half-life, high potency, and inter-individual variations in pharmacokinetics, methadone, when indicated, should be started by or in consultation with an experienced pain or palliative care specialist.

Although many recommendations for methadone rotation exist, the NCCN Panel members find the recommendations on the starting doses of methadone outlined in the Hospice and Palliative Medicine White Paper to be the easiest to implement.¹³⁹ The evidence is mixed regarding superiority of one method to initiate methadone. Some evidence supports lower adverse events with the 3-day switch method compared to the rapid conversion (stop-and-go) or ad libitum methods. In the 3-day switch method for methadone initiation, the original opioid is discontinued gradually by lowering the daily dose by 1/3 over a 3-day period. Simultaneously methadone is initiated at 1/3 of the calculated dose, and increased by 1/3 over 3 days to the calculated amount.^{140,141}

Because the starting dose may need to be titrated up, it is essential to provide the patient with access to adequate, short-acting, breakthrough pain medications during the titration period. The NCCN Guidelines recommend monitoring for drug accumulation and adverse effects, particularly over the first 4 to 7 days, and caution that a steady state may not be reached for several days to 2 weeks. Furthermore, these recommendations should not be applied when converting from methadone to morphine.

Generally, RCT data have demonstrated that appropriately titrated methadone, although harder to manage than morphine, has similar



NCCN Guidelines Version 2.2022

Adult Cancer Pain

efficacy and tolerability and has a role in treating cancer pain.¹⁴² Studies show that outpatient initiation and rotation to methadone can be successfully done in patients with cancer without serious adverse effects.¹⁴³ Retrospective studies have also reported that low-dose methadone may improve pain control when used as a coanalgesic in patients with cancer-related pain that were receiving a different, regularly scheduled opioid analgesic.^{144,145}

There is evidence suggesting that high doses of methadone (120 mg and above) may lead to QTc prolongation and torsades de pointes, which may lead to sudden cardiac death.¹⁴⁶⁻¹⁴⁸ A study conducted in patients with cancer suggests that QT interval changes exist commonly at baseline and are not changed with the addition of methadone.¹⁴⁹ The NCCN Panel supports the use of baseline and follow-up electrocardiogram (ECG) for patients treated with methadone as outlined in published recommendations and for patients with cardiac disease, or when methadone is used in patients taking other medications also known to prolong QTc (including TCAs).^{139,150} ECG monitoring should be considered within the patient's goals of care and risk/benefit ratio as discussed with the patient. The following measures may be considered to correct QTc prolongation:

- 1) Correction of hypokalemia, hypomagnesemia, or hypocalcemia
- 2) Avoidance of other drugs that can prolong QTc
- 3) Avoidance of other drugs that can inhibit the biotransformation of methadone such as CYP3A4 inhibitors

Alternate opioids are needed for patients with QTc greater than 500 msec, and are recommended for those with QTc of 450 to 500 msec, concurrently with interventions to correct any reversible causes of prolonged QTc.¹⁵⁰ The decision must be tailored to the individual clinical situation and goals of care. Good communication among the patient, family, and care providers is a critical component of the decision process.

Patients and their families may need to be educated about analgesic utility of methadone. Some may only be familiar with methadone use for maintenance of addiction and be unaware of its utility as a potent opioid analgesic. Patients and caregivers should be educated on the signs of delayed sedation and respiratory depression that may occur 4 to 7 days or longer after initiation of methadone or after titrating the dose upwards.

Levorphanol

Levorphanol is a mu-, delta-, and kappa-opioid receptor agonist. Like methadone, levorphanol also acts as an antagonist at NMDA receptors, but it has a shorter half-life and more predictable metabolism.¹⁵¹ Similar to methadone, levorphanol varies in its dosing equivalence with morphine. In a case series of 20 patients receiving palliative or hospice care, the morphine to levorphanol conversion factors were listed as 12:1 for morphine doses of less than 100 mg, 15:1 for morphine doses between 100 mg and 299 mg, 20:1 for morphine doses between 300 mg and 599 mg, and 25:1 for morphine doses greater than 600 mg.¹⁵¹ For certain populations (eg, the elderly), levorphanol may offer similar benefits to methadone but with lessened prescribing complexities and adverse effects.¹⁵² One study also demonstrated potential efficacy of levorphanol for treating neuropathic pain.¹⁵³

Miscellaneous Analgesics and Mixed Mechanism Drugs

Tramadol and Tapentadol

Tramadol and tapentadol are atypical opioids with a dual mechanism of action on opioid receptors and neurotransmitter reuptake (eg, norepinephrine, serotonin). Tramadol and tapentadol should be used with caution or avoided in patients taking other serotonergic or monoamine oxidase inhibitors (MAOI)-like medications (eg, TCAs, SSRIs, SNRIs, and MAOIs) due to risk of serotonin syndrome.¹⁵⁴

Tramadol is a weak mu-opioid receptor agonist with some norepinephrine and serotonin reuptake inhibition that is indicated for treating moderate to



NCCN Guidelines Version 2.2022

Adult Cancer Pain

moderately severe pain.¹⁵⁵ Tramadol is available as IR and ER formulations. The NCCN Panel recommends a maximum daily dose of 400 mg for IR formulations (100 mg four times a day), or 300 mg/d for ER formulations, for adults with normal hepatic and renal function. Lower doses are recommended for older adults (≥ 75 years) and those with hepatic and/or renal dysfunction to reduce the risk of seizures. Tramadol is less potent than other opioids and is considered to be approximately one tenth as potent as morphine.¹⁵⁵ One nonrandomized, observational study in patients with cancer found comparable analgesic efficacy of high-dose tramadol (ie, ≥ 300 mg/d) and low-dose morphine (ie, ≤ 60 mg/d), but observed higher rates of constipation, neuropsychological symptoms, and pruritus in patients receiving low-dose morphine.¹⁵⁶ However, in a double-blind study of patients with cancer, tramadol produced more adverse effects, including vomiting, dizziness, and weakness, than hydrocodone and codeine.¹⁵⁷ A Cochrane review of tramadol (with or without acetaminophen) concluded that limited evidence supports the use of tramadol for treatment of cancer pain and that tramadol is likely not as effective as morphine in this setting.¹⁵⁸

Tapentadol is an opioid that binds to the μ opioid receptor and inhibits norepinephrine reuptake.^{159,160} It is available as ER and IR formulations and is used for treatment of moderate to severe pain as well as for neuropathic pain. Typical doses start at 50 to 100 mg orally every 4 hours as needed, with a maximal daily dose of 500 mg per day (if using the ER) or 600 mg per day (if using the IR only) due to a lack of published data regarding higher doses. Lower doses are recommended for patients with moderate hepatic impairment, and tapentadol should be avoided in patients with severe hepatic or renal impairment. In comparative phase II–III studies, the efficacy and safety of tapentadol have been demonstrated compared with placebo and oxycodone for non-cancer pain.¹⁶¹⁻¹⁶³ Data on tapentadol for treating non-cancer pain have also suggested that it may have a lower incidence of gastrointestinal adverse effects than

oxycodone.¹⁶¹ Limited data suggest that there may be a role for tapentadol in the management of cancer pain,^{164,165} but further clinical trials are needed.

Buprenorphine

Buprenorphine, a partial μ -agonist, has been approved for chronic pain in opioid-naïve or opioid-tolerant patients. Although RCT data on buprenorphine for treating cancer pain are somewhat limited, several case series, prospective uncontrolled studies, and a few randomized trials support its use in cancer-related pain.¹⁶⁶⁻¹⁷⁰ Therefore, transdermal buprenorphine may be used at a dose of 5 mcg/h in opioid-naïve patients requiring initiation of LA opioid therapy. In some instances, transmucosal buprenorphine may be more appropriate given a wider range of available doses, a higher maximum dose, and a lower likelihood of causing skin reactions compared to transdermal buprenorphine.

Based on its pharmacokinetics, buprenorphine may be especially appropriate for treating cancer pain in patients with renal impairment.¹⁶⁹ Studies of buprenorphine suggest that, being a partial μ -receptor agonist, it exhibits a ceiling to analgesic efficacy and may precipitate withdrawal symptoms if administered to individuals currently taking a high-dose opioid.¹⁷¹ While transdermal buprenorphine may have some advantages over methadone in the context of cancer treatments that prolong QT, FDA guidelines recommend limiting dose to a maximum of 20 mcg/h due to concern for QT prolongation. Because the dose conversion from other opioids to buprenorphine can be complex, the NCCN Panel suggests that providers consider a pain specialty consultation for complex cases.

Ketamine

Ketamine is a non-competitive N-methyl D-aspartate receptor antagonist that blocks glutamate. Low (sub-anesthetic) doses produce analgesia and may limit central sensitization, hyperalgesia, and opioid tolerance. There



are only limited data regarding the use of ketamine as an adjuvant to opioids for management of cancer pain.¹⁷² A double-blind, randomized, placebo-controlled trial found no significant difference between the outcomes of patients treated for cancer pain with ketamine versus placebo.¹⁷³ However, a subsequent systematic review of the evidence on ketamine for treating cancer-related pain concluded that the data, although limited, did suggest modest analgesic potential for ketamine.¹⁷⁴ There are also some data suggesting that ketamine may improve mood in individuals with depressive disorders.¹⁷⁵⁻¹⁷⁷

Lidocaine

While it is most often used as a local analgesic, lidocaine may also be administered intravenously in patients with refractory cancer pain. Although data supporting the use of IV lidocaine for treatment of cancer pain are limited, there are case reports and smaller studies that support its use for opioid-refractory cancer pain or postsurgical pain.¹⁷⁸⁻¹⁸¹ One phase 2, randomized, double-blind crossover study of 50 patients with opioid-refractory cancer pain found that pain relief was better with IV lidocaine compared to placebo ($P < .001$). Additionally, more patients were able to decrease their analgesic requirements following administration of IV lidocaine than placebo ($P = .0012$). Side effects including tinnitus, perioral numbness, sedation, lightheadedness, and headache were self-limiting and did not require intervention except for discontinuation of the lidocaine infusion in one patient.¹⁷⁸ A meta-analysis of pooled data from RCTs demonstrated a significant reduction in cancer pain with lidocaine infusion compared to placebo in 60 patients.¹⁸² IV lidocaine may be started as a bolus infusion of 1 to 3 mg/kg over 20 to 30 minutes. If this bolus is tolerated and effective at reducing pain, a continuous infusion of IV lidocaine may be started at 0.5 to 2 mg/kg/h (maximum 100 mg/h), using the lowest dose that controls the patient's pain.¹⁸⁰ Some reports suggest that IV lidocaine may be especially useful for cancer-related neuropathic pain.¹⁷⁹⁻¹⁸¹

Selecting a Route of Administration for Opioid Analgesics and Mixed Mechanism Drugs

The least invasive, easiest, and safest route of opioid administration should be provided to ensure adequate analgesia.

Oral is the preferred route of administration for chronic opioid therapy.^{51,183,184} The oral route should be considered first in patients who can take oral medications unless a rapid onset of analgesia is required or the patient experiences adverse effects associated with the oral administration. Continuous parenteral infusion, IV or SC, is recommended for patients who cannot swallow or absorb opioids enterally. Opioids, given parenterally, may produce fast and effective plasma concentrations in comparison with oral or transdermal opioids. IV route is considered for faster analgesia because of the short lag-time between injection and effect (peak 15 minutes) in comparison with oral dosing (peak 60 minutes).¹⁸⁵ The SC route has a slower onset and lower peak (30 minutes) effect when compared with IV route.

Analgesic Agents That Are Not Recommended

The following agents are not recommended for patients with cancer: 1) mixed agonist-antagonists (eg, butorphanol, pentazocine); 2) meperidine; and 3) placebos. Mixed agonist-antagonists should not be used in combination with opioid agonist drugs for cancer pain management. Converting from an agonist to an agonist-antagonist could precipitate the abstinence syndrome (a withdrawal crisis) if given to a patient who is physically dependent on a pure opioid agonist. Meperidine is contraindicated for chronic pain, especially in patients with impaired renal function or dehydration, because accumulation of metabolites that are cleared renally may result in neurotoxicity (seizures) or cardiac arrhythmias.¹⁸³ Use of placebo in the treatment of pain is unethical.



Opioid Prescription, Titration, and Maintenance

The appropriate dose of opioid is based on the patient's pain intensity and goals, while limiting undesirable and unmanageable adverse drug effects.

The physicians should be aware of potential drug-drug and drug-disease interactions while determining the treatment plan. For a summary of common drug-drug interactions between chemotherapeutics, analgesics, and other commonly prescribed medications, see Table 1. The patient's goals and quality of life should also be considered when modifying the treatment plan.

The following methods of ongoing analgesic administration are widely used in clinical practice: “around the clock,” “as needed,” and “patient-controlled analgesia.” For most patients, LA dosing should be used for continuous pain relief. Additional doses of opioid may be required for pain not relieved by a regular schedule of LA (eg, ER) opioid.

The NCCN Panel recommends considering opioid rotation if pain is inadequately managed despite adequate dose titration, or if persistent adverse effects from current therapy occur. Other indications for switching to a different opioid include a change in the patient's condition (dysphagia, NPO [*nil per os*] status, or initiation of tube feeding), and out-of-pocket costs and limitations based on insurance formularies.

For patients who have intermittent pain with pain-free intervals, IR opioids can be administered on an “as needed” basis, with the exception of methadone due to its long duration of effect. The “as needed” method is also used when rapid dose titration is required. The patient-controlled analgesia technique allows a patient to control a device that delivers a bolus of analgesic “on demand” (according to, and limited by, parameters set by a physician).¹⁸⁶ However, if the patient persistently requires doses of “as-needed” opioids, or if the “around-the-clock” opioid regimen fails to

relieve pain at peak effect or at end of dose, increased dose of the regularly scheduled opioid should be considered.

Breakthrough pain is defined as pain that fails to be adequately managed or “breaks through” a regimen of regularly scheduled opioid and may be further categorized as:

- incident pain that is associated with specific activities or events (eg, physical therapy, exercise, or routine procedures that may induce pain), potentially managed with “rescue doses” of short-acting opioid given in anticipation of those events;
- end-of-dose failure pain that recurs toward the end of dosing interval for regularly scheduled opioid, potentially managed by increasing the dose or frequency of regularly scheduled opioid; or
- persistent pain that is routinely inadequately managed by existing regularly scheduled opioid, potentially managed by adjusting dose of regularly scheduled opioid.

Breakthrough pain is commonly reported among patients with cancer. In a survey of 1000 oncology patients, 44% reported incident pain, 41.5% reported spontaneous pain, and 14.5% reported both incident-related and spontaneous breakthrough pain.¹⁸⁷ Although the literature on useful therapies for breakthrough cancer pain is relatively small, multiple RCTs suggest that buccal, sublingual, or oral/nasal transmucosal formulations of fentanyl are effective options for managing episodic breakthrough pain.¹⁸⁸⁻¹⁹¹

Initiating Short-Acting Opioids in Opioid-Naïve Patients

The route of administration of an opioid (oral or IV) must be selected based on the patient's needs. The NCCN Guidelines for Adult Cancer Pain management provide guidance for initiating short-acting opioids in opioid-naïve patients.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

For opioid-naïve patients experiencing moderate/severe pain, or mild pain when non-opioid therapies are contraindicated, an initial dose of 5 to 7.5 mg of oral morphine sulfate every 3 to 4 hours as needed, or equivalent, is recommended. The algorithm also recommends starting doses for oxycodone immediate release, hydrocodone, and hydromorphone in opioid-naïve patients. Pain should be reassessed regularly and the dose titrated further as needed. If four or more doses of a short-acting opioid are consistently needed each day, the addition of a long-acting opioid may be considered based on the total daily dose.

Opioid Dose Reduction

The NCCN Panel recommends monitoring patients for situations that may warrant opioid dose reduction. Scenarios where opioid dose reduction may be considered include the patient rarely or never needing breakthrough analgesics, completion of an acute pain event, response to cancer-directed therapies, or improvement of pain control through use of non-opioid or interventional pain management therapies. In these situations, the dose of opioid may be reduced by 10% to 20% after which the adequacy of pain control may be re-evaluated and further dose reductions may be considered if appropriate. Opioid dose reduction may also be considered when the pain is mild and the patient is experiencing unmanageable opioid-related adverse effects and/or rapid clinical deterioration. For more information on tapering opioids, see *Principles of Opioid Dose Reduction* in the algorithm and the [VA/DoD Clinical Practice Guideline for Opioid Therapy for Chronic Pain](#).¹⁹²

Preventing Opioid Misuse and Abuse

The NCCN Panel recommends monitoring for aberrant medication drug-related behaviors over the course of treatment using tools such as COMM (Current Opioid Misuse Measure). The COMM tool helps clinicians identify whether a patient, currently on long-term opioid therapy, is exhibiting aberrant behaviors associated with misuse of opioid medications.^{193,194} It examines concurrent misuse; in contrast, SOAPP-R or ORT are helpful in

predicting which patients being considered for long-term opioid therapy may exhibit aberrant medications behaviors in the future. Potential risk factors for abuse/misuse of prescribed analgesics include the following patient characteristics:³⁷

- History of prescription, illicit drug, or alcohol dependence or misuse prior to cancer diagnosis/treatment
- History of binge drinking or peers who binge drink
- Family history of substance abuse
- History of psychiatric disorder including anxiety, depression, attention-deficit hyperactivity disorder, post-traumatic stress disorder, bipolar disorder, or schizophrenia
- History of sexual abuse victimization
- Young age (younger than 45 years of age)
- History of legal problems or incarceration

If signs of aberrant opioid use are present, providers should consider limiting or restricting use to avoid risk of diversion. Patients who are actively receiving treatment for addiction should be encouraged to continue with therapy and care should be coordinated with their addiction specialist. See additional recommendations in *Strategies to Maintain Patient Safety and Minimize the Risk of Opioid Misuse and Abuse During Chronic Opioid Use* in the algorithm.

Opioid Adverse Effects

A number of adverse effects are associated with the use of opioid analgesics. Constipation, nausea and vomiting, pruritus, delirium, respiratory depression, motor and cognitive impairment, and sedation are fairly common, especially when multiple agents are used.¹⁹⁵⁻²⁰⁰ Chronic opioid therapy may depress the hypothalamic-pituitary axis and cause hypogonadism.²⁰¹ Each adverse effect requires a careful assessment and treatment strategy. Management of opioid-induced adverse effects is integral to opioid pain management.^{195,202-210}



NCCN Guidelines Version 2.2022 Adult Cancer Pain

The details of prophylactic regimens and other measures to prevent opioid-induced adverse effects are provided in *Management of Opioid Adverse Effects* in the algorithm.

Constipation

Constipation can almost always be anticipated with opioid treatment and patients do not develop tolerance to constipation; therefore, administration of a prophylactic bowel regimen is recommended for nearly all patients taking opioids. However, there is limited evidence on which to base the selection of the most appropriate prophylactic bowel regimen. One study showed that addition of the stool softener, docusate, to the laxative, sennosides, was less effective than administering sennosides alone.²¹¹ More recently, an RCT in hospice patients showed that there was no benefit in adding docusate to sennosides compared to sennosides alone.²¹² Therefore, for prophylaxis, the NCCN Guidelines for Adult Cancer Pain Panel Members recommend a stimulant laxative or a heaping tablespoon (17 g) of polyethylene glycol (PEG) with 8 oz of water one to two times daily along with maintaining adequate fluid intake. Based on the available literature, docusate has not shown benefit and is therefore not recommended. While maintaining adequate dietary fiber intake is recommended, supplemental medicinal fiber, such as psyllium, is ineffective and may worsen constipation.

Once constipation develops, the cause and severity of constipation must be assessed to rule out obstruction. Laxatives may be titrated as needed with the goal of achieving one non-forced bowel movement every 1 to 2 days. Adjuvant analgesics may be considered to allow reduction of the opioid dose.

If constipation persists, the cause and severity of constipation must be assessed again to rule out bowel obstruction and hypercalcemia. Providers should assess other medications with the potential to cause constipation. Adding stimulant laxatives, such as magnesium-based

products, bisacodyl (available in tablets or suppositories), or osmotic laxatives (such as sorbitol, lactulose, and PEG) may be helpful. Opioid rotation to fentanyl or methadone may be considered. Enema with sodium phosphate, saline, or tap water may be helpful as it dilates the bowel, stimulates peristalsis, and lubricates the stool to encourage a bowel movement. However, these types of enemas should be used sparingly with awareness of possible electrolyte abnormalities. The use of rectal suppositories or enemas should be avoided in patients with neutropenia or thrombocytopenia. Additionally, oral laxatives or enemas that contain sodium phosphate should be limited to a maximum dose of once daily in patients at risk for renal dysfunction; optimally, alternative agents can be employed.

When response to laxative therapy has not been sufficient, peripherally acting mu-opioid receptor antagonists (PAMORAs) such as oral methylnaltrexone,²¹³⁻²¹⁸ naloxegol,²¹⁹ or naldemedine,²²⁰ opioid antagonists that work on receptors in the gastrointestinal system can be used as a rescue when constipation is clearly related to opioid therapy²²¹ (methylnaltrexone is FDA approved for opioid-induced constipation in adults with advanced illness who are receiving palliative care; naloxegol and naldemedine are FDA approved for opioid-induced constipation in adults with chronic non-cancer pain, including those with chronic pain related to prior cancer or its treatment). Other second-line agents may be considered, including lubiprostone which is FDA approved for opioid-induced constipation in adults with non-cancer pain including those with chronic pain related to prior cancer or its treatment.^{222,223} These agents should not be used in patients with known or suspected mechanical bowel obstruction, recent bowel surgery, transmural bowel metastases, or other processes affecting integrity of the GI lumen due to the potential increased risk of perforation.²²⁴ Guidelines from the American Gastroenterological Association Institute for the medical management of opioid-induced



constipation may be referenced for more information and recommendations on use of these agents.²²⁵

Neuraxial analgesics, neuroablative techniques, or other interventions to decrease pain and/or reduce systemic opioid dose may also be considered to reduce opioid-related adverse effects.

Nausea and Vomiting

For patients with a prior history of opioid-induced nausea, prophylactic treatment with antiemetic agents is highly recommended. If nausea develops, other causes of nausea (eg, constipation, CNS pathology, chemotherapy, radiation therapy, hypercalcemia, bowel obstruction) must be assessed. Effective agents that may be considered include phenothiazines such as prochlorperazine or thiethylperazine or dopamine receptor antagonists such as metoclopramide or haloperidol.

If nausea persists despite an as-needed regimen, administer antiemetics around the clock for 1 week and then change dosing as needed. When managing opioid-induced persistent nausea, instead of replacing one antiemetic with another, adding therapies that target different mechanisms of action resulting in a synergistic effect may be helpful. Adding serotonin receptor antagonists such as granisetron or ondansetron may be helpful and have a lower rate of CNS effects. Alternative agents such as scopolamine or olanzapine may also be considered for management of nausea. Olanzapine may be especially helpful for patients with bowel obstruction.^{226,227} Corticosteroids, such as dexamethasone, can also be quite beneficial for reducing opioid-induced nausea and vomiting, and in particular have been found to be effective in combination with metoclopramide and ondansetron.²²⁸

If nausea persists for longer than a week, the cause of nausea needs to be reassessed and opioid rotation must be considered.²²⁹ If opioid rotation and the above measures have been tried and nausea still persists,

neuraxial analgesics, neuroablative techniques, and other interventions could be performed to potentially reduce the opioid dose. Cannabinoids that have been FDA-approved for chemotherapy-induced nausea and vomiting (eg, dronabinol, nabilone) may also be considered in this situation.²³⁰⁻²³³ It should be noted that in the context of shifting legality, many patients with cancer are using medical cannabis for treatment of nausea and other cancer- or cancer treatment-related symptoms.^{88,89}

While medical cannabis has been legalized in many states, it has not been FDA-approved.⁸⁹ Education on state and federal regulations for medical cannabis should be provided (see *Adjuvant Analgesics* and *Cannabinoids and Medical Marijuana/Cannabis* for more information).

Pruritus

Pruritus or itchiness is a particularly common and distressing complaint. Pruritus occurs in 10% to 50% of patients receiving opioids. Even in the presence of attentive skin care, opioids can produce recalcitrant pruritus. If pruritus develops, other causes of pruritus such as use of any other medication must first be assessed. Pruritus is more likely to occur early in the course of treatment. If it is persistent despite attempted symptom management, consider changing to another opioid. Careful titration of mixed opioid agonist-antagonists (eg, nalbuphine) or μ -opioid receptor antagonists (eg, naloxone) may help reduce opioid-induced adverse effects while maintaining analgesic efficacy. The μ -receptor antagonists (eg, naloxone) are also used to reverse the effects of opioid-induced adverse effects,²³⁴ and careful dose titration can produce relief without reversing analgesic efficacy. A serotonin antagonist such as ondansetron may also be considered. Antihistamines such as cetirizine (non-sedating), diphenhydramine (sedating), or promethazine (sedating) may be beneficial. Hydroxyzine, administered by mouth or intramuscular injection, may also be useful.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

Delirium

Delirium is a pathophysiologic condition characterized by altered consciousness and inattention, cognitive dysfunction, and disturbed psychomotor behavior. Delirium may be prevented or decreased with various non-pharmacologic interventions or, when delirium is severe and hyperactive, may be managed with a neuroleptic drug such as haloperidol, olanzapine, or risperidone on an as-needed basis or by switching to another opioid.²³⁵⁻²³⁸ Studies have shown that stable doses of opioids (>2 weeks) are not likely to interfere with psychomotor and cognitive function, but these functions should be monitored during analgesic administration and titration.²³⁹ Quetiapine may be useful, especially in patients with Parkinson's syndrome.²⁴⁰ Patients taking opioids may be screened for driving impairment, if indicated. Driving fitness screens are often performed through occupational therapy.

Sedation

It is critical to recognize the difference between cancer-related fatigue and opioid-induced sedation, as some techniques to manage sedation may not work for fatigue. The Pasero Opioid-induced Sedation Scale (POSS) has been validated and is recommended for monitoring of opioid-induced sedation, particularly for inpatient care and when increasing the dose of opioids.^{241,242} For more information on managing cancer-related fatigue, see the [NCCN Guidelines for Cancer-Related Fatigue](#). Sedation may hinder the achievement of dose titration of opioids to levels that provide adequate analgesia.⁴⁹ If opioid-induced sedation develops, it may be managed by administration of psychostimulants such as methylphenidate, dextroamphetamine, modafinil, or armodafinil or by adding caffeine. When using CNS stimulants for sedation, the dosing should be limited to morning and early afternoon to avoid insomnia at night. Sedation often precedes respiratory depression; therefore, progressive sedation should be noted and adjustments in care should be made.

Use caution when combining opioid medications with other medications that have a sedating effect (eg, benzodiazepines). The FDA has issued a black box warning about possible serious effects from this combination, including slowed or difficult breathing and death.²⁴³

Respiratory Depression

Respiratory depression is another adverse effect that is a concern for both physicians and patients. Physicians should be aware that patients with limited cardiopulmonary reserve are more susceptible and hypercarbia occurs before hypoxia. Initial steps that may be taken when there is a concern about respiratory depression include reduction of the opioid dose, increasing the interval of opioid administration, confirming that the patient does not have any forgotten transdermal opioid patches, and close monitoring. If the patient is medically stable, noninvasive respiratory support may be considered and/or additional doses of opioid analgesic may be temporarily halted until the patient's respiratory status improves.

In cases where the patient is unstable or response to the above interventions is inadequate, naloxone remains a useful antidote for the reversal of opioid-induced respiratory and CNS depression, but should be administered cautiously so as not to precipitate acute opioid withdrawal syndrome in the opioid-tolerant patient. Abrupt reversal of opioid depression in opioid-tolerant patients may result in nausea, vomiting, sweating, tachycardia, increased blood pressure, tremulousness, and seizures. Pulmonary edema, cardiac arrhythmias, and cardiac arrest have also been associated with naloxone administration.²⁴⁴ At end-of-life in patients receiving comfort measures only, slowed respiration is expected. Naloxone administration may be inconsistent with goals of care in these patients.

Naloxone may be made available to caregivers of patients receiving opioid analgesics to administer in the event of respiratory depression and sedation. While there are no RCTs, the results of a nonrandomized



NCCN Guidelines Version 2.2022

Adult Cancer Pain

intervention study showed that patients receiving long-term opioid analgesia who were co-prescribed naloxone had fewer opioid-related emergency department visits compared to those who were not prescribed naloxone.²⁴⁵ Providers should become familiar with state regulations regarding the prescription of naloxone. The availability of needle-free naloxone preparations (eg, nasal spray) may facilitate use of naloxone in the outpatient setting. Importantly, caregivers who are provided naloxone must be educated in the proper indications and usage to prevent inappropriate administration. Patients and caregivers should be instructed to call emergency services (911) if naloxone is administered. Naloxone may be available without a prescription in some localities.

Opioid Rotation

No single opioid is optimal for all patients.²⁴⁶ If opioid adverse effects are significant, an improved balance between analgesia and adverse effects might be achieved by changing to an alternative opioid. This approach is known as opioid rotation.^{195,247,248} A post hoc analysis of 498 patients from a randomized phase IV clinical trial found that the opioid was switched in 79 patients, with 51.45% of switches resulting in better pain control and 43.5% resulting in better control of opioid side effects.²⁴⁹ Establishing equianalgesic dosing can be challenging; studies have sought to establish safe conversion ratios and methods.^{141,250-254} It is important to consider relative effectiveness when switching between oral and parenteral routes to avoid subsequent overdosing or underdosing. Known equianalgesic dose ratios, opioid titration and maintenance, and clinical examples of converting from one opioid to another are listed in *Opioid Principles, Prescribing, Titration, Maintenance, and Safety* in the algorithm.

Opioids and Risk Evaluation and Mitigation Strategy

While opioids are the principal analgesics for management of moderate to severe pain in the context of a cancer diagnosis, they pose risks to patients and society. The abuse of opioids is an increasing concern. In

2018, there were 67,367 drug overdose deaths in the United States, including 46,802 drug overdose deaths involving opioid analgesics.²⁵⁵ Drug poisoning remains the number one cause of injury-related death in the United States.²⁵⁶ While it is important to ensure that opioids continue to be prescribed for patients for whom they are appropriate, it is also essential to ensure that these drugs are prescribed carefully. To reduce addiction, misuse, abuse, overdose, and death the FDA has established Risk Evaluation and Mitigation Strategy (REMS) programs for opioid products.²⁵⁷ The principal recommendations of opioid REMS programs are educating the provider, patient, and family/caregiver.

The highlights of provider responsibilities included in the REMS are:

- Establishing patient-specific goals of opioid analgesic therapy and regularly evaluating therapeutic opioid response to guide further therapy.
- Evaluating each patient for risk factors associated with opioid misuse or abuse.
- Educating each patient on safe use, storage, and disposal of opioid.
- Routinely monitoring patients for opioid misuse, abuse, or diversion.

On September 18, 2018, the FDA approved the Opioid Analgesic REMS program, which covers all opioid analgesics intended for use in an outpatient setting.²⁵⁸ This program requires that training be made available to all health care providers who are involved in the management of patients with pain (eg, nurses, pharmacists) and requires that education cover broader information about pain management, including non-opioid analgesics and non-pharmacologic interventions.²⁵⁹ The complete list of currently approved REMS programs is available on the FDA website.²⁶⁰

All prescribers are encouraged to discuss the risks and benefits of opioid products with their patients. A patient counseling document approved with the REMS will be made available by the manufacturers to assist the prescribers in having these discussions. Providers should also routinely



NCCN Guidelines Version 2.2022

Adult Cancer Pain

screen for signs of opioid misuse, abuse, or diversion. Various screening tools have been described for this purpose, but have not yet been evaluated in patients with cancer.³⁷ One exception is the ORT, the use of which was evaluated in a retrospective chart review of 114 patients with cancer.²⁶¹ More research is warranted to determine the best practice for screening methods. If signs of aberrant drug use are observed, urine drug testing and limiting or restricting use accordingly is strongly encouraged. However, urine drug testing results can be challenging to interpret and are susceptible to false positive and negatives; therefore, unexpected results should be followed up with confirmatory testing and/or input from a certified laboratory professional or toxicologist.²⁶²

The panel recommends that clinicians utilize state PDMPs. The National Association of State Controlled Substances Authorities (NASCA) maintains a database of state PMP contacts (see [NASCA's website](#)). Written agreements or guidelines may help to clarify expectations and parameters for safe use of opioid analgesics. While further research is needed to evaluate their utility in patients with cancer, such agreements are consistent with evolving CDC and FDA recommendations and may be required in certain states.

Management Strategies for Specific Cancer Pain Syndromes

Moderate to severe cancer pain is treated with opioids as indicated; however, opioids alone may not provide optimal analgesia. When a specific cancer pain syndrome is suspected or documented, additional interventions may be targeted to that pain syndrome. Nonopioid analgesics (such as an NSAID), adjuvant analgesics (antidepressants, anticonvulsants, topical agents, and corticosteroids), integrative interventions (psychologic and physical approaches), and/or interventional strategies may be used in conjunction with opioids to help improve patient outcomes.⁴⁹

Neuropathic Pain

Cancer-related neuropathic pain is common and can be related to the cancer itself or the acute or chronic effects of cancer treatment.²⁶³ Adjuvant analgesics are particularly important in treating neuropathic pain.^{75,76} The most common adjuvant analgesics used for treating neuropathic cancer pain include anticonvulsants, antidepressants, and topical treatments. See *Adjuvant Analgesics* for more information on these agents, including important cautions for their use. Corticosteroids have also long been used to relieve neuropathic pain syndromes, particularly radiculopathies associated with vertebral body compression fractures.

While there are a limited number of RCTs supporting the role of antidepressants as adjuvant analgesics for neuropathic cancer pain, the effectiveness of TCAs for relief of neuropathic cancer pain may be extrapolated from studies conducted in non-cancer-related neuropathic pain.²⁶⁴⁻²⁶⁶ Several RCTs have shown that anticonvulsants (pregabalin or gabapentin) provided relief of neuropathic cancer-related pain.^{85,267}

Likewise, some systematic reviews of trials of patients with cancer pain suggest that adjuvant analgesics (antidepressants and antiepileptics) added to opioids provided additional neuropathic pain relief,²⁶⁸ although another concluded that combining opioid analgesia with gabapentinoids did not provide significantly improved pain relief (data on amitriptyline, fluvoxamine, and phenytoin were inconclusive).²⁶⁹ The likelihood of benefit should be balanced with the risk of adverse effects by clinicians considering adjuvant analgesics for neuropathic pain.

Topical local anesthetic agents can be useful in preventing procedural pain and in relieving some types of cancer-related neuropathic pain. They act locally and are also thought to have some central inhibitory effect on pain. They may be used as an analgesic in combination with an opioid, antidepressant, and/or an anticonvulsant. Both the gel and patch forms of



NCCN Guidelines Version 2.2022

Adult Cancer Pain

lidocaine have been shown to reduce the pain of postherpetic neuropathy and cancer-related pain.²⁷⁰⁻²⁷²

Management of Bone Pain Without an Oncologic Emergency

The clinical complications of bone metastases include debilitating bone pain, which tends to be most prominent with movement, pathologic fractures, spinal cord compression, neurologic complications, and hypercalcemia of malignancy. The term skeletal-related events (SREs) refers to a constellation of skeletal complications including fracture, need for surgery to bone, need for radiation to bone, and spinal cord compression. In some situations, hypercalcemia of malignancy is also included as an SRE. Administration of NSAIDs, acetaminophen, or steroids may improve bone pain control when combined with opioid analgesics.²⁷³⁻²⁷⁵ Topical diclofenac, including gel or patch, may provide relief for pain due to bone metastases with minimal system effects.²⁷³

Although bone-modifying agents such as bisphosphonates and RANKL (receptor activator of nuclear factor-kappa-B ligand) inhibitors are primarily used for the reduction of overall SREs, clinical trials have established that these agents can have an analgesic effect on patients with metastatic bone pain from a variety of tumors. Clinical trials have demonstrated the palliative effects of bisphosphonates (eg, zoledronic acid, ibandronate)²⁷⁶⁻²⁸⁰ and denosumab (a RANKL inhibitor)^{278,281} on pain related to bone metastases. Randomized trials suggest that, compared with zoledronic acid, denosumab provides comparable palliation of existing bone pain and may be superior for preventing worsening of bone pain,^{278,281,282} although evidence is insufficient to recommend one of these agents over the others.²⁸³ Due to differences in patient populations and the methods for assessing bone pain, direct comparison of bisphosphonates to determine their relative effects on bone pain across studies is difficult. Review of the literature shows that the analgesic effects of bone-modifying agents are

modest and, therefore, these agents should not be used as a primary therapy for treatment of bone pain.²⁸³

Surgical and radiation treatment for bone metastases is performed to relieve local bone pain, provide stabilization, and prevent impending fracture or spinal cord compression.²⁸⁴ In some situations, interventions such as vertebral augmentation provide a greater likelihood of return to ambulatory status than radiation alone. Plain radiographs may be used to identify impending fractures so that the patient can be referred to an orthopedic specialist for stabilization. Consultation with a pain or palliative care specialist and/or an interventional therapist is recommended to determine optimal management strategy for vertebral augmentation.

Ablative strategies such as image-guided ablation may also be performed to reduce pain and prevent SREs. Image-guided ablation of bone lesions has proven successful in pain management, especially for those failing to achieve adequate analgesia without intolerable effects.²⁸⁵⁻²⁸⁸ Several small studies have also demonstrated the palliative effects of high-intensity focused US (HIFU) treatment of bone lesions.²⁸⁹⁻²⁹¹

Physical and occupational therapy may also be beneficial in the prevention of complications associated with SREs.²⁹²⁻²⁹⁴

Management of Pain from Mucositis

Certain treatments for cancer—including systemic therapy, head and neck radiation, or hematopoietic stem cell transplant—can cause pain in the mouth, pharynx, and esophagus.²⁹⁵ Resources and guidelines from the Oncology Nursing Society (ONS),²⁹⁶ Multinational Association of Supportive Care in Cancer/International Society of Oral Oncology (MASCC/ISOO),²⁹⁷ and European Society for Medical Oncology (ESMO)²⁹⁸ detail strategies for prevention and management of oral and gastrointestinal mucositis.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

To prevent oral mucositis, cryotherapy may be performed by having the patient suck on ice chips or hold ice water in his/her mouth before, during, and/or after rapid infusions of systemic therapies that are associated with mucositis. Studies have shown this approach to be effective in patients receiving melphalan for multiple myeloma and 5-fluorouracil for solid tumors.^{299,300} Gabapentin may be used in combination with opioid or non-opioid analgesics for treatment of mucositis, although studies on the effectiveness of this approach have reported mixed results.^{84,301}

Oral care protocols, consisting of good oral hygiene and prophylactic mouth rinses, may be used for prevention of mucositis.³⁰² Prophylactic mouth rinses (also called “magic mouthwash”) compositions vary significantly, including ingredients such as antibiotics, antihistamines, antifungals, corticosteroids, and antacids.^{303,304} The effectiveness of these ingredients for preventing or treating mucositis and the evidence supporting their use varies. Because of this, bland mouth rinses using ingredients such as sodium bicarbonate are often recommended.²⁹⁵ The NSAID, benzydamine, also has some data supporting its use in an oral rinse for the prevention and treatment of mucositis.^{305,306} Local anesthetics (eg, lidocaine) may be used to treat mucositis either as component of a mouth rinse or separately, in a liquid or gel formulation.

Pharmacogenetic Considerations

Pharmacogenetics can be defined as the study of how genetic differences may determine drug metabolism, and therefore response to analgesics. On the other hand, pharmacogenomics generally refers to a broader understanding of how all the genes in the genome relate to pain perception and response to analgesic drugs. There is considerable overlap in these terms, however, and they are often used interchangeably.³⁰⁷ For the purposes of this Guideline, the term pharmacogenetics will be used as it is referring to detection of specific allelic variants, rather than the entire genome. This terminology is in

agreement with that which is used in the Clinical Pharmacogenetics Implementation Consortium (CPIC) Guidelines.^{308,309}

Several classes of analgesic drugs have pharmacogenetic implications that may lead to little or no analgesic response or significant adverse effects. Notable among these are opioids,³¹⁰ TCAs,^{311,312} and NSAIDs,³¹³ which are all metabolized by the cytochrome P450s (CYPs), a family of enzymes that represent the major system for oxidative metabolism of drugs. In humans, there are at least 57 documented CYP genes across 18 different families and 44 subfamilies. CYP families 1–3 are most relevant to the pharmacogenetics of drug metabolism.³¹⁴ CYP enzymes are named with the letters “CYP” followed by a numeral designating the family, then a letter designating the sub-family, and finally another numeral that represents the individual gene or isoform. Recent evidence also supports the roles of *OPRM1* (mu receptor) and *COMT* (catechol-o-methyltransferase) in response to opioids, although current implications for clinical practice are unclear.³¹⁰

The FDA has published a table of pharmacogenetic associations that it believes have sufficient evidence to suggest that subgroups of patients with certain genetic variants have altered drug metabolism and/or therapeutic effects.³¹⁵ Of note for this guideline, codeine (*CYP2D6*), tramadol (*CYP2D6*), amitriptyline (*CYP2D6*), doxepin (*CYP2D6, CYP2C19*), celecoxib (*CYP2C9*), and meloxicam (*CYP2C9*) are included on this list. See Table 1 for potential drug-drug interactions of commonly prescribed medications and Table 2 for a more complete list of CYP 450 opioid metabolic pathways and related drug-to-drug interactions. FDA-approved pharmacogenetic tests for *CYP2D6*, *CYP2C19*, and *CYP2C9* are currently available. However, insurance reimbursement, availability of pharmacogenetic tests, and other implementation barriers may limit their use in clinical practice.³¹⁶ See *Principles of Pharmacogenetics* in the algorithm for recommendations on how to adjust



NCCN Guidelines Version 2.2022

Adult Cancer Pain

drug dosages or selection based on the results of specific pharmacogenetic tests.

The NCCN Panel recommends that pharmacogenetic testing may be considered either prior to initiation of therapy or when concerns of toxicity or lack of analgesic response arise during analgesic treatment. Consultation with a clinical pharmacist or clinical pharmacogenomics/pharmacogenetics specialist may be helpful in the interpretation of pharmacogenetic test results and to aid in drug selection and/or dose adjustments, when necessary.

Specialty Consultations

Continued pain assessments should be obtained and documented in the medical record to ensure that the patient's pain remains well-managed and goals of treatment are achieved. Specialty consultations can be helpful in providing interventions to assist with difficult cancer pain problems. The major indication for referral to a specialty service provider is if the pain is likely to be relieved or will help patients become functional in their daily activities. These modalities are delivered by a specialty service provider, and pain management is accomplished by establishing individualized goals and then providing specific treatment and education for patients. The specialties include physical/occupational therapy; integrative medicine; mental health support services (including psychiatric consultation, psychology consultation, and/or substance abuse consultation, as needed); pain and palliative care services; depression/distress consultation; spiritual care consultation; or social work services. Consultation with an addiction specialist may be helpful for managing aberrant drug behavior in patients with a history of or risk factors for abuse or misuse of pain medication.

Non-Pharmacologic Interventions for Cancer Pain Management

Integrative Interventions

Since pain encompasses physical, psychosocial, and spiritual dimensions, the treatment of cancer pain inherently requires integration of therapies inclusive of non-pharmacologic interventions. There is a growing body of evidence suggesting that the use of nonpharmacologic interventions (physical, cognitive, psychosocial, and spiritual) may serve as valuable additions to pharmacologic interventions.³¹⁷⁻³¹⁹ The integration of physical, cognitive, psychosocial, and spiritual modalities should be based on assessment of cultural and financial considerations, and are best presented as part of joint and informed decision-making.

Physical Interventions

Physical interventions include, but aren't limited to, therapeutic or conditioning exercise, physical or occupational therapy, massage, use of heat and/or cold, acupuncture, and acupressure.³²⁰⁻³²⁴

Cognitive-Behavioral Interventions

Cognitive interventions are aimed at enhancing a sense of control over the pain or underlying disease. Mindfulness-based stress reduction (MBSR), breathing exercises, relaxation, imagery, hypnosis, biofeedback, music, and other behavioral therapies can be very useful.³²⁵⁻³³¹ Patient-based educational interventions have a significant impact in providing pain relief.³³² Skills training helps modify the patient's experience of pain and helps patients acquire techniques of pain management such as deep muscle relaxation. Patients who may benefit from skills training may be referred to a licensed mental health professional trained in cognitive behavioral therapy, hypnosis, biofeedback, or MBSR. Education provides patients and family/caregivers with the knowledge to use analgesics correctly and to address side effects or unrelieved pain.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

Psychosocial Interventions

Attention should focus on psychosocial support and providing education to patients and families.^{333,334} Psychosocial support can greatly enhance patients' sense of control as well as greatly reduce the family/caregivers' feeling of helplessness.³²⁹ A meta-analysis of the effect of psychosocial interventions on cancer pain highlights the importance of a multimodal approach to the management of cancer pain.³³⁵

Spiritual Interventions

In cancer care, there is growing interest in attention to spiritual needs and the existential concerns often associated with pain. Many patients hold cultural beliefs about such treatments, and home remedies, rituals, prayer, and other spiritual practices may be most helpful in relieving or coping with pain. Involvement of spiritual care providers from a range of culturally appropriate spiritual backgrounds is essential.³³⁶ Spiritual needs should be routinely assessed and spiritual care should be incorporated as a component of comprehensive pain management.

Interventional Strategies

Some patients experience inadequate pain management despite pharmacologic therapy or may not tolerate an opioid titration program because of side effects. Some patients may prefer interventional therapies instead of a chronic medication regimen. Interventional techniques have been demonstrated, in some cases, to eliminate or significantly reduce the level of pain, and/or may allow a significant decrease in systemic analgesics. Interventional therapies that can be useful in the relief of cancer pain include nerve blocks, vertebral augmentation, regional infusion of analgesics, image-guided ablation, and other techniques.^{49,287,288,337-341}

The major indications for referral for interventional therapies include a patient suffering from pain that is likely to be relieved with nerve block (eg,

pancreas/upper abdomen with celiac plexus block, lower abdomen with superior hypogastric plexus block, intercostal nerve, peripheral/plexus nerve) and/or patients unable to achieve adequate analgesia and/or the presence of intolerable side effects. For example, a patient with pancreatic cancer who was not tolerating opioids or not receiving adequate analgesia could be offered a neurolytic celiac plexus block. Neurolytic celiac plexus block may offer some improvement in pain management over systemic analgesics, but is generally associated with a reduction in adverse effects.^{342,343}

Regional infusion of analgesics (epidural, intrathecal, and regional plexus) minimizes the distribution of drugs to receptors in the brain, as well as lowering serum opioid levels, potentially avoiding adverse effects of systemic administration. Therefore, the intrathecal route of opioid administration should be considered in patients with intolerable sedation, confusion, constipation, and/or inadequate pain management with systemic opioid administration.³⁴⁴⁻³⁴⁶ This approach is a valuable tool to improve analgesia for patients who have pain from a variety of anatomical locations (eg, head and neck, upper and lower extremities, trunk).³⁴⁷⁻³⁵⁰ However, due to the risk of catheter migration and infection risk, consider limiting the duration of use to several days.

Percutaneous vertebral augmentation and/or cementoplasty might be useful for the treatment of lytic osteoclastic spinal metastases or in cases of vertebral compression fractures or spinal instability for which surgery is not feasible or indicated. Vertebral augmentation helps restore mechanical stability while reducing pain and neurologic symptoms.³⁵¹⁻³⁵⁶ Ablation techniques may also be helpful for pain management in patients who receive inadequate relief from pharmacologic therapy. Prospective trials of percutaneous ablative techniques, many using thermal energy, have shown decreased patient pain from bone metastases in patients who did or did not receive prior radiation therapy.³⁵⁷⁻³⁶⁶ Non-ionizing thermal



NCCN Guidelines Version 2.2022

Adult Cancer Pain

ablative techniques may serve as an alternative and/or adjunct to radiation therapy or be offered in patients who refuse or cannot receive radiation therapy. Early data suggest a synergistic effect with radiation therapy and these different treatment modalities may prove to be complementary.^{367,368} Similarly, vertebral augmentation/cementoplasty provides pain relief with the additional benefit of improved stabilization, which may prevent or halt pathologic fracture.³⁶⁹

Neurodestructive procedures may be used for well-localized pain syndromes (eg, back pain due to facet or sacroiliac joint arthropathy; visceral pain due to abdominal or pelvic malignancy). Ablation therapy (eg, image-guided ablation, US ablation) for bone lesions can also be helpful in reducing pain.²⁸⁵⁻²⁹¹ See *Management Strategies for Specific Cancer Pain Syndromes, Bone Pain Without Oncologic Emergency* in the algorithm for more information. Neurostimulation procedures have been suggested to be useful for painful chemotherapy-induced peripheral neuropathies, neuralgias, and complex regional pain syndrome.³⁷⁰

Interventional strategies listed above are not appropriate if patients are unwilling or in patients with infections, coagulopathy, or with very short life expectancies. Also, the experts performing the interventions must be made aware of any medications that the patient is taking that might increase bleeding risk (ie, anticoagulants [warfarin, heparin], antiplatelet agents [clopidogrel, dipyridamole], anti-angiogenesis agents [bevacizumab]). If this occurs, the patient may have to be off the medication for an appropriate amount of time prior to the pain intervention and may need to continue to stay off the medication for a specified amount of time after the procedure. Interventions are not appropriate if technical expertise is not available. Additionally, if interventional treatment is undertaken and successfully improves pain control, significant opioid dose reduction may be required.

Summary

In most patients, cancer pain can be successfully managed with appropriate techniques and safe drugs. The overall approach to pain management encompassed in these guidelines is multimodal and comprehensive. It is based on routine pain assessments, utilizes both pharmacologic and nonpharmacologic interventions, and requires ongoing reevaluation of the patient. The NCCN Adult Cancer Pain Guidelines Panel advises that cancer pain can be well managed in the vast majority of patients if the algorithms presented are systematically applied, carefully monitored, and tailored to the needs of the individual patient.

Recommended Readings

- Brant JM, Rodgers BB, Gallagher E, Sundaramurthi T. Breakthrough cancer pain: A systematic review of pharmacologic management. *Clin J Oncol Nurs* 2017;21:71-80.
- Cherny N. Cancer Pain Syndromes. In: Cherny N, Fallon M, Kaasa S, et al., eds. *Oxford Textbook of Palliative Medicine* (5th ed). Oxford: Oxford University Press; 2015:819-840.
- Kwon JH. Overcoming barriers in cancer pain management. *J Clin Oncol* 2014;32:1727-1733.
- Liu WC, Zheng ZX, Tan KH, Meredith GJ. Multidimensional treatment of cancer pain. *Curr Oncol Rep* 2017;19:10.
- Paice JA, Ferrell B. The management of cancer pain. *CA Cancer J Clin* 2011;61:157-182.
- Schmidt BL. The neurobiology of cancer pain. *Neuroscientist* 2014;20:546-562.
- Wiffen PJ, Wee B, Derry S, et al. Opioids for cancer pain - an overview of Cochrane reviews. *Cochrane Database Syst Rev* 2017;7:CD012592.

Table 1: Potential Drug-Drug Interactions: Chemotherapeutics, Analgesics, and Other Commonly Prescribed Medications^{*,β}

Drug	Buprenorphine, Fentanyl, Hydrocodone, Methadone, & Oxycodone	Methadone & Buprenorphine	Enzalutamide, Apalutamide, Mitotane, & Dexamethasone ^{ε,φ}
Interaction	Potential to <u>increase</u> plasma levels of the above opioids	Potential for QTc prolongation when used with above opioids	Potential to <u>decrease</u> plasma levels of the agents below
Interacting Drugs	Clarithromycin Cobicistat Conivaptan Erythromycin Fluconazole Imatinib Indinavir Itraconazole Ketoconazole (systemic) Nelfinavir Nefazodone Posaconazole Ritonavir Saquinavir Verapamil Voriconazole	Abarelix Ceritinib Citalopram Crizotinib Bortezomib Bevacizumab Dabrafenib Dasatinib Degarelix Dolasetron Doxorubicin Epirubicin Fluoroquinolones Granisetron Haloperidol Ibrutinib Idelalisib Lapatinib Lenvatinib Metoclopramide Nilotinib Olanzapine Ondansetron Pazopanib Prochlorperazine Ribociclib Risperdal Ruxolitinib Sorafenib Sunitinib Toremifene Vandetanib Vemurafenib Voriconazole Ziprasidone	Aprepitant Buprenorphine Bortezomib Ceritinib Crizotinib Dabrafenib Erlotinib Everolimus Fentanyl Gefitinib Hydrocodone Ibrutinib Idelalisib Imatinib Lapatinib Methadone Naldemedine Naloxegol Oxycodone Pazopanib Ruxolitinib Sirolimus Sorafenib Sunitinib Tacrolimus Temsirolimus Vandetanib Vemurafenib

*Data within this table were obtained from the FDA drug Development and Drug Interactions: Table of Substrates, Inhibitors, and Inducers³⁷¹ and Lexicomp Online (Hudson, Ohio: Lexicomp, Inc.), available published literature, and prescribing information for drug products. Information updated on May 10, 2021.

β This list is not comprehensive and may not represent new data or other agents recently introduced into practice. Clinicians are advised to refer to the individual drug labeling or seek expert consultation.

ε Many chemotherapeutic agents produce immunosuppression that can be exacerbated by concomitant dexamethasone use; physicians should consider goals of care, rationale for dexamethasone use, duration of use, and other factors when considering use with other immunosuppressive agents.

φ Dexamethasone is an inducer of cytochrome P450 3A.

NCCN Guidelines Version 2.2022

Adult Cancer Pain

Table 2: CYP P450 Opioid Metabolism and Potential Drug-Drug Interactions

Opioid	CYP P450 Metabolic Pathway ³⁷²⁻³⁸¹	CYP 3A4 Inducers ³⁷¹	CYP 3A4 Inhibitors ³⁷¹		
Buprenorphine	CYP 3A4	Carbamazepine ^Ω Rifampin ^Ω Efavirenz Ritonavir Oxcarbazepine St john's wort Phenobarbital Enzalutamide ^Ω Phenytoin ^Ω Apalutamide Primidone Mitotane Rifabutin Dexamethasone	Itraconazole** Nelfinavir** Ketoconazole** Indinavir** Nefazodone** Clarithromycin** Posaconazole** Verapamil* Voriconazole** Aprepitant* Atazanavir** Conivaptan* Boceprevir** Diltiazem* Ritonavir** Dronedarone* Saquinavir** Erythromycin* Telaprevir** Fluconazole*		
Codeine	CYP2D6				
Fentanyl	CYP3A4				
Hydrocodone	CYP2D6, CYP3A4				
Methadone	CYP2B6, CYP 3A4 CYP2C8, CYP2C19 CYP2D6, CYP2C9				
Oxycodone	CYP2D6, CYP3A4				
Tapentadol	CYP2C9, CYP2C19 CYP2D6 ^β			CYP 2D6 Inducers ³⁷¹	CYP 2D6 Inhibitors ³⁷¹
Tramadol	CYP2D6, CYP3A4			No Data	Quinidine** Paroxetine** Fluoxetine** Bupropion**
Morphine	None				
Hydromorphone	None				
Oxymorphone	None				

Created in collaboration by Rebecca Nelson, PharmD and David Craig, PharmD and reviewed by the NCCN Panel (last updated 5/10/2021)

**Denotes strong inhibitors (>5-fold increase in exposure, or >80% decrease in clearance of substrate)

*Denotes moderate inhibitors (>2 to <5-fold increase in exposure, or 50% to 80% decrease in clearance of substrate).

β Tapentadol is not extensively metabolized via CYP 450 enzymes, therefore, clinically relevant interactions mediated by this metabolic pathway are unlikely to occur³⁸⁰

Ω Denotes strong inducer (> 80% reduction in AUC of substrate)



References

1. Raja SN, Carr DB, Cohen M, et al. The revised International Association for the Study of Pain definition of pain: concepts, challenges, and compromises. *Pain* 2020;161:1976-1982. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32694387>.
2. van den Beuken-van Everdingen MH, de Rijke JM, Kessels AG, et al. Prevalence of pain in patients with cancer: a systematic review of the past 40 years. *Ann Oncol* 2007;18:1437-1449. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17355955>
3. Te Boveldt N, Vernooij-Dassen M, Burger N, et al. Pain and its interference with daily activities in medical oncology outpatients. *Pain Physician* 2013;16:379-389. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23877454>.
4. Temel JS, Greer JA, Muzikansky A, et al. Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010;363:733-742. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20818875>.
5. Zimmermann C, Swami N, Krzyzanowska M, et al. Early palliative care for patients with advanced cancer: a cluster-randomised controlled trial. *Lancet* 2014;383:1721-1730. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24559581>.
6. Grudzen CR, Richardson LD, Johnson PN, et al. Emergency department-initiated palliative care in advanced cancer: A randomized clinical trial. *JAMA Oncol* 2016;2:591-598. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26768772>.
7. Ferrell B, Sun V, Hurria A, et al. Interdisciplinary palliative care for patients with lung cancer. *J Pain Symptom Manage* 2015;50:758-767. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26296261>.
8. Bakitas MA, Tosteson TD, Li Z, et al. Early versus delayed initiation of concurrent palliative oncology care: patient outcomes in the ENABLE III randomized controlled trial. *J Clin Oncol* 2015;33:1438-1445. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25800768>.
9. Bakitas M, Lyons KD, Hegel MT, et al. Effects of a palliative care intervention on clinical outcomes in patients with advanced cancer: the Project ENABLE II randomized controlled trial. *Jama* 2009;302:741-749. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19690306>.
10. Greco MT, Roberto A, Corli O, et al. Quality of cancer pain management: an update of a systematic review of undertreatment of patients with cancer. *J Clin Oncol* 2014;32:4149-4154. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25403222>.
11. Fairchild A. Under-treatment of cancer pain. *Curr Opin Support Palliat Care* 2010;4:11-15. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20040878>.
12. Passik SD, Weinreb HJ. Managing chronic nonmalignant pain: overcoming obstacles to the use of opioids. *Adv Ther* 2000;17:70-83. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11010058>.
13. Cleeland CS, Gonin R, Hatfield AK, et al. Pain and its treatment in outpatients with metastatic cancer. *N Engl J Med* 1994;330:592-596. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7508092>.
14. Martin LA, Hagen NA. Neuropathic pain in cancer patients: mechanisms, syndromes, and clinical controversies. *J Pain Symptom Manage* 1997;14:99-9117. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9262040>.
15. Mercadante S. Malignant bone pain: pathophysiology and treatment. *Pain* 1997;69:1-18. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9060007>.
16. Stjernsward J. WHO cancer pain relief programme. *Cancer Surv* 1988;7:195-208. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2454740>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

17. Stjernsward J, Colleau SM, Ventafridda V. The World Health Organization Cancer Pain and Palliative Care Program. Past, present, and future. *J Pain Symptom Manage* 1996;12:65-72. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8754982>.
18. Caraceni A, Weinstein SM. Classification of cancer pain syndromes. *Oncology (Williston Park)* 2001;15:1627-1640. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11780704>.
19. Hewitt DJ. The management of pain in the oncology patient. *Obstet Gynecol Clin North Am* 2001;28:819-846. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11766154>.
20. Portenoy RK. Cancer pain. Epidemiology and syndromes. *Cancer* 1989;63:2298-2307. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2655867>.
21. Hicks CL, von Baeyer CL, Spafford PA, et al. The Faces Pain Scale-Revised: toward a common metric in pediatric pain measurement. *Pain* 2001;93:173-183. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11427329>.
22. Serlin RC, Mendoza TR, Nakamura Y, et al. When is cancer pain mild, moderate or severe? Grading pain severity by its interference with function. *Pain* 1995;61:277-284. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7659438>.
23. Soetenga D, Frank J, Pellino TA. Assessment of the validity and reliability of the University of Wisconsin Children's Hospital Pain Scale for Preverbal and Nonverbal Children. *Pediatr Nurs* 1999;25:670-676. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12024390>.
24. Ware LJ, Epps CD, Herr K, Packard A. Evaluation of the Revised Faces Pain Scale, Verbal Descriptor Scale, Numeric Rating Scale, and Iowa Pain Thermometer in older minority adults. *Pain Manag Nurs* 2006;7:117-125. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16931417>.
25. Cleeland CS, Nakamura Y, Mendoza TR, et al. Dimensions of the impact of cancer pain in a four country sample: new information from multidimensional scaling. *Pain* 1996;67:267-273. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8951920>.
26. Cleeland CS, Ryan KM. Pain assessment: global use of the Brief Pain Inventory. *Ann Acad Med Singapore* 1994;23:129-138. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8080219>.
27. Holen JC, Lydersen S, Klepstad P, et al. The Brief Pain Inventory: pain's interference with functions is different in cancer pain compared with noncancer chronic pain. *Clin J Pain* 2008;24:219-225. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18287827>.
28. Sonis ST, Eilers JP, Epstein JB, et al. Validation of a new scoring system for the assessment of clinical trial research of oral mucositis induced by radiation or chemotherapy. Mucositis Study Group. *Cancer* 1999;85:2103-2113. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10326686>.
29. Kushner JA, Lawrence HP, Shoal I, et al. Development and validation of a Patient-Reported Oral Mucositis Symptom (PROMS) scale. *J Can Dent Assoc* 2008;74:59. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18298885>.
30. Gussgard AM, Hope AJ, Jokstad A, et al. Assessment of cancer therapy-induced oral mucositis using a patient-reported oral mucositis experience questionnaire. *PLoS One* 2014;9:e91733. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24614512>.
31. Callin S, Bennett MI. Assessment of neuropathic pain. *Continuing Education in Anaesthesia Critical Care & Pain* 2008;8:210-213. Available at: <http://www.sciencedirect.com/science/article/pii/S1743181617304559>.
32. Bennett MI, Attal N, Backonja MM, et al. Using screening tools to identify neuropathic pain. *Pain* 2007;127:199-203. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17182186>.



33. Broom R, Du H, Clemons M, et al. Switching breast cancer patients with progressive bone metastases to third-generation bisphosphonates: measuring impact using the Functional Assessment of Cancer Therapy-Bone Pain. *J Pain Symptom Manage* 2009;38:244-257. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19364633>.
34. Amtmann D, Cook KF, Jensen MP, et al. Development of a PROMIS item bank to measure pain interference. *Pain* 2010;150:173-182. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20554116>.
35. Al-Atiyyat HNM. Cultural diversity and cancer pain. *Journal of Hospice & Palliative Nursing* 2009;11:154-164. Available at: http://journals.lww.com/jhpn/Abstract/2009/05000/Cultural_Diversity_and_Cancer_Pain.9.aspx.
36. Ezenwa MO, Ameringer S, Ward SE, Serlin RC. Racial and ethnic disparities in pain management in the United States. *J Nurs Scholarsh* 2006;38:225-233. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17044339>.
37. Angheliescu DL, Ehrentraut JH, Faughnan LG. Opioid misuse and abuse: risk assessment and management in patients with cancer pain. *J Natl Compr Canc Netw* 2013;11:1023-1031. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23946178>.
38. Akbik H, Butler SF, Budman SH, et al. Validation and clinical application of the Screener and Opioid Assessment for Patients with Pain (SOAPP). *J Pain Symptom Manage* 2006;32:287-293. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16939853>.
39. Butler SF, Fernandez K, Benoit C, et al. Validation of the revised Screener and Opioid Assessment for Patients with Pain (SOAPP-R). *J Pain* 2008;9:360-372. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18203666>.
40. Webster LR, Webster RM. Predicting aberrant behaviors in opioid-treated patients: preliminary validation of the Opioid Risk Tool. *Pain Med* 2005;6:432-442. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16336480>.
41. Passik SD, Kirsh KL. The interface between pain and drug abuse and the evolution of strategies to optimize pain management while minimizing drug abuse. *Exp Clin Psychopharmacol* 2008;16:400-404. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18837636>.
42. Chou R, Fanciullo GJ, Fine PG, et al. Opioids for chronic noncancer pain: prediction and identification of aberrant drug-related behaviors: a review of the evidence for an American Pain Society and American Academy of Pain Medicine clinical practice guideline. *J Pain* 2009;10:131-146. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19187890>.
43. Arthur JA, Tang M, Lu Z, et al. Random urine drug testing among patients receiving opioid therapy for cancer pain. *Cancer* 2021;127:968-975. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33231885>.
44. U.S. Food and Drug Administration. Transmucosal Immediate Release Fentanyl (TIRF) Risk Evaluation and Mitigation Strategy (REMS). Silver Spring, MD: 2014. Available at: <http://www.fda.gov/downloads/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/UCM289730.pdf>. Accessed April 6, 2021.
45. U.S. Food and Drug Administration. Extended-Release (ER) and Long-Acting (LA) Opioid Analgesics Risk Evaluation and Mitigation Strategy (REMS). Silver Spring, MD: 2015. Available at: <http://www.fda.gov/downloads/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/UCM311290.pdf>. Accessed April 6, 2021.
46. Stewart M, Brown JB, Donner A, et al. The impact of patient-centered care on outcomes. *J Fam Pract* 2000;49:796-804. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11032203>.
47. Syrjala KL, Abrams JR, Polissar NL, et al. Patient training in cancer pain management using integrated print and video materials: a multisite randomized controlled trial. *Pain* 2008;135:175-186. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18093738>.



48. Mercadante SL, Berchovich M, Casuccio A, et al. A prospective randomized study of corticosteroids as adjuvant drugs to opioids in advanced cancer patients. *Am J Hosp Palliat Care* 2007;24:13-19. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17347500>.
49. American Pain Society. *Principles of Analgesic Use*. (ed 7th). Glenview, IL: American Pain Society; 2016.
50. Patel PM, Goodman LF, Knepel SA, et al. Evaluation of emergency department management of opioid-tolerant cancer patients with acute pain. *J Pain Symptom Manage* 2017;54:501-507. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28729010>.
51. Portenoy RK, Lesage P. Management of cancer pain. *Lancet* 1999;353:1695-1700. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10335806>.
52. Mercadante S, Arcuri E, Ferrera P, et al. Alternative treatments of breakthrough pain in patients receiving spinal analgesics for cancer pain. *J Pain Symptom Manage* 2005;30:485-491. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16310622>.
53. Green CR, Anderson KO, Baker TA, et al. The unequal burden of pain: confronting racial and ethnic disparities in pain. *Pain Med* 2003;4:277-294. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12974827>.
54. Gordon DB, Pellino TA, Miaskowski C, et al. A 10-year review of quality improvement monitoring in pain management: recommendations for standardized outcome measures. *Pain Manag Nurs* 2002;3:116-130. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12454804>.
55. Sun VC, Borneman T, Ferrell B, et al. Overcoming barriers to cancer pain management: an institutional change model. *J Pain Symptom Manage* 2007;34:359-369. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17616336>.
56. Lin CC, Chou PL, Wu SL, et al. Long-term effectiveness of a patient and family pain education program on overcoming barriers to management of cancer pain. *Pain* 2006;122:271-281. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16545909>.
57. Chang MC, Chang YC, Chiou JF, et al. Overcoming patient-related barriers to cancer pain management for home care patients. A pilot study. *Cancer Nurs* 2002;25:470-476. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12464839>.
58. Key findings summary: Opioid access research project. American Cancer Society Cancer Action Network (ACS CAN) and the Patient Quality of Life Coalition (PQLC); 2018. Available at: <https://www.acscan.org/sites/default/files/ACS%20CAN%20PQLC%20Opioid%20Research%20Project%20Key%20Findings%20Summary%20Memo%20FINAL.pdf>. Accessed April 06, 2021.
59. Schumacher KL, Koresawa S, West C, et al. The usefulness of a daily pain management diary for outpatients with cancer-related pain. *Oncol Nurs Forum* 2002;29:1304-1313. Available at: <https://pubmed.ncbi.nlm.nih.gov/12370700/>.
60. Pachman DR, Barton DL, Swetz KM, Loprinzi CL. Troublesome symptoms in cancer survivors: fatigue, insomnia, neuropathy, and pain. *J Clin Oncol* 2012;30:3687-3696. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23008320>.
61. Paice JA, Ferrell B. The management of cancer pain. *CA Cancer J Clin* 2011;61:157-182. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21543825>.
62. Paice JA, Lacchetti C, Bruera E. Management of Chronic Pain in Survivors of Adult Cancers: ASCO Clinical Practice Guideline Summary. *J Oncol Pract* 2016;12:757-762. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27460497>.
63. Stockler M, Vardy J, Pillai A, Warr D. Acetaminophen (paracetamol) improves pain and well-being in people with advanced cancer already receiving a strong opioid regimen: a randomized, double-blind, placebo-controlled cross-over trial. *J Clin Oncol* 2004;22:3389-3394. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15310785>.



64. Pharmacological management of persistent pain in older persons. *J Am Geriatr Soc* 2009;57:1331-1346. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19573219>.
65. Israel FJ, Parker G, Charles M, Reymond L. Lack of benefit from paracetamol (acetaminophen) for palliative cancer patients requiring high-dose strong opioids: a randomized, double-blind, placebo-controlled, crossover trial. *J Pain Symptom Manage* 2010;39:548-554. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20083373>.
66. U.S. Food and Drug Administration. FDA Drug Safety Communication: Prescription Acetaminophen Products to be Limited to 325 mg Per Dosage Unit; Boxed Warning Will Highlight Potential for Severe Liver Failure. 2011. Available at: <http://www.fda.gov/drugs/drugsafety/ucm239821.htm>. Accessed April 06, 2021.
67. U.S. Food and Drug Administration. Information for Healthcare Professionals: Concomitant Use of Ibuprofen and Aspirin. 2006. Available at: <https://www.fda.gov/downloads/drugs/drugsafety/postmarketdrugsafety/informationforpatientsandproviders/ucm161282.pdf>. Accessed April 06, 2021.
68. Tielemans MM, Eikendal T, Jansen JB, van Oijen MG. Identification of NSAID users at risk for gastrointestinal complications: a systematic review of current guidelines and consensus agreements. *Drug Saf* 2010;33:443-453. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20486727>.
69. Laine L, Curtis SP, Cryer B, et al. Risk factors for NSAID-associated upper GI clinical events in a long-term prospective study of 34 701 arthritis patients. *Aliment Pharmacol Ther* 2010;32:1240-1248. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20955443>.
70. Magee DJ, Jhanji S, Poulogiannis G, et al. Nonsteroidal anti-inflammatory drugs and pain in cancer patients: a systematic review and reappraisal of the evidence. *Br J Anaesth* 2019;123:e412-e423. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31122736>.
71. Derry S, Wiffen PJ, Moore RA, et al. Oral nonsteroidal anti-inflammatory drugs (NSAIDs) for cancer pain in adults. *Cochrane Database Syst Rev* 2017;7:CD012638. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28700091>.
72. U.S. Food and Drug Administration. FDA Drug Safety Communication: FDA strengthens warning that non-aspirin nonsteroidal anti-inflammatory drugs (NSAIDs) can cause heart attacks or strokes. 2015. Available at: <https://www.fda.gov/drugs/drug-safety-and-availability/fda-drug-safety-communication-fda-strengthens-warning-non-aspirin-nonsteroidal-anti-inflammatory>. Accessed April 06, 2021.
73. Bally M, Dendukuri N, Rich B, et al. Risk of acute myocardial infarction with NSAIDs in real world use: bayesian meta-analysis of individual patient data. *Bmj* 2017;357:j1909. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28487435>.
74. Lussier D, Huskey AG, Portenoy RK. Adjuvant analgesics in cancer pain management. *Oncologist* 2004;9:571-591. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15477643>.
75. Manfredi PL, Gonzales GR, Sady R, et al. Neuropathic pain in patients with cancer. *J Palliat Care* 2003;19:115-118. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12955928>.
76. Jongen JL, Huijsman ML, Jessurun J, et al. The evidence for pharmacologic treatment of neuropathic cancer pain: beneficial and adverse effects. *J Pain Symptom Manage* 2013;46:581-590 e581. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23415040>.
77. Jin Y, Desta Z, Stearns V, et al. CYP2D6 genotype, antidepressant use, and tamoxifen metabolism during adjuvant breast cancer treatment. *J Natl Cancer Inst* 2005;97:30-39. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15632378>.
78. Haque R, Shi J, Schottinger JE, et al. Tamoxifen and antidepressant drug interaction in a cohort of 16,887 breast cancer survivors. *J Natl Cancer Inst* 2016;108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26631176>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

79. Azoulay L, Dell'Aniello S, Huiart L, et al. Concurrent use of tamoxifen with CYP2D6 inhibitors and the risk of breast cancer recurrence. *Breast Cancer Res Treat* 2011;126:695-703. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20848186>.
80. Johannessen Landmark C. Antiepileptic drugs in non-epilepsy disorders: relations between mechanisms of action and clinical efficacy. *CNS Drugs* 2008;22:27-47. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18072813>.
81. Derry S, Bell RF, Straube S, et al. Pregabalin for neuropathic pain in adults. *Cochrane Database Syst Rev* 2019;1:CD007076. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30673120>.
82. Chen DL, Li YH, Wang ZJ, Zhu YK. The research on long-term clinical effects and patients' satisfaction of gabapentin combined with oxycontin in treatment of severe cancer pain. *Medicine (Baltimore)* 2016;95:e5144. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27759644>.
83. Dou Z, Jiang Z, Zhong J. Efficacy and safety of pregabalin in patients with neuropathic cancer pain undergoing morphine therapy. *Asia Pac J Clin Oncol* 2017;13:e57-e64. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25530068>.
84. Bar Ad V, Weinstein G, Dutta PR, et al. Gabapentin for the treatment of pain syndrome related to radiation-induced mucositis in patients with head and neck cancer treated with concurrent chemoradiotherapy. *Cancer* 2010;116:4206-4213. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20564146>.
85. Raptis E, Vadalouca A, Stavropoulou E, et al. Pregabalin vs. opioids for the treatment of neuropathic cancer pain: a prospective, head-to-head, randomized, open-label study. *Pain Pract* 2014;14:32-42. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23464813>.
86. Wooldridge JE, Anderson CM, Perry MC. Corticosteroids in advanced cancer. *Oncology (Williston Park)* 2001;15:225-234; discussion 234-226. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11252935>.
87. Haywood A, Good P, Khan S, et al. Corticosteroids for the management of cancer-related pain in adults. *Cochrane Database Syst Rev* 2015;4:CD010756. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25908299>.
88. Donovan KA, Chang YD, Oberoi-Jassal R, et al. Relationship of cannabis use to patient-reported symptoms in cancer patients seeking supportive/palliative care. *J Palliat Med* 2019;22:1191-1195. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30794025>.
89. Steele G, Arneson T, Zylla D. A comprehensive review of cannabis in patients with cancer: Availability in the USA, general efficacy, and safety. *Curr Oncol Rep* 2019;21:10. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30707319>.
90. National Center for Complementary and Integrative Health. Cannabis (Marijuana) and Cannabinoids: What You Need To Know. 2019. Available at: <https://nccih.nih.gov/health/marijuana>. Accessed April 06, 2021.
91. Drugs of Abuse: a DEA Resource Guide; 2017. Available at: https://www.dea.gov/sites/default/files/sites/getsmartaboutdrugs.com/files/publications/DoA_2017Ed_Updated_6.16.17.pdf.
92. Pergam SA, Woodfield MC, Lee CM, et al. Cannabis use among patients at a comprehensive cancer center in a state with legalized medicinal and recreational use. *Cancer* 2017;123:4488-4497. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28944449>.
93. Tringale KR, Huynh-Le MP, Salans M, et al. The role of cancer in marijuana and prescription opioid use in the United States: A population-based analysis from 2005 to 2014. *Cancer* 2019. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31006849>.
94. Johnson JR, Burnell-Nugent M, Lossignol D, et al. Multicenter, double-blind, randomized, placebo-controlled, parallel-group study of the efficacy, safety, and tolerability of THC:CBD extract and THC extract in patients with intractable cancer-related pain. *J Pain Symptom Manage* 2010;39:167-179. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19896326>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

95. Portenoy RK, Ganae-Motan ED, Allende S, et al. Nabiximols for opioid-treated cancer patients with poorly-controlled chronic pain: a randomized, placebo-controlled, graded-dose trial. *J Pain* 2012;13:438-449. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22483680>.
96. Lynch ME, Cesar-Rittenberg P, Hohmann AG. A double-blind, placebo-controlled, crossover pilot trial with extension using an oral mucosal cannabinoid extract for treatment of chemotherapy-induced neuropathic pain. *J Pain Symptom Manage* 2014;47:166-173. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23742737>.
97. Hauser W, Welsch P, Klose P, et al. Efficacy, tolerability and safety of cannabis-based medicines for cancer pain : A systematic review with meta-analysis of randomised controlled trials. *Schmerz* 2019;33:424-436. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31073761>.
98. Boland EG, Bennett MI, Allgar V, Boland JW. Cannabinoids for adult cancer-related pain: systematic review and meta-analysis. *BMJ Support Palliat Care* 2020;10:14-24. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31959586>.
99. Rabgay K, Waranuch N, Chaiyakunapruk N, et al. The effects of cannabis, cannabinoids, and their administration routes on pain control efficacy and safety: A systematic review and network meta-analysis. *J Am Pharm Assoc (2003) 2020;60:225-234 e226*. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31495691>.
100. Monte AA, Shelton SK, Mills E, et al. Acute illness associated with cannabis use, by route of exposure: An observational study. *Ann Intern Med* 2019;170:531-537. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30909297>.
101. Cherny NI. The pharmacologic management of cancer pain. *Oncology (Williston Park)* 2004;18:1499-1515. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15609474>.
102. Bandieri E, Romero M, Ripamonti CI, et al. Randomized trial of low-dose morphine versus weak opioids in moderate cancer pain. *J Clin Oncol* 2016;34:436-442. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26644526>.
103. Andersen G, Jensen NH, Christrup L, et al. Pain, sedation and morphine metabolism in cancer patients during long-term treatment with sustained-release morphine. *Palliat Med* 2002;16:107-114. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11969141>.
104. Smith MT. Neuroexcitatory effects of morphine and hydromorphone: evidence implicating the 3-glucuronide metabolites. *Clin Exp Pharmacol Physiol* 2000;27:524-528. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10874511>.
105. Sande TA, Laird BJ, Fallon MT. The use of opioids in cancer patients with renal impairment-a systematic review. *Support Care Cancer* 2017;25:661-675. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27744535>.
106. Mercadante S. Intravenous morphine for management of cancer pain. *Lancet Oncol* 2010;11:484-489. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20434717>.
107. Klepstad P, Kaasa S, Borchgrevink PC. Start of oral morphine to cancer patients: effective serum morphine concentrations and contribution from morphine-6-glucuronide to the analgesia produced by morphine. *Eur J Clin Pharmacol* 2000;55:713-719. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10663448>.
108. Klepstad P, Kaasa S, Skauge M, Borchgrevink PC. Pain intensity and side effects during titration of morphine to cancer patients using a fixed schedule dose escalation. *Acta Anaesthesiol Scand* 2000;44:656-664. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10903012>.
109. Foley KM. The treatment of pain in the patient with cancer. *CA Cancer J Clin* 1986;36:194-215. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2425909>.
110. Tiseo PJ, Thaler HT, Lapin J, et al. Morphine-6-glucuronide concentrations and opioid-related side effects: a survey in cancer patients.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

Pain 1995;61:47-54. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/7644248>.

111. Portenoy RK, Foley KM, Stulman J, et al. Plasma morphine and morphine-6-glucuronide during chronic morphine therapy for cancer pain: plasma profiles, steady-state concentrations and the consequences of renal failure. Pain 1991;47:13-19. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/1771088>.

112. Trescot AM, Datta S, Lee M, Hansen H. Opioid pharmacology. Pain Physician 2008;11:S133-153. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18443637>.

113. Mercadante S, Vellucci R, Cuomo A, et al. Long-term efficacy and tolerability of intranasal fentanyl in the treatment of breakthrough cancer pain. Support Care Cancer 2015;23:1349-1354. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25351457>.

114. Caraceni A, Hanks G, Kaasa S, et al. Use of opioid analgesics in the treatment of cancer pain: evidence-based recommendations from the EAPC. Lancet Oncol 2012;13:e58-68. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22300860>.

115. Hadley G, Derry S, Moore RA, Wiffen PJ. Transdermal fentanyl for cancer pain. Cochrane Database Syst Rev 2013;10:CD010270. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24096644>.

116. Wang DD, Ma TT, Zhu HD, Peng CB. Transdermal fentanyl for cancer pain: Trial sequential analysis of 3406 patients from 35 randomized controlled trials. J Cancer Res Ther 2018;14:S14-s21. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29578144>.

117. Kornick CA, Santiago-Palma J, Khojainova N, et al. A safe and effective method for converting cancer patients from intravenous to transdermal fentanyl. Cancer 2001;92:3056-3061. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11753984>.

118. Portenoy RK, Taylor D, Messina J, Tremmel L. A randomized, placebo-controlled study of fentanyl buccal tablet for breakthrough pain in

opioid-treated patients with cancer. Clin J Pain 2006;22:805-811. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17057563>.

119. Weinstein SM, Messina J, Xie F. Fentanyl buccal tablet for the treatment of breakthrough pain in opioid-tolerant patients with chronic cancer pain: A long-term, open-label safety study. Cancer 2009;115:2571-2579. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19373888>.

120. Kleeberg UR, Filbet M, Zeppetella G. Fentanyl buccal tablet for breakthrough cancer pain: why titrate? Pain Pract 2011;11:185-190. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20807349>.

121. Srinivasan V, Wielbo D, Tebbett IR. Analgesic effects of codeine-6-glucuronide after intravenous administration. Eur J Pain 1997;1:185-190. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15102399>.

122. Kirchheiner J, Schmidt H, Tzvetkov M, et al. Pharmacokinetics of codeine and its metabolite morphine in ultra-rapid metabolizers due to CYP2D6 duplication. Pharmacogenomics J 2007;7:257-265. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16819548>.

123. Murray A, Hagen N. Hydromorphone. J Pain Symptom Manage. 2005;29(5 suppl):S57-S66. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/15907647>.

124. Thwaites D, McCann S, Broderick P. Hydromorphone neuroexcitation. J Palliat Med 2004;7:545-550. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15353098>.

125. Wright AW, Mather LE, Smith MT. Hydromorphone-3-glucuronide: a more potent neuro-excitant than its structural analogue, morphine-3-glucuronide. Life Sci 2001;69:409-420. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11459432>.

126. Han HS, Lee KH, Lee KH, et al. A prospective, open-label, multicenter study of the clinical efficacy of extended-release hydromorphone in treating cancer pain inadequately controlled by other analgesics. Support Care Cancer 2014;22:741-750. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24203087>.

**NCCN Guidelines Version 2.2022**
Adult Cancer Pain

127. Yu S, Shen W, Yu L, et al. Safety and efficacy of once-daily hydromorphone extended-release versus twice-daily oxycodone hydrochloride controlled-release in chinese patients with cancer pain: a phase 3, randomized, double-blind, multicenter study. *J Pain* 2014;15:835-844. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24846822>.

128. Inoue S, Saito Y, Tsuneto S, et al. A randomized, double-blind, non-inferiority study of hydromorphone hydrochloride immediate-release tablets versus oxycodone hydrochloride immediate-release powder for cancer pain: efficacy and safety in Japanese cancer patients. *Jpn J Clin Oncol* 2018;48:542-547. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29659913>.

129. Bao YJ, Hou W, Kong XY, et al. Hydromorphone for cancer pain. *Cochrane Database Syst Rev* 2016;10:Cd011108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27727452>.

130. Davis MP, Varga J, Dickerson D, et al. Normal-release and controlled-release oxycodone: pharmacokinetics, pharmacodynamics, and controversy. *Support Care Cancer* 2003;11:84-92. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12560936>.

131. Ordonez Gallego A, Gonzalez Baron M, Espinosa Arranz E. Oxycodone: a pharmacological and clinical review. *Clin Transl Oncol* 2007;9:298-307. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17525040>.

132. Gabrail NY, Dvergsten C, Ahdieh H. Establishing the dosage equivalency of oxymorphone extended release and oxycodone controlled release in patients with cancer pain: a randomized controlled study. *Curr Med Res Opin* 2004;20:911-918. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15200750>.

133. Schmidt-Hansen M, Bennett MI, Arnold S, et al. Oxycodone for cancer-related pain. *Cochrane Database Syst Rev* 2017;8:Cd003870. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28829910>.

134. Schmidt-Hansen M, Bennett MI, Arnold S, et al. Efficacy, tolerability and acceptability of oxycodone for cancer-related pain in adults: an

updated Cochrane systematic review. *BMJ Support Palliat Care* 2018;8:117-128. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29331953>.

135. Guo KK, Deng CQ, Lu GJ, Zhao GL. Comparison of analgesic effect of oxycodone and morphine on patients with moderate and advanced cancer pain: a meta-analysis. *BMC Anesthesiol* 2018;18:132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30249205>.

136. Ahmedzai SH, Nauck F, Bar-Sela G, et al. A randomized, double-blind, active-controlled, double-dummy, parallel-group study to determine the safety and efficacy of oxycodone/naloxone prolonged-release tablets in patients with moderate/severe, chronic cancer pain. *Palliat Med* 2012;26:50-60. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21937568>.

137. Ahmedzai SH, Leppert W, Janecki M, et al. Long-term safety and efficacy of oxycodone/naloxone prolonged-release tablets in patients with moderate-to-severe chronic cancer pain. *Support Care Cancer* 2015;23:823-830. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25218610>.

138. Davis MP, Homsy J. The importance of cytochrome P450 monooxygenase CYP2D6 in palliative medicine. *Support Care Cancer* 2001;9:442-451. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11585271>.

139. McPherson ML, Walker KA, Davis MP, et al. Safe and appropriate use of methadone in hospice and palliative care: Expert consensus white paper. *J Pain Symptom Manage* 2019;57:635-645.e634. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30578934>.

140. Moksnes K, Dale O, Rosland JH, et al. How to switch from morphine or oxycodone to methadone in cancer patients? a randomised clinical phase II trial. *Eur J Cancer* 2011;47:2463-2470. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21775131>.

141. McLean S, Twomey F. Methods of rotation from another strong opioid to methadone for the management of cancer pain: a systematic review of



NCCN Guidelines Version 2.2022

Adult Cancer Pain

the available evidence. *J Pain Symptom Manage* 2015;50:248-259 e241. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25896106>.

142. Nicholson AB, Watson GR, Derry S, Wiffen PJ. Methadone for cancer pain. *Cochrane Database Syst Rev* 2017;2:Cd003971. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28177515>.

143. Parsons HA, de la Cruz M, El Osta B, et al. Methadone initiation and rotation in the outpatient setting for patients with cancer pain. *Cancer* 2010;116:520-528. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19924788>.

144. Courtemanche F, Dao D, Gagne F, et al. Methadone as a coanalgesic for palliative care cancer patients. *J Palliat Med* 2016;19:972-978. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27399839>.

145. Furst P, Lundstrom S, Klepstad P, et al. Improved pain control in terminally ill cancer patients by introducing low-dose oral methadone in addition to ongoing opioid treatment. *J Palliat Med* 2018;21:177-181. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28792784>.

146. Krantz MJ, Lewkowicz L, Hays H, et al. Torsade de pointes associated with very-high-dose methadone. *Ann Intern Med* 2002;137:501-504. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12230351>.

147. Krantz MJ, Kutinsky IB, Robertson AD, Mehler PS. Dose-related effects of methadone on QT prolongation in a series of patients with torsade de pointes. *Pharmacotherapy* 2003;23:802-805. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12820821>.

148. Kornick CA, Kilborn MJ, Santiago-Palma J, et al. QTc interval prolongation associated with intravenous methadone. *Pain* 2003;105:499-506. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14527710>.

149. Reddy S, Hui D, El Osta B, et al. The effect of oral methadone on the QTc interval in advanced cancer patients: a prospective pilot study. *J Palliat Med* 2010;13:33-38. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19824814>.

150. Chou R, Cruciani RA, Fiellin DA, et al. Methadone safety: a clinical practice guideline from the American Pain Society and College on Problems of Drug Dependence, in collaboration with the Heart Rhythm Society. *J Pain* 2014;15:321-337. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24685458>.

151. McNulty JP. Can levorphanol be used like methadone for intractable refractory pain? *J Palliat Med* 2007;10:293-296. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17472497>.

152. Atkinson TJ, Fudin J, Pandula A, Mirza M. Medication pain management in the elderly: unique and underutilized analgesic treatment options. *Clin Ther* 2013;35:1669-1689. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24161287>.

153. Rowbotham MC, Twilling L, Davies PS, et al. Oral opioid therapy for chronic peripheral and central neuropathic pain. *N Engl J Med* 2003;348:1223-1232. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12660386>.

154. Beakley BD, Kaye AM, Kaye AD. Tramadol, pharmacology, side effects, and serotonin syndrome: a review. *Pain Physician* 2015;18:395-400. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26218943>.

155. Grond S, Sablotzki A. Clinical pharmacology of tramadol. *Clin Pharmacokinet* 2004;43:879-923. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15509185>.

156. Grond S, Radbruch L, Meuser T, et al. High-dose tramadol in comparison to low-dose morphine for cancer pain relief. *J Pain Symptom Manage* 1999;18:174-179. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10517038>.

157. Rodriguez RF, Bravo LE, Castro F, et al. Incidence of weak opioids adverse events in the management of cancer pain: a double-blind comparative trial. *J Palliat Med* 2007;10:56-60. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17298254>.



158. Wiffen PJ, Derry S, Moore RA. Tramadol with or without paracetamol (acetaminophen) for cancer pain. *Cochrane Database Syst Rev* 2017;5:Cd012508. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28510996>.
159. Wade WE, Spruill WJ. Tapentadol hydrochloride: a centrally acting oral analgesic. *Clin Ther* 2009;31:2804-2818. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20110020>.
160. Hartrick CT, Rodriguez Hernandez JR. Tapentadol for pain: a treatment evaluation. *Expert Opin Pharmacother* 2012;13:283-286. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22192161>.
161. Afilalo M, Etropolski MS, Kuperwasser B, et al. Efficacy and safety of Tapentadol extended release compared with oxycodone controlled release for the management of moderate to severe chronic pain related to osteoarthritis of the knee: a randomized, double-blind, placebo- and active-controlled phase III study. *Clin Drug Investig* 2010;30:489-505. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20586515>.
162. Buynak R, Shapiro DY, Okamoto A, et al. Efficacy and safety of tapentadol extended release for the management of chronic low back pain: results of a prospective, randomized, double-blind, placebo- and active-controlled Phase III study. *Expert Opin Pharmacother* 2010;11:1787-1804. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20578811>.
163. Schwartz S, Etropolski M, Shapiro DY, et al. Safety and efficacy of tapentadol ER in patients with painful diabetic peripheral neuropathy: results of a randomized-withdrawal, placebo-controlled trial. *Curr Med Res Opin* 2011;27:151-162. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21162697>.
164. Mercadante S, Porzio G, Ferrera P, et al. Tapentadol in cancer pain management: a prospective open-label study. *Curr Med Res Opin* 2012;28:1775-1779. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23057488>.
165. Mercadante S, Porzio G, Adile C, et al. Tapentadol at medium to high doses in patients previously receiving strong opioids for the management of cancer pain. *Curr Med Res Opin* 2014;30:2063-2068. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24926734>.
166. Naing C, Aung K, Raclou V, Yeoh PN. Safety and efficacy of transdermal buprenorphine for the relief of cancer pain. *J Cancer Res Clin Oncol* 2013;139:1963-1970. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23922192>.
167. Pergolizzi JV, Jr., Mercadante S, Echaburu AV, et al. The role of transdermal buprenorphine in the treatment of cancer pain: an expert panel consensus. *Curr Med Res Opin* 2009;25:1517-1528. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19435402>.
168. Deandrea S, Corli O, Moschetti I, Apolone G. Managing severe cancer pain: the role of transdermal buprenorphine: a systematic review. *Ther Clin Risk Manag* 2009;5:707-718. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19774212>.
169. Melilli G, Samolsky Dekel BG, Frenquelli C, et al. Transdermal opioids for cancer pain control in patients with renal impairment. *J Opioid Manag* 2014;10:85-93. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24715663>.
170. Lundorff L, Sjogren P, Hansen OB, et al. Switching from high doses of pure mu-opioid agonists to transdermal buprenorphine in patients with cancer: a feasibility study. *J Opioid Manag* 2013;9:255-262. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24353018>.
171. Johnson RE, Fudala PJ, Payne R. Buprenorphine: considerations for pain management. *J Pain Symptom Manage* 2005;29:297-326. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15781180>.
172. Bell RF, Eccleston C, Kalso EA. Ketamine as an adjuvant to opioids for cancer pain. *Cochrane Database Syst Rev* 2017;6:Cd003351. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28657160>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

173. Hardy J, Quinn S, Fazekas B, et al. Randomized, double-blind, placebo-controlled study to assess the efficacy and toxicity of subcutaneous ketamine in the management of cancer pain. *J Clin Oncol* 2012;30:3611-3617. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22965960>.

174. Bredlau AL, Thakur R, Korones DN, Dworkin RH. Ketamine for pain in adults and children with cancer: a systematic review and synthesis of the literature. *Pain Med* 2013;14:1505-1517. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23915253>.

175. DeWilde KE, Levitch CF, Murrugh JW, et al. The promise of ketamine for treatment-resistant depression: current evidence and future directions. *Ann N Y Acad Sci* 2015;1345:47-58. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25649308>.

176. Fond G, Loundou A, Rabu C, et al. Ketamine administration in depressive disorders: a systematic review and meta-analysis. *Psychopharmacology (Berl)* 2014;231:3663-3676. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25038867>.

177. Lee EE, Della Selva MP, Liu A, Himelhoch S. Ketamine as a novel treatment for major depressive disorder and bipolar depression: a systematic review and quantitative meta-analysis. *Gen Hosp Psychiatry* 2015;37:178-184. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25698228>.

178. Sharma S, Rajagopal MR, Palat G, et al. A phase II pilot study to evaluate use of intravenous lidocaine for opioid-refractory pain in cancer patients. *J Pain Symptom Manage* 2009;37:85-93. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18599258>.

179. Jendoubi A, Naceur IB, Bouzouita A, et al. A comparison between intravenous lidocaine and ketamine on acute and chronic pain after open nephrectomy: A prospective, double-blind, randomized, placebo-controlled study. *Saudi J Anaesth* 2017;11:177-184. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28442956>.

180. Ferrini R, Paice JA. How to initiate and monitor infusional lidocaine for severe and/or neuropathic pain. *J Support Oncol* 2004;2:90-94. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15330376>.

181. Carroll I. Intravenous lidocaine for neuropathic pain: diagnostic utility and therapeutic efficacy. *Curr Pain Headache Rep* 2007;11:20-24. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17214917>.

182. Lee JT, Sanderson CR, Xuan W, Agar M. Lidocaine for cancer pain in adults: A systematic review and meta-analysis. *J Palliat Med* 2019;22:326-334. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30614748>.

183. Bruera E, Kim HN. Cancer pain. *JAMA* 2003;290:2476-2479. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14612485>.

184. Stevens RA, Ghazi SM. Routes of opioid analgesic therapy in the management of cancer pain. *Cancer Control* 2000;7:132-141. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10783817>.

185. Harris JT, Suresh Kumar K, Rajagopal MR. Intravenous morphine for rapid control of severe cancer pain. *Palliat Med* 2003;17:248-256. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12725478>.

186. Nijland L, Schmidt P, Frosch M, et al. Subcutaneous or intravenous opioid administration by patient-controlled analgesia in cancer pain: a systematic literature review. *Support Care Cancer* 2019;27:33-42. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30056529>.

187. Davies A, Buchanan A, Zeppetella G, et al. Breakthrough cancer pain: an observational study of 1000 European oncology patients. *J Pain Symptom Manage* 2013;46:619-628. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23523361>.

188. Zeppetella G, Davies AN. Opioids for the management of breakthrough pain in cancer patients. *Cochrane Database Syst Rev* 2013;10:CD004311. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24142465>.



189. Mercadante S. Pharmacotherapy for breakthrough cancer pain. *Drugs* 2012;72:181-190. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22233484>.

190. Jandhyala R, Fullarton JR, Bennett MI. Efficacy of rapid-onset oral fentanyl formulations vs. oral morphine for cancer-related breakthrough pain: a meta-analysis of comparative trials. *J Pain Symptom Manage* 2013;46:573-580. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23380337>.

191. Zeppetella G, Davies A, Eijgelshoven I, Jansen JP. A network meta-analysis of the efficacy of opioid analgesics for the management of breakthrough cancer pain episodes. *J Pain Symptom Manage* 2014;47:772-785 e775. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23981487>.

192. VA/DoD clinical practice guideline for opioid therapy for chronic pain. Version 3.0. 2017. Available at: <https://www.healthquality.va.gov/guidelines/Pain/cot/VADoDOTCPG022717.pdf>. Accessed April 06, 2021.

193. Butler SF, Budman SH, Fernandez KC, et al. Development and validation of the Current Opioid Misuse Measure. *Pain* 2007;130:144-156. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17493754>.

194. Meltzer EC, Rybin D, Saitz R, et al. Identifying prescription opioid use disorder in primary care: diagnostic characteristics of the Current Opioid Misuse Measure (COMM). *Pain* 2011;152:397-402. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21177035>.

195. McNicol E, Horowicz-Mehler N, Fisk RA, et al. Management of opioid side effects in cancer-related and chronic noncancer pain: a systematic review. *J Pain* 2003;4:231-256. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14622694>.

196. Mercadante S. Comments on Wang et al., *PAIN*, 67 (1996) 407-416. *Pain* 1998;74:106-107. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9514568>.

197. Mercadante S. Pathophysiology and treatment of opioid-related myoclonus in cancer patients. *Pain* 1998;74:5-9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9514554>.

198. Wilson RK, Weissman DE. Neuroexcitatory effects of opioids: patient assessment #57. *J Palliat Med* 2004;7:579. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15353102>.

199. Moryl N, Carver, A, Foley, KM. , ed Pain and palliation. In: Holland JF, Frei E, eds. *Cancer Medicine*. Vol. 17th ed. Hamilton, ON: BC Decker Inc; 2006:1113-1124.

200. Moryl N, Obbens EA, Ozigbo OH, Kris MG. Analgesic effect of gefitinib in the treatment of non-small cell lung cancer. *J Support Oncol* 2006;4:111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16553135>.

201. Rajagopal A, Vassilopoulou-Sellin R, Palmer JL, et al. Symptomatic hypogonadism in male survivors of cancer with chronic exposure to opioids. *Cancer* 2004;100:851-858. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14770444>.

202. Boettger S, Breitbart W. Atypical antipsychotics in the management of delirium: a review of the empirical literature. *Palliat Support Care* 2005;3:227-237. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16594462>.

203. Breitbart W, Marotta R, Platt MM, et al. A double-blind trial of haloperidol, chlorpromazine, and lorazepam in the treatment of delirium in hospitalized AIDS patients. *Am J Psychiatry* 1996;153:231-237. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8561204>.

204. Bruera E, Belzile M, Neumann C, et al. A double-blind, crossover study of controlled-release metoclopramide and placebo for the chronic nausea and dyspepsia of advanced cancer. *J Pain Symptom Manage* 2000;19:427-435. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10908823>.



205. Challoner KR, McCarron MM, Newton EJ. Pentazocine (Talwin) intoxication: report of 57 cases. *J Emerg Med* 1990;8:67-74. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2351801>.

206. Katcher J, Walsh D. Opioid-induced itching: morphine sulfate and hydromorphone hydrochloride. *J Pain Symptom Manage* 1999;17:70-72. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9919868>.

207. Marinella MA. Acute colonic pseudo-obstruction complicated by cecal perforation in a patient with Parkinson's disease. *South Med J* 1997;90:1023-1026. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9347813>.

208. Reissig JE, Rybarczyk AM. Pharmacologic treatment of opioid-induced sedation in chronic pain. *Ann Pharmacother* 2005;39:727-731. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15755795>.

209. Tarcatu D, Tamasdan C, Moryl N, Obbens E. Are we still scratching the surface? A case of intractable pruritus following systemic opioid analgesia. *J Opioid Manag* 2007;3:167-170. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18027543>.

210. Prommer E. Modafinil: is it ready for prime time? *J Opioid Manag* 2006;2:130-136. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17319446>.

211. Hawley PH, Byeon JJ. A comparison of sennosides-based bowel protocols with and without docusate in hospitalized patients with cancer. *J Palliat Med* 2008;11:575-581. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18454610>.

212. Tarumi Y, Wilson MP, Szafran O, Spooner GR. Randomized, double-blind, placebo-controlled trial of oral docusate in the management of constipation in hospice patients. *J Pain Symptom Manage* 2013;45:2-13. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22889861>.

213. Rauck R, Slatkin NE, Stambler N, et al. Randomized, Double-Blind Trial of Oral Methylnaltrexone for the Treatment of Opioid-Induced Constipation in Patients with Chronic Noncancer Pain. *Pain Pract*

2017;17:820-828. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27860208>.

214. Michna E, Blonsky ER, Schulman S, et al. Subcutaneous methylnaltrexone for treatment of opioid-induced constipation in patients with chronic, nonmalignant pain: a randomized controlled study. *J Pain* 2011;12:554-562. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21429809>.

215. Portenoy RK, Thomas J, Moehl Boatwright ML, et al. Subcutaneous methylnaltrexone for the treatment of opioid-induced constipation in patients with advanced illness: a double-blind, randomized, parallel group, dose-ranging study. *J Pain Symptom Manage* 2008;35:458-468. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18440447>.

216. Chappell D, Rehm M, Conzen P. Methylnaltrexone for opioid-induced constipation in advanced illness. *N Engl J Med* 2008;359:1071; author reply 1071. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18777614>.

217. Sanz Rubiales A, del Valle Rivero ML. Methylnaltrexone for opioid-induced constipation in advanced illness. *N Engl J Med* 2008;359:1070-1071; author reply 1071. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18768955>.

218. Thomas J, Karver S, Cooney GA, et al. Methylnaltrexone for opioid-induced constipation in advanced illness. *N Engl J Med* 2008;358:2332-2343. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18509120>.

219. Chey WD, Webster L, Sostek M, et al. Naloxegol for opioid-induced constipation in patients with noncancer pain. *N Engl J Med* 2014;370:2387-2396. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24896818>.

220. Katakami N, Harada T, Murata T, et al. Randomized phase III and extension studies of naldemedine in patients with opioid-induced constipation and cancer. *J Clin Oncol* 2017;35:3859-3866. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28968171>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

221. Nee J, Zakari M, Sugarman MA, et al. Efficacy of treatments for opioid-induced constipation: systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2018;16:1569-1584.e1562. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29374616>.
222. Cryer B, Katz S, Vallejo R, et al. A randomized study of lubiprostone for opioid-induced constipation in patients with chronic noncancer pain. *Pain Med* 2014;15:1825-1834. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24716835>.
223. Jamal MM, Adams AB, Jansen JP, Webster LR. A randomized, placebo-controlled trial of lubiprostone for opioid-induced constipation in chronic noncancer pain. *Am J Gastroenterol* 2015;110:725-732. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25916220>.
224. Mackey AC, Green L, Greene P, Avigan M. Methylnaltrexone and gastrointestinal perforation. *J Pain Symptom Manage* 2010;40:e1-3. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20619194>.
225. Crockett SD, Greer KB, Heidelbaugh JJ, et al. American Gastroenterological Association Institute Guideline on the Medical Management of Opioid-Induced Constipation. *Gastroenterology* 2019;156:218-226. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30340754>.
226. Kaneishi K, Kawabata M, Morita T. Olanzapine for the relief of nausea in patients with advanced cancer and incomplete bowel obstruction. *J Pain Symptom Manage* 2012;44:604-607. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22771132>.
227. Navari RM, Nagy CK, Gray SE. The use of olanzapine versus metoclopramide for the treatment of breakthrough chemotherapy-induced nausea and vomiting in patients receiving highly emetogenic chemotherapy. *Support Care Cancer* 2013;21:1655-1663. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23314603>.
228. Bruera E, Seifert L, Watanabe S, et al. Chronic nausea in advanced cancer patients: a retrospective assessment of a metoclopramide-based antiemetic regimen. *J Pain Symptom Manage* 1996;11:147-153. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8851371>.
229. Sande TA, Laird BJA, Fallon MT. The management of opioid-induced nausea and vomiting in patients with cancer: A systematic review. *J Palliat Med* 2019;22:90-97. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30239277>.
230. Herman TS, Einhorn LH, Jones SE, et al. Superiority of nabilone over prochlorperazine as an antiemetic in patients receiving cancer chemotherapy. *N Engl J Med* 1979;300:1295-1297. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/375088>.
231. Steele N, Gralla RJ, Braun DW, Jr., Young CW. Double-blind comparison of the antiemetic effects of nabilone and prochlorperazine on chemotherapy-induced emesis. *Cancer Treat Rep* 1980;64:219-224. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/6250699>.
232. Sallan SE, Cronin C, Zelen M, Zinberg NE. Antiemetics in patients receiving chemotherapy for cancer: a randomized comparison of delta-9-tetrahydrocannabinol and prochlorperazine. *N Engl J Med* 1980;302:135-138. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/6985702>.
233. Sallan SE, Zinberg NE, Frei E, 3rd. Antiemetic effect of delta-9-tetrahydrocannabinol in patients receiving cancer chemotherapy. *N Engl J Med* 1975;293:795-797. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1099449>.
234. Chamberlain JM, Klein BL. A comprehensive review of naloxone for the emergency physician. *Am J Emerg Med* 1994;12:650-660. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7945608>.
235. Gagnon P, Allard P, Masse B, DeSerres M. Delirium in terminal cancer: a prospective study using daily screening, early diagnosis, and continuous monitoring. *J Pain Symptom Manage* 2000;19:412-426. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10908822>.
236. Agar MR, Lawlor PG, Quinn S, et al. Efficacy of oral risperidone, haloperidol, or placebo for symptoms of delirium among patients in



palliative care: A randomized clinical trial. JAMA Intern Med 2017;177:34-42. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27918778>.

237. Hirst JM, Vaughan CL, Irwin SA. Delirium: Use antipsychotics when appropriate and appropriately. J Palliat Med 2017;20:799. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28410450>.

238. Hui D, Frisbee-Hume S, Wilson A, et al. Effect of lorazepam with haloperidol vs haloperidol alone on agitated delirium in patients with advanced cancer receiving palliative care: A randomized clinical trial. Jama 2017;318:1047-1056. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28975307>.

239. Bruera E, Macmillan K, Hanson J, MacDonald RN. The cognitive effects of the administration of narcotic analgesics in patients with cancer pain. Pain 1989;39:13-16. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2812850>.

240. Vardy ER, Teodorczuk A, Yarnall AJ. Review of delirium in patients with Parkinson's disease. J Neurol 2015;262:2401-2410. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25957635>.

241. Pasero C. Assessment of sedation during opioid administration for pain management. J Perianesth Nurs 2009;24:186-190. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19500754>.

242. Nisbet AT, Mooney-Cotter F. Comparison of selected sedation scales for reporting opioid-induced sedation assessment. Pain Manag Nurs 2009;10:154-164. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19706353>.

243. U.S. Food and Drug Administration. FDA warns about serious risks and death when combining opioid pain or cough medicines with benzodiazepines; requires its strongest warning. 2016. Available at: <https://www.fda.gov/downloads/drugs/drugsafety/ucm518672.pdf>. Accessed April 06, 2021.

244. U.S. Food and Drug Administration. Prescribing information. Naloxone hydrochloride nasal spray. 2020. Available at:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/208411Orig1s004lbl.pdf. Accessed April 06, 2021.

245. Coffin PO, Behar E, Rowe C, et al. Nonrandomized intervention study of naloxone coprescription for primary care patients receiving long-term opioid therapy for pain. Ann Intern Med 2016;165:245-252. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27366987>.

246. Slatkin NE. Opioid switching and rotation in primary care: implementation and clinical utility. Curr Med Res Opin 2009;25:2133-2150. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19601703>.

247. Vissers KC, Besse K, Hans G, et al. Opioid rotation in the management of chronic pain: where is the evidence? Pain Pract 2010;10:85-93. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20070552>.

248. Schuster M, Bayer O, Heid F, Laufenberg-Feldmann R. Opioid rotation in cancer pain treatment. Dtsch Arztebl Int 2018;115:135-142. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29563006>.

249. Corli O, Roberto A, Corsi N, et al. Opioid switching and variability in response in pain cancer patients. Support Care Cancer 2019;27:2321-2327. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30357556>.

250. Reddy A, Yennurajalingam S, Desai H, et al. The opioid rotation ratio of hydrocodone to strong opioids in cancer patients. Oncologist 2014;19:1186-1193. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25342316>.

251. Davis MP, McPherson ML. Tabling hydromorphone: do we have it right? J Palliat Med 2010;13:365-366. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20233019>.

252. Reddy A, Tayjasanant S, Haider A, et al. The opioid rotation ratio of strong opioids to transdermal fentanyl in cancer patients. Cancer 2016;122:149-156. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26451687>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

253. Reddy A, Yennurajalingam S, Reddy S, et al. The opioid rotation ratio from transdermal fentanyl to "strong" opioids in patients with cancer pain. *J Pain Symptom Manage* 2016;51:1040-1045. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26826675>.
254. Reddy A, Vidal M, Stephen S, et al. The conversion ratio from intravenous hydromorphone to oral opioids in cancer patients. *J Pain Symptom Manage* 2017;54:280-288. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28711751>.
255. Hedegaard H, Miniño AM, Warner M. Drug Overdose Deaths in the United States, 1999-2018. *NCHS Data Brief* 2020:1-8. Available at: <https://pubmed.ncbi.nlm.nih.gov/32487285/>.
256. Dart RC, Surratt HL, Cicero TJ, et al. Trends in opioid analgesic abuse and mortality in the United States. *N Engl J Med* 2015;372:241-248. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25587948>.
257. U.S. Food and Drug Administration. Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS). 2018. Available at: <https://www.fda.gov/drugs/drugsafety/informationbydrugclass/ucm163647.htm>. Accessed April 06, 2021.
258. FDA takes important steps to encourage appropriate and rational prescribing of opioids through final approval of new safety measures governing the use of immediate-release opioid analgesic medications; 2018. Available at: <https://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm620935.htm>.
259. FDA Education Blueprint for Health Care Providers Involved in the Treatment and Monitoring of Patients with Pain; 2018. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/remis/Opioid_analgesic_2018_09_18_FDA_Blueprint.pdf.
260. U.S. Food and Drug Administration. Approved Risk Evaluation and Mitigation Strategies (REMS). Available at: <http://www.accessdata.fda.gov/scripts/cder/remis/index.cfm>. Accessed April 06, 2021.
261. Barclay JS, Owens JE, Blackhall LJ. Screening for substance abuse risk in cancer patients using the Opioid Risk Tool and urine drug screen. *Support Care Cancer* 2014;22:1883-1888. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24563103>.
262. Arthur JA. Urine drug testing in cancer pain management. *Oncologist* 2020;25:99-104. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32043770>.
263. Fallon MT. Neuropathic pain in cancer. *Br J Anaesth* 2013;111:105-111. Available at:
264. Saarto T, Wiffen PJ. Antidepressants for neuropathic pain. *Cochrane Database Syst Rev* 2005:CD005454. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16034979>.
265. Saarto T, Wiffen PJ. Antidepressants for neuropathic pain: a Cochrane review. *J Neurol Neurosurg Psychiatry* 2010;81:1372-1373. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20543189>.
266. Finnerup NB, Sindrup SH, Jensen TS. The evidence for pharmacological treatment of neuropathic pain. *Pain* 2010;150:573-581. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20705215>.
267. Caraceni A, Zecca E, Bonezzi C, et al. Gabapentin for neuropathic cancer pain: a randomized controlled trial from the Gabapentin Cancer Pain Study Group. *J Clin Oncol* 2004;22:2909-2917. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15254060>.
268. Bennett MI. Effectiveness of antiepileptic or antidepressant drugs when added to opioids for cancer pain: systematic review. *Palliat Med* 2011;25:553-559. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20671006>.
269. Kane CM, Mulvey MR, Wright S, et al. Opioids combined with antidepressants or antiepileptic drugs for cancer pain: Systematic review and meta-analysis. *Palliat Med* 2018;32:276-286. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28604172>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

270. Fleming JA, O'Connor BD. Use of lidocaine patches for neuropathic pain in a comprehensive cancer centre. *Pain Res Manag* 2009;14:381-388. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19862373>.

271. Gammaitoni AR, Alvarez NA, Galer BS. Safety and tolerability of the lidocaine patch 5%, a targeted peripheral analgesic: a review of the literature. *J Clin Pharmacol* 2003;43:111-117. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12616661>.

272. Garzon-Rodriguez C, Casals Merchan M, Calsina-Berna A, et al. Lidocaine 5 % patches as an effective short-term co-analgesic in cancer pain. Preliminary results. *Support Care Cancer* 2013;21:3153-3158. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24000041>.

273. Liu Z, Xu Y, Liu ZL, et al. Combined application of diclofenac and celecoxib with an opioid yields superior efficacy in metastatic bone cancer pain: a randomized controlled trial. *Int J Clin Oncol* 2017;22:980-985. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28484877>.

274. Sima L, Fang WX, Wu XM, Li F. Efficacy of oxycodone/paracetamol for patients with bone-cancer pain: a multicenter, randomized, double-blinded, placebo-controlled trial. *J Clin Pharm Ther* 2012;37:27-31. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21208247>.

275. Chow E, Meyer RM, Ding K, et al. Dexamethasone in the prophylaxis of radiation-induced pain flare after palliative radiotherapy for bone metastases: a double-blind, randomised placebo-controlled, phase 3 trial. *Lancet Oncol* 2015;16:1463-1472. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26489389>.

276. Body JJ, Diel IJ, Lichinitzer M, et al. Oral ibandronate reduces the risk of skeletal complications in breast cancer patients with metastatic bone disease: results from two randomised, placebo-controlled phase III studies. *Br J Cancer* 2004;90:1133-1137. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15026791>.

277. Body JJ, Diel IJ, Bell R, et al. Oral ibandronate improves bone pain and preserves quality of life in patients with skeletal metastases due to

breast cancer. *Pain* 2004;111:306-312. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15363874>.

278. Cleeland CS, Body JJ, Stopeck A, et al. Pain outcomes in patients with advanced breast cancer and bone metastases: Results from a randomized, double-blind study of denosumab and zoledronic acid. *Cancer* 2013;119:832-838. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22951813>.

279. Rosen LS, Gordon D, Kaminski M, et al. Zoledronic acid versus pamidronate in the treatment of skeletal metastases in patients with breast cancer or osteolytic lesions of multiple myeloma: a phase III, double-blind, comparative trial. *Cancer J* 2001;7:377-387. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11693896>.

280. Wardley A, Davidson N, Barrett-Lee P, et al. Zoledronic acid significantly improves pain scores and quality of life in breast cancer patients with bone metastases: a randomised, crossover study of community vs hospital bisphosphonate administration. *Br J Cancer* 2005;92:1869-1876. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15870721>.

281. Vadhan-Raj S, von Moos R, Fallowfield LJ, et al. Clinical benefit in patients with metastatic bone disease: results of a phase 3 study of denosumab versus zoledronic acid. *Ann Oncol* 2012;23:3045-3051. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22851406>.

282. Martin M, Bell R, Bourgeois H, et al. Bone-related complications and quality of life in advanced breast cancer: results from a randomized phase III trial of denosumab versus zoledronic acid. *Clin Cancer Res* 2012;18:4841-4849. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22893628>.

283. Van Poznak C, Somerfield MR, Barlow WE, et al. Role of Bone-Modifying Agents in Metastatic Breast Cancer: An American Society of Clinical Oncology-Cancer Care Ontario Focused Guideline Update. *J Clin Oncol* 2017;35:3978-3986. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29035643>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

284. Malviya A, Gerrand C. Evidence for orthopaedic surgery in the treatment of metastatic bone disease of the extremities: a review article. *Palliat Med* 2012;26:788-796. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21930647>.

285. Dupuy DE, Liu D, Hartfeil D, et al. Percutaneous radiofrequency ablation of painful osseous metastases: a multicenter American College of Radiology Imaging Network trial. *Cancer* 2010;116:989-997. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20041484>.

286. Lutz S, Berk L, Chang E, et al. Palliative radiotherapy for bone metastases: an ASTRO evidence-based guideline. *Int J Radiat Oncol Biol Phys* 2011;79:965-976. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21277118>.

287. Goetz MP, Callstrom MR, Charboneau JW, et al. Percutaneous image-guided radiofrequency ablation of painful metastases involving bone: a multicenter study. *J Clin Oncol* 2004;22:300-306. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14722039>.

288. Kashima M, Yamakado K, Takaki H, et al. Radiofrequency ablation for the treatment of bone metastases from hepatocellular carcinoma. *AJR Am J Roentgenol* 2010;194:536-541. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20093621>.

289. Li C, Zhang W, Fan W, et al. Noninvasive treatment of malignant bone tumors using high-intensity focused ultrasound. *Cancer* 2010;116:3934-3942. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20564113>.

290. Napoli A, Anzidei M, Marincola BC, et al. Primary pain palliation and local tumor control in bone metastases treated with magnetic resonance-guided focused ultrasound. *Invest Radiol* 2013;48:351-358. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/?term=23571832>.

291. Liberman B, Gianfelice D, Inbar Y, et al. Pain palliation in patients with bone metastases using MR-guided focused ultrasound surgery: a multicenter study. *Ann Surg Oncol* 2009;16:140-146. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/?term=19002530>.

292. Silver JK, Gilchrist LS. Cancer rehabilitation with a focus on evidence-based outpatient physical and occupational therapy interventions. *Am J Phys Med Rehabil* 2011;90:S5-15. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21765263>.

293. Silver JK, Baima J, Mayer RS. Impairment-driven cancer rehabilitation: an essential component of quality care and survivorship. *CA Cancer J Clin* 2013;63:295-317. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23856764>.

294. Jones L, Fitzgerald G, Leurent B, et al. Rehabilitation in advanced, progressive, recurrent cancer: a randomized controlled trial. *J Pain Symptom Manage* 2013;46:315-325.e313. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23182307>.

295. Eilers J, Harris D, Henry K, Johnson LA. Evidence-based interventions for cancer treatment-related mucositis: putting evidence into practice. *Clin J Oncol Nurs* 2014;18 Suppl:80-96. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25427611>.

296. Oncology Nursing Society. Mucositis. 2019. Available at: <https://www.ons.org/pep/mucositis>. Accessed February 1, 2021.

297. Elad S, Cheng KKF, Lalla RV, et al. MASCC/ISOO clinical practice guidelines for the management of mucositis secondary to cancer therapy. *Cancer* 2020;126:4423-4431. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32786044>.

298. Peterson DE, Boers-Doets CB, Bensadoun RJ, et al. Management of oral and gastrointestinal mucosal injury: ESMO Clinical Practice Guidelines for diagnosis, treatment, and follow-up. *Ann Oncol* 2015;26 Suppl 5:v139-151. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26142468>.

299. Riley P, Glenny AM, Worthington HV, et al. Interventions for preventing oral mucositis in patients with cancer receiving treatment: oral cryotherapy. *Cochrane Database Syst Rev* 2015: Cd011552. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26695736>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

300. Idayu Mat Nawi R, Lei Chui P, Wan Ishak WZ, Hsien Chan CM. Oral cryotherapy: Prevention of oral mucositis and pain among patients with colorectal cancer undergoing chemotherapy. *Clin J Oncol Nurs* 2018;22:555-560. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30239519>.
301. Kataoka T, Kiyota N, Shimada T, et al. Randomized trial of standard pain control with or without gabapentin for pain related to radiation-induced mucositis in head and neck cancer. *Auris Nasus Larynx* 2016;43:677-684. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26992271>.
302. McGuire DB, Fulton JS, Park J, et al. Systematic review of basic oral care for the management of oral mucositis in cancer patients. *Support Care Cancer* 2013;21:3165-3177. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24018908>.
303. Kwong KK. Prevention and treatment of oropharyngeal mucositis following cancer therapy: are there new approaches? *Cancer Nurs* 2004;27:183-205. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15238805>.
304. Shih A, Miaskowski C, Dodd MJ, et al. A research review of the current treatments for radiation-induced oral mucositis in patients with head and neck cancer. *Oncol Nurs Forum* 2002;29:1063-1080. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12183755>.
305. Chitapanarux I, Tungkasamit T, Petsuksiri J, et al. Randomized control trial of benzydamine HCl versus sodium bicarbonate for prophylaxis of concurrent chemoradiation-induced oral mucositis. *Support Care Cancer* 2018;26:879-886. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28942587>.
306. Epstein JB, Silverman S, Jr., Paggiarino DA, et al. Benzydamine HCl for prophylaxis of radiation-induced oral mucositis: results from a multicenter, randomized, double-blind, placebo-controlled clinical trial. *Cancer* 2001;92:875-885. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11550161>.
307. Pirmohamed M. Pharmacogenetics and pharmacogenomics. *Br J Clin Pharmacol* 2001;52:345-347. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11678777>.
308. Clinical Pharmacogenetics Implementation Consortium (CPIC). Guidelines. 2021. Available at: <https://cpicpgx.org/guidelines/>. Accessed March 30, 2021.
309. Relling MV, Klein TE, Gammal RS, et al. The clinical pharmacogenetics implementation consortium: 10 years later. *Clin Pharmacol Ther* 2020;107:171-175. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31562822>.
310. Crews KR, Monte AA, Huddart R, et al. Clinical pharmacogenetics implementation consortium guideline for CYP2D6, OPRM1, and COMT genotypes and select opioid therapy. *Clin Pharmacol Ther* 2021. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33387367>.
311. Hicks JK, Swen JJ, Thorn CF, et al. Clinical pharmacogenetics implementation consortium guideline for CYP2D6 and CYP2C19 genotypes and dosing of tricyclic antidepressants. *Clin Pharmacol Ther* 2013;93:402-408. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23486447>.
312. Hicks JK, Sangkuhl K, Swen JJ, et al. Clinical pharmacogenetics implementation consortium guideline (CPIC) for CYP2D6 and CYP2C19 genotypes and dosing of tricyclic antidepressants: 2016 update. *Clin Pharmacol Ther* 2017;102:37-44. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27997040>.
313. Theken KN, Lee CR, Gong L, et al. Clinical Pharmacogenetics Implementation Consortium Guideline (CPIC) for CYP2C9 and Nonsteroidal Anti-Inflammatory Drugs. *Clin Pharmacol Ther* 2020;108:191-200. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32189324>.
314. Manikandan P, Nagini S. Cytochrome P450 structure, function and clinical significance: A review. *Curr Drug Targets* 2018;19:38-54. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28124606>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

315. U.S. Food & Drug Administration. Table of Pharmacogenetic Associations. 2020. Available at: <https://www.fda.gov/medical-devices/precision-medicine/table-pharmacogenetic-associations>. Accessed March 31, 2021.

316. Smith DM, Weitzel KW, Cavallari LH, et al. Clinical application of pharmacogenetics in pain management. *Per Med* 2018;15:117-126. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29714124>.

317. Deng GE, Rausch SM, Jones LW, et al. Complementary therapies and integrative medicine in lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e420S-e436S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649450>.

318. Greenlee H, DuPont-Reyes MJ, Balneaves LG, et al. Clinical practice guidelines on the evidence-based use of integrative therapies during and after breast cancer treatment. *CA Cancer J Clin* 2017;67:194-232. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28436999>.

319. Lyman GH, Greenlee H, Bohlke K, et al. Integrative Therapies During and After Breast Cancer Treatment: ASCO Endorsement of the SIO Clinical Practice Guideline. *J Clin Oncol* 2018;36:2647-2655. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29889605>.

320. Chiu HY, Hsieh YJ, Tsai PS. Systematic review and meta-analysis of acupuncture to reduce cancer-related pain. *Eur J Cancer Care (Engl)* 2017;26. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26853524>.

321. Hershman DL, Unger JM, Greenlee H, et al. Effect of acupuncture vs sham acupuncture or waitlist control on joint pain related to aromatase inhibitors among women with early-stage breast cancer: A randomized clinical trial. *Jama* 2018;320:167-176. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29998338>.

322. Pfister DG, Cassileth BR, Deng GE, et al. Acupuncture for pain and dysfunction after neck dissection: results of a randomized controlled trial. *J Clin Oncol* 2010;28:2565-2570. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20406930>.

323. Zick SM, Sen A, Hassett AL, et al. Impact of self-acupressure on co-occurring symptoms in cancer survivors. *JNCI Cancer Spectr* 2018;2:pk064. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30687806>.

324. He Y, Guo X, May BH, et al. Clinical evidence for association of acupuncture and acupressure with improved cancer pain: A systematic review and meta-analysis. *JAMA Oncol* 2020;6:271-278. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31855257>.

325. Raphael J, Hester J, Ahmedzai S, et al. Cancer pain: part 2: physical, interventional and complimentary therapies; management in the community; acute, treatment-related and complex cancer pain: a perspective from the British Pain Society endorsed by the UK Association of Palliative Medicine and the Royal College of General Practitioners. *Pain Med* 2010;11:872-896. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20456069>.

326. Stoelb BL, Molton IR, Jensen MP, Patterson DR. The efficacy of hypnotic analgesia in adults: a review of the literature. *Contemp Hypn* 2009;26:24-39. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20161034>.

327. Huang ST, Good M, Zauszniewski JA. The effectiveness of music in relieving pain in cancer patients: a randomized controlled trial. *Int J Nurs Stud* 2010;47:1354-1362. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20403600>.

328. Kwekkeboom KL, Cherwin CH, Lee JW, Wanta B. Mind-body treatments for the pain-fatigue-sleep disturbance symptom cluster in persons with cancer. *J Pain Symptom Manage* 2010;39:126-138. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19900778>.

329. Cassileth BR, Keefe FJ. Integrative and behavioral approaches to the treatment of cancer-related neuropathic pain. *Oncologist* 2010;15 Suppl 2:19-23. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20489193>.

330. Montgomery GH, Weltz CR, Seltz M, Bovbjerg DH. Brief presurgery hypnosis reduces distress and pain in excisional breast biopsy patients. *Int*



NCCN Guidelines Version 2.2022

Adult Cancer Pain

J Clin Exp Hypn 2002;50:17-32. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11778705>.

331. De Paolis G, Naccarato A, Cibelli F, et al. The effectiveness of progressive muscle relaxation and interactive guided imagery as a pain-reducing intervention in advanced cancer patients: A multicentre randomised controlled non-pharmacological trial. Complement Ther Clin Pract 2019;34:280-287. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/30712739>.

332. Bennett MI, Bagnall AM, Jose Closs S. How effective are patient-based educational interventions in the management of cancer pain? Systematic review and meta-analysis. Pain 2009;143:192-199. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19285376>.

333. Keefe FJ, Abernethy AP, L CC. Psychological approaches to understanding and treating disease-related pain. Annu Rev Psychol 2005;56:601-630. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15709948>.

334. Lovell MR, Lockett T, Boyle FM, et al. Patient education, coaching, and self-management for cancer pain. J Clin Oncol 2014;32:1712-1720. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24799486>.

335. Sheinfeld Gorin S, Krebs P, Badr H, et al. Meta-analysis of psychosocial interventions to reduce pain in patients with cancer. J Clin Oncol 2012;30:539-547. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22253460>.

336. Puchalski C, Ferrell B, Virani R, et al. Improving the quality of spiritual care as a dimension of palliative care: the report of the Consensus Conference. J Palliat Med 2009;12:885-904. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19807235>.

337. Brogan S, Junkins S. Interventional therapies for the management of cancer pain. J Support Oncol 2010;8:52-59. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20464881>.

338. Eidelman A, White T, Swarm RA. Interventional therapies for cancer pain management: important adjuvants to systemic analgesics. J Natl Compr Canc Netw 2007;5:753-760. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17927931>.

339. Tay W, Ho KY. The role of interventional therapies in cancer pain management. Ann Acad Med Singapore 2009;38:989-997. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19956822>.

340. Wong GY, Schroeder DR, Carns PE, et al. Effect of neurolytic celiac plexus block on pain relief, quality of life, and survival in patients with unresectable pancreatic cancer: a randomized controlled trial. JAMA 2004;291:1092-1099. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/14996778>.

341. Chevillat AL, Basford JR. Role of rehabilitation medicine and physical agents in the treatment of cancer-associated pain. J Clin Oncol 2014;32:1691-1702. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24799472>.

342. Arcidiacono PG, Calori G, Carrara S, et al. Celiac plexus block for pancreatic cancer pain in adults. Cochrane Database Syst Rev 2011;CD007519. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21412903>.

343. Zhang CL, Zhang TJ, Guo YN, et al. Effect of neurolytic celiac plexus block guided by computerized tomography on pancreatic cancer pain. Dig Dis Sci 2008;53:856-860. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17676392>.

344. Zheng S, He L, Yang X, et al. Evaluation of intrathecal drug delivery system for intractable pain in advanced malignancies: A prospective cohort study. Medicine (Baltimore) 2017;96:e6354. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28296770>.

345. Stearns LM, Abd-Elseyed A, Perruchoud C, et al. Intrathecal drug delivery systems for cancer pain: An analysis of a prospective, multicenter product surveillance registry. Anesth Analg 2020;130:289-297. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31567325>.

**NCCN Guidelines Version 2.2022**
Adult Cancer Pain

346. Brogan SE, Sindt JE, Jackman CM, et al. Prospective association of serum opioid levels and clinical outcomes in patients with cancer pain treated with intrathecal opioid therapy. *Anesth Analg* 2020;130:1035-1044. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31206429>.
347. Smith TJ, Staats PS, Deer T, et al. Randomized clinical trial of an implantable drug delivery system compared with comprehensive medical management for refractory cancer pain: impact on pain, drug-related toxicity, and survival. *J Clin Oncol* 2002;20:4040-4049. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12351602>.
348. Deer TR, Prager J, Levy R, et al. Polyanalgesic Consensus Conference 2012: recommendations for the management of pain by intrathecal (intraspinal) drug delivery: report of an interdisciplinary expert panel. *Neuromodulation* 2012;15:436-464; discussion 464-436. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22748024>.
349. Gulati A, Puttanniah V, Hung J, Malhotra V. Considerations for evaluating the use of intrathecal drug delivery in the oncologic patient. *Curr Pain Headache Rep* 2014;18:391. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24407749>.
350. Lauretti GR, Rizzo CC, Mattos AL, Rodrigues SW. Epidural methadone results in dose-dependent analgesia in cancer pain, further enhanced by epidural dexamethasone. *Br J Cancer* 2013;108:259-264. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23322191>.
351. Rastogi R, Patel T, Swarm RA. Vertebral augmentation for compression fractures caused by malignant disease. *J Natl Compr Canc Netw* 2010;8:1095-1102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20876546>.
352. Tancioni F, Lorenzetti MA, Navarria P, et al. Percutaneous vertebral augmentation in metastatic disease: state of the art. *J Support Oncol* 2011;9:4-10. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21465731>.
353. Gofeld M, Bhatia A, Burton AW. Vertebroplasty in the management of painful bony metastases. *Curr Pain Headache Rep* 2009;13:288-294. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19586592>.
354. Berenson J, Pflugmacher R, Jarzem P, et al. Balloon kyphoplasty versus non-surgical fracture management for treatment of painful vertebral body compression fractures in patients with cancer: a multicentre, randomised controlled trial. *Lancet Oncol* 2011;12:225-235. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21333599>.
355. Eleraky M, Papanastassiou I, Setzer M, et al. Balloon kyphoplasty in the treatment of metastatic tumors of the upper thoracic spine. *J Neurosurg Spine* 2011;14:372-376. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21250808>.
356. Zou J, Mei X, Gan M, Yang H. Kyphoplasty for spinal fractures from multiple myeloma. *J Surg Oncol* 2010;102:43-47. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20578077>.
357. Wallace AN, McWilliams SR, Connolly SE, et al. Percutaneous image-guided cryoablation of musculoskeletal metastases: Pain palliation and local tumor control. *J Vasc Interv Radiol* 2016;27:1788-1796. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27745968>.
358. Tomasian A, Gangi A, Wallace AN, Jennings JW. Percutaneous thermal ablation of spinal metastases: Recent advances and review. *AJR Am J Roentgenol* 2018;210:142-152. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29112473>.
359. Scipione R, Anzidei M, Bazzocchi A, et al. HIFU for bone metastases and other musculoskeletal applications. *Semin Intervent Radiol* 2018;35:261-267. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30402009>.
360. Kurup AN, Morris JM, Callstrom MR. Ablation of Musculoskeletal Metastases. *AJR Am J Roentgenol* 2017;209:713-721. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28777648>.
361. Hurwitz MD, Ghanouni P, Kanaev SV, et al. Magnetic resonance-guided focused ultrasound for patients with painful bone metastases: phase III trial results. *J Natl Cancer Inst* 2014;106. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24760791>.



NCCN Guidelines Version 2.2022

Adult Cancer Pain

362. Huisman M, ter Haar G, Napoli A, et al. International consensus on use of focused ultrasound for painful bone metastases: Current status and future directions. *Int J Hyperthermia* 2015;31:251-259. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25677840>.
363. Deib G, Deldar B, Hui F, et al. Percutaneous microwave ablation and cementoplasty: Clinical utility in the treatment of painful extraspinal osseous metastatic disease and myeloma. *AJR Am J Roentgenol* 2019;1-8. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30917019>.
364. Callstrom MR, Dupuy DE, Solomon SB, et al. Percutaneous image-guided cryoablation of painful metastases involving bone: multicenter trial. *Cancer* 2013;119:1033-1041. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23065947>.
365. Bagla S, Sayed D, Smirniotopoulos J, et al. Multicenter prospective clinical series evaluating radiofrequency ablation in the treatment of painful spine metastases. *Cardiovasc Intervent Radiol* 2016;39:1289-1297. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27343124>.
366. Anchala PR, Irving WD, Hillen TJ, et al. Treatment of metastatic spinal lesions with a navigational bipolar radiofrequency ablation device: a multicenter retrospective study. *Pain Physician* 2014;17:317-327. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25054391>.
367. Greenwood TJ, Wallace A, Friedman MV, et al. Combined ablation and radiation therapy of spinal metastases: A novel multimodality treatment approach. *Pain Physician* 2015;18:573-581. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26606009>.
368. Di Staso M, Zugaro L, Gravina GL, et al. A feasibility study of percutaneous Radiofrequency Ablation followed by Radiotherapy in the management of painful osteolytic bone metastases. *Eur Radiol* 2011;21:2004-2010. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21533865>.
369. Sorensen ST, Kirkegaard AO, Carreon L, et al. Vertebroplasty or kyphoplasty as palliative treatment for cancer-related vertebral compression fractures: a systematic review. *Spine J* 2019;19:1067-1075. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30822527>.
370. Flagg A, 2nd, McGreevy K, Williams K. Spinal cord stimulation in the treatment of cancer-related pain: "back to the origins". *Curr Pain Headache Rep* 2012;16:343-349. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22610506>.
371. U.S. Food & Drug Administration. Drug Development and Drug Interactions: Table of Substrates, Inhibitors and Inducers. 2020. Available at: <https://www.fda.gov/drugs/drug-interactions-labeling/drug-development-and-drug-interactions-table-substrates-inhibitors-and-inducers>. Accessed May 10, 2021.
372. U.S. Food & Drug Administration. Prescribing information. Tramadol hydrochloride extended-release tablets for oral use. 2017. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/021692s015bl.pdf. Accessed May 11, 2021.
373. U.S. Food & Drug Administration. Prescribing information. Methadone hydrochloride tablets, for oral use. 2018. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/006134s045bl.pdf. Accessed May 11, 2021.
374. U.S. Food & Drug Administration. Prescribing information. Hydromorphone hydrochloride for intravenous, intramuscular, or subcutaneous use. 2019. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/019034s035bl.pdf. Accessed May 11, 2021.
375. U.S. Food & Drug Administration. Prescribing Information. Codeine sulfate tablets, for oral use. 2021. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/022402s014bl.pdf. Accessed May 11, 2021.
376. U.S. Food & Drug Administration. Prescribing information. Hydrocodone bitartrate extended-release capsules, for oral use. 2021. Available at:



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2022 Adult Cancer Pain

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/202880s0201bl.pdf. Accessed May 11, 2021.

377. U.S. Food & Drug Administration. Prescribing information. Oxycodone hydrochloride extended-release tablets, for oral use. 2021.

Available at:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/022272s0461bl.pdf. Accessed May 11, 2021.

378. U.S. Food & Drug Administration. Prescribing information. Fentanyl transdermal system. 2021. Available at:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/019813s0811bl.pdf. Accessed May 11, 2021.

379. U.S. Food & Drug Administration. Prescribing information. Oxymorphone hydrochloride extended-release tablets, for oral use. 2021.

Available at:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/021610s0281bl.pdf. Accessed May 11, 2021.

380. U.S. Food & Drug Administration. Prescribing information.

Tapentadol tablets for oral use. 2021. Available at:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/022304s0231bl.pdf. Accessed May 11, 2021.

381. Coffman BL, King CD, Rios GR, Tephly TR. The glucuronidation of opioids, other xenobiotics, and androgens by human UGT2B7Y(268) and UGT2B7H(268). *Drug Metab Dispos* 1998;26:73-77. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/9443856>.