

EAU Guidelines on Urological Trauma

N.D. Kitrey (Chair), N. Djakovic, P. Hallscheidt, F.E. Kuehhas,
N. Lumen, E. Serafetinidis, D.M. Sharma
Guidelines Associates: Y. Abu-Ghanem, A. Sujenthiran,
M. Waterloos

TABLE OF CONTENTS

PAGE

1.	INTRODUCTION	5
	1.1 Aim and objectives	5
	1.2 Panel composition	5
	1.3 Available publications	5
	1.4 Publication history	5
2.	METHODS	5
	2.1 Evidence sources	5
	2.2 Peer review	6
3.	EPIDEMIOLOGY, CLASSIFICATION & GENERAL MANAGEMENT PRINCIPALS	6
	3.1 Definition and Epidemiology	6
	3.2 Classification of trauma	6
	3.3 General management principals	6
	3.3.1 The Initial evaluation	6
	3.3.2 Polytrauma managed in major trauma centres leads to improved survival	7
	3.3.3 Damage control	7
	3.3.4 Mass casualty events and Triage	7
	3.3.5 The role of thromboprophylaxis and bed rest	7
	3.3.6 Antibiotic stewardship	7
	3.3.7 Urinary catheterisation	7
4.	UROGENITAL TRAUMA GUIDELINES	8
	4.1 Renal Trauma	8
	4.1.1 Epidemiology, aetiology and pathophysiology	8
	4.1.2 Evaluation	8
	4.1.3 Imaging: criteria for radiographic assessment	9
	4.1.3.1 Computed tomography	9
	4.1.3.2 Ultrasonography (US)	9
	4.1.3.3 Intravenous pyelography (IVP)	9
	4.1.3.4 Magnetic resonance imaging (MRI)	9
	4.1.3.5 Radionuclide scans	9
	4.1.4 Disease management	9
	4.1.4.1 Non-operative management	9
	4.1.4.1.1 Blunt renal injuries	9
	4.1.4.1.2 Penetrating renal injuries	10
	4.1.4.1.3 Selective angioembolisation	10
	4.1.4.1.4 Urinary catheterisation	10
	4.1.4.1.5 Repeat imaging (early)	10
	4.1.4.2 Surgical management	10
	4.1.4.2.1 Indications for renal exploration	10
	4.1.4.2.2 Operative findings and reconstruction	11
	4.1.5 Follow-up	11
	4.1.5.1 Complications	11
	4.1.6 Iatrogenic renal injuries	11
	4.1.7 Summary of evidence and recommendations for evaluation and management of renal trauma	12
	4.1.8 Treatment algorithms	13
	4.2 Ureteral Trauma	14
	4.2.1 Incidence	14
	4.2.2 Epidemiology, aetiology, and pathophysiology	14
	4.2.3 Diagnosis	14
	4.2.3.1 Clinical diagnosis	14
	4.2.3.2 Radiological diagnosis	15
	4.2.4 Prevention of iatrogenic trauma	15
	4.2.5 Management	15
	4.2.5.1 Proximal and mid-ureteral injury	15
	4.2.5.2 Distal ureteral injury	15

	4.2.5.3	Long segment ureteral injury	16
	4.2.6	Summary of evidence and recommendations for the management of ureteral trauma	16
	4.2.7	Treatment algorithms	17
4.3		Bladder Trauma	17
	4.3.1	Classification	17
	4.3.2	Epidemiology, aetiology and pathophysiology	17
	4.3.2.1	Iatrogenic bladder trauma (IBT)	18
	4.3.3	Diagnostic evaluation	18
	4.3.3.1	Cystography	19
	4.3.3.2	Cystoscopy	19
	4.3.3.3	Ultrasound	19
	4.3.4	Prevention	19
	4.3.5	Disease management	19
	4.3.5.1	Conservative management	19
	4.3.5.2	Surgical management	19
	4.3.5.2.1	Blunt non-iatrogenic trauma	20
	4.3.5.2.2	Penetrating non-iatrogenic trauma	20
	4.3.5.2.3	Iatrogenic bladder trauma	20
	4.3.6	Follow-up	20
	4.3.7	Summary of evidence and recommendations for bladder injury	20
4.4		Urethral Trauma	21
	4.4.1	Epidemiology, aetiology and pathophysiology	21
	4.4.1.1	Anterior male urethral injury	21
	4.4.1.2	Posterior male urethral injuries	21
	4.4.1.3	Female urethral injuries	22
	4.4.2	Evaluation	22
	4.4.2.1	Clinical signs	22
	4.4.2.2	Urethrography	22
	4.4.2.3	Cysto-urethroscopy	22
	4.4.2.4	Ultrasound and magnetic resonance imaging	23
	4.4.3	Disease Management	23
	4.4.3.1	Male anterior urethral injuries	23
	4.4.3.1.1	Immediate exploration and urethral reconstruction	23
	4.4.3.1.2	Urinary diversion	23
	4.4.3.2	Male posterior urethral injuries	23
	4.4.3.2.1	Emergency room management	23
	4.4.3.2.2	Early urethral management (less than six weeks after injury)	23
	4.4.3.2.2.1	Immediate urethroplasty	23
	4.4.3.2.2.2	Early urethroplasty	24
	4.4.3.2.2.3	Early re-alignment	24
	4.4.3.2.3	Deferred management (greater than three months after injury)	24
	4.4.3.3	Female urethral injuries	25
	4.4.4	Summary of evidence and recommendations for the evaluation and management of urethral trauma	26
	4.4.5	Treatment algorithms	27
4.5		Genital Trauma	28
	4.5.1	Epidemiology, aetiology and pathophysiology	28
	4.5.2	Diagnostic evaluation	29
	4.5.2.1	Patient history and physical examination	29
	4.5.3	Imaging	29
	4.5.4	Disease management	30
	4.5.4.1	Animal bites	30
	4.5.4.2	Human bites	30
	4.5.4.3	Blunt penile trauma	30
	4.5.4.4	Penile fracture	30
	4.5.4.5	Penetrating penile trauma	30
	4.5.4.6	Penile avulsion injuries and amputation	31

4.5.4.7	Testicular dislocation	31
4.5.4.8	Haematocoele	31
4.5.4.9	Testicular rupture	31
4.5.4.10	Penetrating scrotal trauma	31
4.5.5	Complications	32
4.5.6	Follow up	32
4.5.7	Summary of evidence and recommendations for evaluation and management of genital trauma.	32
5.	REFERENCES	33
6.	CONFLICT OF INTEREST	48
7.	CITATION INFORMATION	48

1. INTRODUCTION

1.1 Aim and objectives

The European Association of Urology (EAU) Guidelines Panel for Urological Trauma have prepared these guidelines in order to assist medical professionals in the management of urological trauma in adults. Paediatric trauma is addressed in the EAU Paediatric Urology Guidelines [1].

It must be emphasised that clinical guidelines present the best evidence available to the experts but following guideline recommendations will not necessarily result in the best outcome. Guidelines can never replace clinical expertise when making treatment decisions for individual patients, but rather help to focus decisions – also taking personal values and preferences/individual circumstances of patients into account. Guidelines are not mandates and do not purport to be a legal standard of care.

1.2 Panel composition

The EAU Urological Trauma Guidelines Panel consists of an international group of urologists and an interventional radiologist, all with particular expertise in urological trauma. All experts involved in the production of this document have submitted potential conflict of interest statements, which can be viewed on the EAU Website Uroweb: <http://uroweb.org/guideline/urological-trauma/?type=panel>.

1.3 Available publications

A quick reference document, the Pocket Guidelines, is available in print and as an app for iOS and Android devices. These are abridged versions which may require consultation together with the full text version. A number of translated versions, alongside several scientific publications in European Urology, the Associations scientific journal, are also available [2-5]. All documents can be viewed through the EAU website: <http://uroweb.org/guideline/urological-trauma/>.

1.4 Publication history

The Urological Trauma Guidelines were first published in 2003. Standard procedure for EAU Guidelines includes an annual assessment of newly published literature in the field to guide future updates. All sections of the 2019 Urological Trauma Guidelines have been fully updated.

2. METHODS

2.1 Evidence sources

For the 2019 Urological Trauma Guidelines, new and relevant evidence has been identified, collated and appraised through a structured assessment of the literature. A broad and comprehensive literature search, covering all sections of the Urological Trauma Guidelines was performed. Databases searched included Medline, EMBASE, and the Cochrane Libraries, covering a time frame between May 31st 2017 and April 1st 2018. A total of 2,104 unique records were identified, retrieved and screened for relevance. A detailed search strategy and summary PRISMA diagram is available online: <http://uroweb.org/guideline/urological-trauma/?type=appendices-publications>. The majority of identified publications were comprised of case reports and retrospective case series. The lack of high-powered randomised controlled trials (RCTs) makes it difficult to draw meaningful conclusions. The panel recognises this critical limitation.

For each recommendation within the guidelines there is an accompanying online strength rating form the bases of which is a modified GRADE methodology [6, 7]. Each strength rating form addresses a number of key elements namely:

1. the overall quality of the evidence which exists for the recommendation, references used in this text are graded according to a classification system modified from the Oxford Centre for Evidence-Based Medicine Levels of Evidence [8];
2. the magnitude of the effect (individual or combined effects);
3. the certainty of the results (precision, consistency, heterogeneity and other statistical or study related factors);
4. the balance between desirable and undesirable outcomes;
5. the impact of patient values and preferences on the intervention;
6. the certainty of those patient values and preferences.

These key elements are the basis which panels use to define the strength rating of each recommendation. The strength of each recommendation is represented by the words 'strong' or 'weak' [9]. The strength of each recommendation is determined by the balance between desirable and undesirable consequences of alternative management strategies, the quality of the evidence (including certainty of estimates), and nature and variability of patient values and preferences. The strength rating forms will be posted online for consultation.

Additional information can be found in the general Methodology section of this print, and online at the EAU website; <http://www.uroweb.org/guideline/>. A list of associations endorsing the EAU Guidelines can also be viewed online at the above address.

2.2 Peer review

The Urological trauma Guidelines has been peer reviewed prior to publication in 2019.

3. EPIDEMIOLOGY, CLASSIFICATION & GENERAL MANAGEMENT PRINCIPALS

3.1 Definition and Epidemiology

Trauma is defined as a physical injury or a wound to living tissue caused by an extrinsic agent. Trauma is the sixth leading cause of death worldwide, accounting for 10% of all mortalities. It accounts for approximately five million deaths each year and causes disability to millions more [10, 11].

About half of all deaths due to trauma are in people aged 15-45 years; trauma is the leading cause of death in this age group [12]. Death from injury is twice as common in males, especially in relation to motor vehicle accidents (MVA's) and interpersonal violence. Trauma is therefore a serious public health problem with significant social and economic costs. Significant variation exists in the causes and the effects of traumatic injuries between geographical areas, and between low, middle, and high-income countries. It should be noted that alcohol and drug abuse increase the rate of traumatic injuries by precipitating interpersonal violence, child and sexual abuse, and MVA's .

3.2 Classification of trauma

Traumatic injuries are classified by the World Health Organization (WHO) into intentional (either interpersonal violence related, war-related or self-inflicted injuries), and unintentional injuries (mainly MVA's, falls, and other domestic accidents). Intentional trauma accounts for approximately half of the trauma-related deaths worldwide [11]. A specific type of unintentional injury is iatrogenic injury which occurs during therapeutic or diagnostic procedures by healthcare personnel. Traumatic insults are classified according to the basic mechanism of the injury into penetrating, when an object pierces the skin, and blunt injuries. Penetrating trauma is further classified according to the velocity of the projectile into:

1. high-velocity projectiles (e.g. rifle bullets - 800-1,000 m/sec);
2. medium-velocity projectiles (e.g. handgun bullets - 200-300 m/sec);
3. low-velocity items (e.g. knife stab).

High-velocity weapons inflict greater damage due to a temporary expansive cavitation that causes destruction in a much larger area than the projectile tract itself. In lower velocity injuries, the damage is usually confined to the projectile tract. Blast injury is a complex cause of trauma which includes blunt and penetrating trauma and burns.

The most commonly used classification grading system is the AAST (American Association for the Surgery of Trauma) injury scoring scale [13]. It is useful for managing renal trauma, but for the other urological organs, the injuries are commonly described by their anatomical site and severity (partial/complete).

3.3 General management principals

3.3.1 The Initial evaluation

The initial emergency assessment of a trauma patient is beyond the focus of these guidelines. It is usually carried out by emergency medicine and trauma specialised personnel following ATLS principles. Detailed further assessment involves cross sectional imaging, laboratory analysis and specialist surgical input. The management of individual organ injury will follow in the sections below.

3.3.2 Polytrauma managed in major trauma centres leads to improved survival

Urological trauma is often associated with significant injuries in the polytraumatised patient [14]. Lessons from civilian trauma networks, military conflict, and mass casualty events have led to many advances in trauma care [15, 16]. These include the widespread acceptance of damage control principles and trauma centralisation to major trauma centres staffed by dedicated trauma teams. The re-organisation of care to these centres has been shown to reduce mortality by 25% and length of stay by four days [15]. Urologists increasingly understand their role in the context of polytrauma with the ultimate aims of improving survivability and decreasing morbidity in these patients.

3.3.3 Damage control

Damage control is a life-saving strategy for severely injured patients that recognises the consequences of the lethal triad of trauma - hypothermia, coagulopathy and acidosis [17-19]. The first of a three phased approach consists of rapid control of haemorrhage and wound contamination. The second phase involves resuscitation in the intensive care unit (ICU), with the goal of restoring normal temperature, coagulation, and tissue oxygenation. The final stage involves definitive surgery when more time-consuming reconstructive procedures are performed in the stabilised patient [20]. Urological intervention needs to be mindful of the phase of management. Temporary abbreviated measures followed by later definitive surgery are required. Complex reconstructive procedures, including organ preservation, are not undertaken. The decision to enter damage control mode is taken by the lead trauma clinician following team discussion.

Urological examples include haemodynamically unstable patients due to suspected renal haemorrhage or pelvic fracture with associated urethral or bladder injury. The options of abdominal packing and temporary urinary drainage by ureteric, bladder or urethral catheterisation are valuable adjuncts to care.

3.3.4 Mass casualty events and Triage

A mass casualty event is one in which the number of injured people and the severity of their injuries exceed the capability of the faculty and staff [21]. Triage, communication and preparedness are important components for a successful response.

Triage after mass casualty events involves difficult moral and ethical considerations. Disaster triage requires differentiation of the few critically injured individuals who can be saved by immediate intervention from the many others with non-life-threatening injuries for whom treatment can be delayed and from those whose injuries are so severe that survival is unlikely in the circumstances [22, 23].

3.3.5 The role of thromboprophylaxis and bed rest

Trauma patients are at high risk of deep venous thrombosis (DVT). Concerns with regard to secondary haemorrhage result in prolonged bed rest post-injury which effectively compounds this risk. Established prophylaxis measures reduce thrombosis and are recommended following systemic review [24]. However, the strength of evidence is not high and as yet there is no evidence to suggest that mortality or pulmonary embolism risk is reduced [25]. Compression stockings and low molecular weight heparins are favoured. The risk of secondary haemorrhage is thought to be low and the practice of strict bed rest has waned in patients who are able to mobilise.

3.3.6 Antibiotic stewardship

Single shot antibiotic doses are common in major trauma. The indication for continuing antibiotics is governed by injury grade, associated injuries and the need for intervention. Patients with urinary extravasation tend to be kept on antibiotics but there is no evidence base for this. Antibiotics should be avoided in lesser trauma e.g. Grade 1-3 renal trauma, and regular review undertaken for those continued on regular dosing.

3.3.7 Urinary catheterisation

Prolonged catheterisation is required in all forms of bladder and urethral injury. Catheterisation is not necessary in stable patients with low-grade renal injury. Patients with heavy haematuria, who require monitoring or ureteric stenting, benefit from catheterisation. This can be removed once haematuria lightens and there is an improvement in the clinical situation. The shortest possible period of catheterisation is advised.

4. UROGENITAL TRAUMA GUIDELINES

4.1 Renal Trauma

4.1.1 *Epidemiology, aetiology and pathophysiology*

Renal trauma is present in up to 5% of all trauma cases [26]. It is most common in young males and has an overall population incidence of 4.9 per 100,000 [27]. Most injuries can be managed non-operatively with successful organ preservation [28-31].

Blunt injuries result from MVAs, falls, sporting injuries, and assault [32]. The kidney and/or hilar structures are directly crushed as a result. Less commonly, sudden deceleration may result in an avulsion injury affecting the vascular structures of the hilum or the ureteropelvic junction (UPJ).

Penetrating injuries are due to stab and gunshot wounds. They tend to be more severe and less predictable than blunt trauma. The prevalence is higher in urban settings [33]. Penetrating injury produces direct tissue disruption of the parenchyma, vascular pedicles, or collecting system. High-velocity bullets or fragments have the potential for greatest parenchymal destruction and are most often associated with multiple-organ injuries [34].

The most commonly used classification system is that of the AAST [13]. It is validated and predicts morbidity and the need for intervention [35, 36]. This remains the most useful of urological trauma classifications; however, the majority of Grade 1 - 4 injuries are now managed conservatively and debate has centred around updating the classification of high-grade injury i.e. identifying the injuries most likely to benefit from early angiographic embolisation, repair or nephrectomy [29, 37].

Table 4.1.1: AAST renal injury grading scale

Grade*	Description of injury
1	Contusion or non-expanding sub-capsular haematoma
	No laceration
2	Non-expanding peri-renal haematoma
	Cortical laceration < 1 cm deep without extravasation
3	Cortical laceration > 1 cm without urinary extravasation
4	Parenchymal laceration: through corticomedullary junction into collecting system <i>or</i> Vascular: segmental renal artery or vein injury with contained haematoma, or partial vessel laceration, or vessel thrombosis
	Parenchymal: shattered kidney <i>or</i> Vascular: renal pedicle or avulsion

*Advance one grade for bilateral injuries up to grade III.

4.1.2 *Evaluation*

The evaluation of stable patients with renal trauma is now based on a trauma protocol computed tomography (CT) scan, often performed prior to involvement of a urologist [38, 39]. It is important to consider all parameters in the evaluation of the patient and to understand the indications for scanning when these are not absolute. Indicators of injury include a direct blow to the flank or rapid deceleration event (fall, high-speed MVAs). Special consideration should be given to pre-existing renal disease [40] or the injured solitary kidney [41]. Pre-existing abnormality e.g. hydronephrosis makes injury more likely following trauma [42].

Vital signs should be recorded throughout the initial evaluation and give the most reliable indication of the urgency of the situation. Physical examination may reveal flank bruising, stab wounds, or bullet entry or exit wounds and abdominal tenderness.

Urinalysis, haematocrit and baseline creatinine are required. Haematuria (visible or non-visible) is the key finding. However major injury such as disruption of the UPJ, pedicle injuries, segmental arterial thrombosis and stab wounds may not have haematuria [43-45]. Haematuria that is out of proportion to the history of trauma may suggest pre-existing pathology [46]. Urine dipstick quickly evaluates for haematuria, but false-negative results can range from 3-10% [47]. An increased creatinine level usually reflects pre-existing renal pathology.

4.1.3 **Imaging: criteria for radiographic assessment**

The goals of imaging are to grade the renal injury, document pre-existing renal pathology, demonstrate presence of the contralateral kidney and identify injuries to other organs. Haemodynamic status will determine the initial imaging pathway with unstable patients potentially requiring immediate intervention. The majority of patients with moderate to major trauma will have had a CT scan performed soon after presentation. In patients who have not had any imaging the indications for renal imaging are [32, 48-51]:

- visible haematuria;
- non-visible haematuria and one episode of hypotension;
- a history of rapid deceleration injury and/or significant associated injuries;
- penetrating trauma;
- clinical signs suggesting renal trauma e.g. flank pain, abrasions, fractured ribs, abdominal distension and/or a mass and tenderness.

4.1.3.1 *Computed tomography*

Computed tomography is the imaging modality of choice in stable patients. It is quick, widely available, and can accurately identify grade of renal injury [52], establish the presence of the contralateral kidney and demonstrate concurrent injuries to other organs. It is ideally performed as a three-phase study:

1. the arterial phase assesses vascular injury and presence of active extravasation of contrast;
2. the nephrographic phase optimally demonstrates parenchymal contusions and lacerations;
3. the delayed phase imaging (5 minutes) identifies collecting system/ureteric injury [53].

In practice, trauma patients usually undergo standardised whole-body imaging protocols and delayed phase imaging of the renal tract is not routinely performed. If there is suspicion that renal injuries have not been fully evaluated, delayed phase imaging is recommended. The rates of contrast-induced nephropathy seen in trauma patients is low [54].

4.1.3.2 *Ultrasonography (US)*

In the primary survey of a critically injured patient, FAST (Focused Assessment Sonography in Trauma) is used to identify hemoperitoneum as cause of haemorrhage and hypovolemia. However, it is not routinely used for the assessment of solid organ injury as it is insensitive, operator dependant, does not define the injury well, and inferior to CT. It is an option for follow-up [55-57].

4.1.3.3 *Intravenous pyelography (IVP)*

Intravenous pyelography has been superseded by cross-sectional imaging and should only be performed when CT is not available [49]. One-shot intra-operative IVP can be used to confirm the presence of a functioning contralateral kidney in patients too unstable to have had pre-operative imaging [58]. The technique consists of a bolus intravenous injection of 2 mL/kg of radiographic contrast followed by a single plain film taken after ten minutes. The quality of the resulting imaging is generally poor. Palpation of the contralateral (unaffected) kidney is a pragmatic surrogate of function [18].

4.1.3.4 *Magnetic resonance imaging (MRI)*

The diagnostic accuracy of MRI in renal trauma is similar to that of CT [59, 60]. However, the logistical challenges of MRI make this modality impractical in acute trauma.

4.1.3.5 *Radionuclide scans*

Radionuclide scans do not play a role in the immediate evaluation of renal trauma patients. In the longer term, follow-up scans can be used to identify areas of scarring, functional loss or obstruction [61].

4.1.4 **Disease management**

4.1.4.1 *Non-operative management*

The non-operative management of renal trauma can be viewed as a “package of care”; a step-wise approach starting with conservative, followed by minimally invasive and/or surgical exploration if necessary. It should be noted that an algorithm for “package of care” will vary in different centres according to available interventions; however, the importance of escalation in treatment interventions should be emphasised [29].

4.1.4.1.1 *Blunt renal injuries*

Haemodynamic stability is the primary criterion for the management of all renal injuries. Non-operative management has become the treatment of choice for most cases. In stable patients, this means a period of bed rest, serial blood tests, regular observation and re-imaging as indicated. Primary conservative management is associated with a lower rate of nephrectomies, and no increase in immediate or long-term morbidity [62].

Grade 1 - 3 injuries are managed non-operatively [63, 64]. Grade 4 injuries are also mostly treated conservatively, but the requirement for subsequent intervention is higher [65]. Persistent urinary extravasation from an otherwise viable kidney after blunt trauma often responds to stent placement and/or percutaneous drainage [66].

Grade 5 injuries often present with haemodynamic instability and major associated injuries. There is thus a higher rate of exploration and nephrectomy [67, 68]. However several studies now support expectant management in patients with Grade 4 and 5 injuries [29, 30, 69-73]. Similarly, unilateral main arterial injuries or arterial thrombosis are normally managed non-operatively in haemodynamically stable patients with surgical repair reserved for bilateral artery injuries or injuries involving a solitary functional kidney [74]. Pre-hospital prolonged warm ischaemia usually results in irreparable damage and renal loss.

4.1.4.1.2 Penetrating renal injuries

Penetrating abdominal wounds have traditionally been managed surgically. However, selective non-operative management of penetrating abdominal wounds is now accepted following detailed assessment in stable patients [65, 75, 76].

For renal injuries, the site of the wound, haemodynamic stability, and diagnostic imaging are the main determinants for intervention. The majority of low-grade stab wounds posterior to the anterior axillary line can be managed non-operatively in stable patients [77]. Grade 3 or higher lesions due to stab wounds in stable patients can be managed expectantly, but warrant closer observation as the clinical course is more unpredictable and associated with a higher rate of delayed intervention [77, 78]. Overall, non-operative management of penetrating injuries in selected stable patients is associated with a successful outcome in up to 50% of stab wounds and up to 40% of gunshot wounds [30, 79-82].

4.1.4.1.3 Selective angioembolisation

Selective angioembolisation (AE) has a key role in the non-operative management of blunt renal trauma in haemodynamically stable patients [83-85]. Currently there are no validated criteria to identify patients who require AE and its use in renal trauma remains heterogeneous. Accepted CT findings indicating the need for AE are active extravasation of contrast, arteriovenous fistula and pseudo-aneurysm [86]. The presence of both active extravasation of contrast and a large haematoma (> 25 mm depth) predict the need for AE with good accuracy [86, 87].

Angioembolisation has been utilised in the non-operative management of all grades of renal injury; however, it is likely to be most beneficial in the setting of high-grade renal trauma (AAST > 3) [83-85]. Non-operative management of high-grade renal trauma, where AE is included in the management algorithm, can be successful in up to 94.9% of Grade 3, 89% of Grade 4 and 52% of Grade 5 injuries [83, 84]. Increasing grade of renal injury is associated with increased risk of failed AE and need for repeat intervention [88].

Repeat embolisation prevents nephrectomy in 67% of patients. Open surgery after failed embolisation usually results in nephrectomy [88, 89]. Despite concerns regarding parenchymal infarction and the use of iodinated contrast media, AE does not appear to affect the occurrence or course of acute kidney injury following renal trauma [90]. In severe polytrauma or high operative risk, the main artery may be embolised, either as a definitive treatment or as a step to a more controlled nephrectomy.

The evidence supporting AE in penetrating renal trauma is sparse. One study found that AE is three times more likely to fail in penetrating trauma [75]. However, AE has been used successfully to treat acute haemorrhage, arteriovenous fistulae and pseudo-aneurysms resulting from penetrating renal trauma [91].

4.1.4.1.4 Urinary catheterisation

Catheterisation is not necessary in stable patients with low-grade injury. Patients with severe visible haematuria, who require monitoring or stenting benefit from catheterisation. A longer period of catheterisation is required if a stent is placed. Once the haematuria lightens and the patient is mobile, the catheter should be removed.

4.1.4.1.5 Repeat imaging (early)

Computed tomography scans should be performed on patients with fever, unexplained decreased haematocrit or significant flank pain. Repeat imaging is also recommended in high-grade injury and in penetrating trauma two to four days after trauma to minimise the risk of missed complications. Repeat imaging can be safely omitted for patients with Grade 1-3 injuries as long as they remain clinically well [92].

4.1.4.2 Surgical management

4.1.4.2.1 Indications for renal exploration

A non- or transient-response to initial fluid resuscitation is an absolute indication for exploration [75, 76]. There is a trend towards ongoing resuscitation and AE [93]. Exploration is influenced by aetiology and grade of injury, transfusion requirements, the need to explore associated abdominal injuries, and the discovery of

an expanding or pulsatile peri-renal haematoma at laparotomy [94]. Grade 5 vascular injury is an absolute indication for exploration [35].

4.1.4.2.2 Operative findings and reconstruction

The overall exploration rate for blunt trauma is low [95]. The goals of exploration following renal trauma are control of haemorrhage and renal salvage. Most series recommend the transperitoneal approach for surgery [96, 97]. Entering the retroperitoneum and leaving the confined haematoma undisturbed within the perinephric fascia is recommended; temporarily packing the fossa tightly with laparotomy pads can salvage the kidney in instances of intra-operative haemorrhage [98]. Access to the pedicle is obtained either through the posterior parietal peritoneum, which is incised over the aorta, just medial to the inferior mesenteric vein or by bluntly dissecting along the plane of the psoas muscle fascia, adjacent to the great vessels, and directly placing a vascular clamp on the hilum [98].

Stable haematomas detected during exploration for associated injuries should not be opened. Central or expanding haematomas indicate injuries of the renal pedicle, aorta, or vena cava and are potentially life-threatening and warrant further exploration [99].

Feasibility of renal reconstruction should be judged during the operation. The overall rate of patients who undergo a nephrectomy during exploration is approximately 30% [100]. Other intra-abdominal injuries also increase the likelihood of nephrectomy [101]. Mortality is associated with overall severity of the injury and not often a consequence of the renal injury itself [102]. High velocity gunshot injuries make reconstruction difficult and nephrectomy is usually required [103].

Renorrhaphy is the most common reconstructive technique. Partial nephrectomy is required when non-viable tissue is detected. Watertight closure of the collecting system is desirable, although closing the parenchyma over the injured collecting system is acceptable.

The use of haemostatic agents and sealants in reconstruction is helpful [104]. In all cases, drainage of the ipsilateral retroperitoneum is recommended.

The repair of vascular injuries is seldom, if ever, effective [105]. Repair should be attempted in patients with a solitary kidney or bilateral injuries [106]. Nephrectomy for main artery injury has outcomes similar to those of vascular repair and does not worsen post-treatment renal function in the short-term. Bleeding or dissection of the main renal artery may also be managed with a stent.

4.1.5 **Follow-up**

The risk of complications relates to aetiology, injury grade, and mode of management [107, 108]. Follow-up includes physical examination, urinalysis, diagnostic imaging, blood pressure measurement and serum creatinine [67]. Potential complications are primarily identified by imaging; however, follow up imaging is not recommended in low-grade uncomplicated injury. Ultrasound can be used to define the post-injury anatomy avoiding further ionising radiation. Nuclear scans are useful for documenting functional recovery following renal injury and reconstruction [61]. Annual blood pressure monitoring is recommended to exclude renovascular hypertension [109].

4.1.5.1 *Complications*

Early (≤ 1 month) complications include bleeding, infection, perinephric abscess, sepsis, urinary fistula, hypertension, urinary extravasation and urinoma. Delayed complications include bleeding, hydronephrosis, calculus formation, chronic pyelonephritis, hypertension, arteriovenous fistulae (AVF), hydronephrosis and pseudo-aneurysms. Bleeding may be life-threatening with elective angiographic embolisation the preferred treatment [110]. Perinephric abscess formation is initially managed by percutaneous drainage [95].

Hypertension is rare [111, 112]. It may occur acutely as a result of external compression from peri-renal haematoma (Page kidney), chronically due to compressive scar formation, or as a result of renal artery thrombosis, segmental arterial thrombosis, renal artery stenosis (Goldblatt kidney), or AVF. Arteriography may be required. Treatment, including medical management, excision of the ischaemic parenchymal segment, vascular reconstruction, or nephrectomy, is indicated if hypertension persists [109].

Arteriovenous fistulae usually present with delayed onset of significant haematuria, most often after penetrating trauma. Percutaneous embolisation is often effective for symptomatic AVF, but larger fistulae may require surgery [113]. The development of pseudo-aneurysm is a rare complication following blunt trauma.

4.1.6 **Iatrogenic renal injuries**

Iatrogenic renal trauma needs to be recognised and managed promptly to minimise morbidity and mortality. The most common causes of iatrogenic renal injuries are percutaneous access to kidney, stone surgery, cancer surgery (laparoscopic and open) and transplantation [3]. The diagnosis and management follow the same principles as outlined previously.


4.1.7 **Summary of evidence and recommendations for evaluation and management of renal trauma**

Summary of evidence	LE
Vital signs on admission give the most reliable indication of the urgency of the situation.	3
Special consideration should be given to patients with a solitary kidney and pre-existing renal disease.	4
Haematuria is a key finding following renal trauma; although, it may not be present in certain situations.	3
A multiphase CT scan is the best method for the diagnosis and staging of renal injuries in haemodynamically stable patients.	3
Haemodynamic stability is the primary criterion for selecting patients for non-operative management.	3
Selective angioembolisation is effective in patients with active bleeding from renal injury, without other indications for immediate abdominal operation.	3
Renal reconstruction should be attempted if haemorrhage is controlled and there is sufficient viable renal parenchyma.	3
Iatrogenic renal injuries are procedure-dependent (1.8-15%); the most common injuries are vascular.	3
Limited literature exists with regard to long-term consequences of renal trauma. Current follow-up includes physical examination, urinalysis, diagnostic imaging, serum creatinine, as well as annual blood pressure monitoring to diagnose renovascular hypertension.	4

Recommendations	Strength rating
Evaluation	
Assess haemodynamic stability upon admission.	Strong
Record past renal surgery, and known pre-existing renal abnormalities (ureteropelvic junction obstruction, solitary kidney, lithiasis).	Strong
Test for haematuria in a patient with suspected renal injury.	Strong
Perform a multiphase computed tomography scan in trauma patients with: <ul style="list-style-type: none"> • visible haematuria; • non-visible haematuria and one episode of hypotension; • a history of rapid deceleration injury and/or significant associated injuries; • penetrating trauma; • clinical signs suggesting renal trauma e.g. flank pain, abrasions, fractured ribs, abdominal distension and/or a mass and tenderness. 	Strong
Management	
Manage stable patients with blunt renal trauma non-operatively with close monitoring and re-imaging as required.	Strong
Manage isolated Grade 1-4 stab and low-velocity gunshot wounds in stable patients non-operatively.	Strong
Use selective angioembolisation for active renal bleeding if there are no other indications for immediate surgical exploration.	Strong
Proceed with renal exploration in the presence of: <ul style="list-style-type: none"> • persistent haemodynamic instability; • Grade 5 vascular or penetrating injury; • expanding or pulsatile peri-renal haematoma. 	Strong
Attempt renal reconstruction if haemorrhage is controlled and there is sufficient viable renal parenchyma.	Weak
Repeat imaging in high-grade injuries and in cases of fever, worsening flank pain, or falling haematocrit.	Strong
Follow-up approximately three months after major renal injury with: <ul style="list-style-type: none"> • physical examination; • urinalysis; • individualised radiological investigation including nuclear scintigraphy; • blood pressure measurement; • renal function tests. 	Weak
Measure blood pressure annually to diagnose renovascular hypertension.	Strong

4.1.8 **Treatment algorithms**
Management of renal trauma

Figure 4.1.1 Management of renal trauma



* Excluding Grade 5 penetrating injuries.

** Antibiotics should be administered for all penetrating injuries.

--- If haemodynamically unstable.

CT = computed tomography; Ht = haematocrit; SAE = selective angioembolisation.

4.2 Ureteral Trauma

4.2.1 Incidence

Trauma to the ureters is relatively rare as they are protected from injury by their small size, mobility, and the adjacent vertebrae, bony pelvis and muscles. Iatrogenic trauma is the most common cause of ureteral injury (approximately 80%) [114]. It is seen in open, laparoscopic or endoscopic surgery and is often missed intra-operatively. Any trauma to the ureter may result in severe sequelae [115].

4.2.2 Epidemiology, aetiology, and pathophysiology

Overall, ureteral trauma accounts for 1-2.5% of urinary tract trauma [114, 116-118], with even higher rates in modern combat injuries [119]. Penetrating external ureteral trauma, mainly caused by gunshot wounds, dominates most of the modern series, both civilian and military [114, 116, 120]. About one-third of cases of external trauma to the ureters are caused by blunt trauma, mostly road traffic injuries [117, 118].

Ureteral injury should be suspected in all cases of penetrating abdominal injury, especially gunshot wounds, as it occurs in 2-3% of cases [114]. It should also be suspected in blunt trauma with a deceleration mechanism, as the renal pelvis can be torn away from the ureter [114]. The distribution of external ureteral injuries along the ureter varies between series, but it is more common in the upper ureter [116-118].

Iatrogenic ureteral trauma can result from various mechanisms: ligation or kinking with a suture, crushing from a clamp, partial or complete transection, thermal injury, or ischaemia from devascularisation [120-122]. It usually involves the lower ureter [114, 120, 121, 123]. Gynaecological operations are the most common cause of iatrogenic trauma (Table 4.2.1), but it may also occur in colorectal operations, especially abdominoperineal resection and low anterior resection [124]. The incidence of urological iatrogenic trauma has decreased in the last twenty years due to improvements in technique, instruments and surgical experience [120, 125]. New methods such as robotic surgery in gynaecology have not further reduced the rate of ureteral injuries [126].

Risk factors for iatrogenic trauma include conditions that alter the normal anatomy, e.g. advanced malignancy, prior surgery or irradiation, diverticulitis, endometriosis, anatomical abnormalities, and major haemorrhage [120, 124, 127, 128]. Occult ureteral injury occurs more often than reported and not all injuries are diagnosed intra-operatively [115].

Table 4.2.1: Incidence of ureteral injury in various procedures

Procedure	Percentage %
Gynaecological [123, 129, 130]	
Vaginal hysterectomy	0.02 – 0.5
Abdominal hysterectomy	0.03 – 2.0
Laparoscopic hysterectomy	0.2 – 6.0
Urogynaecological (anti-incontinence/prolapse)	1.7 – 3.0
Colorectal [122, 129, 131]	0.15 – 10
Ureteroscopy [125]	
Mucosal abrasion	0.3 – 4.1
Ureteral perforation	0.2 – 2.0
Intussusception/avulsion	0 – 0.3
Radical prostatectomy [132]	
Open retropubic	0.05 – 1.6
Robot-assisted	0.05 – 0.4

4.2.3 Diagnosis

The diagnosis of ureteral trauma is challenging; therefore, a high index of suspicion should be maintained. In penetrating external trauma, it is usually made intra-operatively during laparotomy [133], while it is delayed in most blunt trauma and iatrogenic cases [120, 123, 134].

4.2.3.1 Clinical diagnosis

External ureteral trauma usually accompanies severe abdominal and pelvic injuries. Penetrating trauma is usually associated with vascular and intestinal injuries, while blunt trauma is associated with damage to the pelvic bones and lumbosacral spine injuries [117, 118]. Haematuria is an unreliable and poor indicator of ureteral injury, as it is present in only 50-75% of patients [114, 120, 135].

Iatrogenic injury may be noticed during the primary procedure, when intravenous dye (e.g. indigo carmine) is injected to exclude ureteral injury. However, it is usually noticed later, when it is discovered by subsequent evidence of upper tract obstruction, urinary fistulae formation or sepsis. The following clinical signs are characteristic of delayed diagnosis flank pain, urinary incontinence, vaginal or drain urinary leakage, haematuria, fever, uraemia or urinoma. When the diagnosis is missed, the complication rate increases [114, 119, 134]. Early recognition facilitates immediate repair and provides better outcome [130, 136].

4.2.3.2 *Radiological diagnosis*

Multi-phase CT is the mainstay imaging technique for trauma patients. Generally, it is widely available and allows for multi-phasic assessment of all of the structures in the pelvis and abdomen. Computed tomography urography (CTU) is the examination of choice when ureteral injuries are suspected [137]. Extravasation of contrast medium in the delayed phase is the hallmark sign of ureteral trauma. However, hydronephrosis, ascites, urinoma or mild ureteral dilation are often the only signs. In unclear cases, a retrograde or antegrade urography is the optimum standard for confirmation [120]. Intravenous pyelography, especially one-shot IVP, is unreliable in diagnosis, as it is negative in up to 60% of patients [114, 120].

4.2.4 **Prevention of iatrogenic trauma**

The prevention of iatrogenic trauma to the ureters depends upon the visual identification of the ureters and careful intra-operative dissection in their proximity [120-122]. The use of prophylactic pre-operative ureteral stent insertion assists in visualisation and palpation and is used in complicated cases (about 4% in a large cohort) [138, 139].

It is probably also advantageous in making it easier to detect ureteral injury [121]; however, it does not decrease the rate of injury [140]. Apart from its evident disadvantages (potential complications and cost), a stent may alter the location of the ureter and diminish its flexibility [121, 131].

4.2.5 **Management**

Management of ureteral trauma depends on many factors concerning the nature, severity and location of the injury. Immediate diagnosis of a ligation injury during an operation can be managed by de-ligation and stent placement. Partial injuries can be repaired immediately with a stent or urinary diversion by a nephrostomy tube. Stenting is helpful because it provides canalisation and may decrease the risk of stricture [120]. On the other hand, its insertion has to be weighed against potentially aggravating the severity of the ureteral injury. Immediate repair of complete ureteral injury is usually advisable. The ureter is mobilised on both ends and a spatulated end-to-end anastomosis is performed. However, in cases of unstable trauma patients, a 'damage control' approach is preferred with ligation of the ureter, diversion of the urine (e.g. by a nephrostomy), and a delayed definitive repair [141]. Injuries that are diagnosed late are usually managed first by a nephrostomy tube or a stent [120].

Endo-urological treatment of delayed-diagnosed ureteral injuries by internal stenting, with or without dilatation, is the first step in most cases. It is performed either retrogradely or antegradely through a percutaneous nephrostomy, and it has a variable success rate of 14-19% in published series [142-144]. An open surgical repair is necessary in case of failure. The basic principles for any surgical repair of a ureteral injury are outlined in Table 4.2.2. Wide debridement is highly recommended for gunshot wound injuries due to the 'blast effect' of the injury.

4.2.5.1 *Proximal and mid-ureteral injury*

Injuries shorter than 2-3 cm can usually be managed by a primary uretero-ureterostomy [114]. When this approach is not feasible, a uretero-calycostomy should be considered. In case of a large extra-renal pelvis and a stricture at the UPJ, a pelvic spiral flap according to Culp-DeWeerd is an option [145]. In extensive ureteral loss, a transuretero-ureterostomy is a valid option, where the proximal stump of the ureter is transposed across the midline and anastomosed to the contralateral ureter. The reported stenosis rate is 4% and intervention or revision occur in 10% of cases [146].

4.2.5.2 *Distal ureteral injury*

Distal injuries are best managed by ureteral re-implantation (uretero-neocystostomy) because the primary trauma usually jeopardises the blood supply to the distal ureter. The question of refluxing vs. non-refluxing ureteral re-implantation remains unresolved in the literature. The risk for clinically significant reflux should be weighed against the risk for ureteral obstruction.

A psoas hitch between the bladder and the ipsilateral psoas tendon is usually needed to bridge the gap and to protect the anastomosis from tension. The contralateral superior vesical pedicle may be divided to improve bladder mobility. The reported success rate is very high (97%) [146]. In extensive mid-lower ureteral

injury, the large gap can be bridged with a tabularised L-shaped bladder flap (Boari flap). It is a time-consuming operation and not usually suitable in the acute setting. The success rate is reported to be 81-88% [147].

4.2.5.3 Long segment ureteral injury

A longer ureteral injury can be replaced using a segment of the intestines, usually the ileum (ileal interposition graft). This should be avoided in patients with impaired renal function or known intestinal disease. Follow-up should include serum chemistry to diagnose hyperchloremic metabolic acidosis [148]. The long-term complications include anastomotic stricture (3%) and fistulae (6%) [149]. In cases of extensive ureteral loss or after multiple attempts at ureteral repair, the kidney can be relocated to the pelvis (auto-transplantation). The renal vessels are anastomosed to the iliac vessels and a ureteral re-implantation is performed [150, 151].

Buccal mucosa ureteroplasty is another option for long segment ureteral injury, especially after a previous failed reconstruction, as an alternative to auto-transplantation. The overall success rate is 90% but experience is limited [152].

Table 4.2.2: Principles of surgical repair of ureteral injury

Debridement of necrotic tissue.
Spatulation of ureteral ends.
Watertight mucosa-to-mucosa anastomosis with absorbable sutures.
Internal stenting.
External drain.
Isolation of injury with peritoneum or omentum.


4.2.6 Summary of evidence and recommendations for the management of ureteral trauma

Summary of evidence	LE
Iatrogenic ureteral trauma is the most common cause of ureteral injury.	3
Gunshot wounds account for the majority of penetrating ureteral injuries, while MVAs account for most blunt injuries.	3
Ureteral trauma usually accompanies severe abdominal and pelvic injuries.	3
Haematuria is an unreliable and poor indicator of ureteral injury.	3
Pre-operative prophylactic stents do not prevent ureteral injury; however, they may assist in its detection.	2
Endo-urological treatment of small ureteral fistulae and strictures is safe and effective.	3
Major ureteral injury requires ureteral reconstruction following temporary urinary diversion.	3

Recommendations	Strength rating
Visually identify the ureters to prevent ureteral trauma during abdominal and pelvic surgery.	Strong
Beware of concomitant ureteral injury in all abdominal penetrating trauma, and in deceleration-type blunt trauma.	Strong
Use pre-operative prophylactic stents in high-risk cases.	Strong
Repair iatrogenic ureteral injuries recognised during surgery immediately.	Strong
Treat iatrogenic ureteral injuries with delayed diagnosis by nephrostomy tube/JJ stent urinary diversion.	Strong
Manage ureteral strictures by ureteral reconstruction according to the location and length of the affected segment.	Strong

4.2.7 **Treatment algorithms**
Management of ureteric injuries

Figure 4.2.1: Management of ureteric injuries



4.3 Bladder Trauma

4.3.1 Classification

Bladder trauma is primarily classified according to the location of the injury: **intra**peritoneal, **extra**peritoneal, and **combined** intra-extra-peritoneal [153], as it guides further management [154]. Bladder trauma is categorised by aetiology: **non-iatrogenic** (blunt and penetrating) and **iatrogenic** (external and internal).

4.3.2 Epidemiology, aetiology and pathophysiology

Motor vehicle accidents are the most common cause of blunt bladder injury, followed by falls and other accidents. The main mechanisms are pelvic crush and blows to the lower abdomen [117, 153, 155]. Most patients with blunt bladder injury have associated pelvic fractures (60-90%) and other intra-abdominal injuries (44-68.5%) [156, 157]. Pelvic fractures are associated with bladder injury in 3.2-3.6% of cases [117, 158]. Bladder injury is associated with urethral injury in 5-20% of cases [154, 157, 159].

The incidence of extra-peritoneal (22.4-61.1%), and intra-peritoneal (38.9-65.8%) injuries varies

among series [160]. **Extraperitoneal injury** is almost always associated with pelvic fractures [155, 157]. It is usually caused by distortion of the pelvic ring, with shearing of the anterolateral bladder wall near the bladder base (at its fascial attachments), or by a contrecoup at the opposite side. The highest risk of bladder injury was found in disruptions of the pelvic circle with displacement > 1 cm, diastasis of the pubic symphysis > 1 cm, and pubic rami fractures [117, 154]. An isolated acetabular fracture is not likely to be associated with bladder injury [154, 157]. Occasionally, the bladder is directly perforated by a sharp bony fragment [154].

Intraperitoneal injury is caused by a sudden rise in intravesical pressure of a distended bladder, secondary to a blow to the pelvis or lower abdomen. The bladder dome is the weakest point of the bladder and ruptures will usually occur there [154]. Penetrating injuries, mainly gunshot wounds, are rare except in conflict zones and violent urban areas [153, 161, 162]. Improvised explosive devices are the main cause of combat related bladder injuries in asymmetric warfare [163].

4.3.2.1 Iatrogenic bladder trauma (IBT)

The bladder is the urological organ that is most commonly affected by iatrogenic injury [164]. Table 4.3.2 shows the incidence of IBT during various procedures. **External IBT** occurs most often during obstetric and gynaecological procedures, followed by urological and general surgical operations [164]. Main risk factors are previous surgery, inflammation and malignancy [164]. Bladder perforations occur in up to 4.9% of mid-urethral sling (MUS) operations for stress urinary incontinence in women. This rate is significantly lower in the obturator route compared to the retropubic route [165].

Internal IBT mainly occurs during transurethral resection of the bladder (TURB). Reported risk factors are larger tumours, older age, pre-treated bladders (previous TURB, intravesical instillations) and location at the bladder dome [166, 167]. Tumours at the lateral wall pose a risk factor because of the obturator jerk, [168, 169]. Extraperitoneal perforations are more frequent than intraperitoneal perforations [167, 170], and perforations requiring intervention are rare (0.16-0.57%) [166].

Table 4.3.2: Incidence of iatrogenic bladder trauma during various procedures

Procedure	Percentage (%)
Laparoscopic/Robotic radical hysterectomy (malignant) [171]	4.19-4.59
Abdominal radical hysterectomy (malignant) [171]	2.37
Laparoscopic/Abdominal hysterectomy (benign) [172, 173]	1-2.7
Vaginal hysterectomy (benign) [172, 173]	0.6-2.5
Caesarean delivery [174]	0.08-0.94
Abdominal cytoreductive surgery [175]	4.5
Rectal procedures [176]	0.27-0.41
Small/large bowel procedures [176]	0.12-0.14
Laparoscopic inguinal hernia repair [177]	0.04-0.14
Urology Specific	
Transurethral resection of the bladder [178, 179]	3.5-58
Retropubic male sling [180]	8.0-19
Mid-urethral sling (Retropubic route) [165, 181]	4.91-5.5
Transvaginal mesh surgery [182]	2.84
Pubovaginal sling [181]	2.8
Laparoscopic sacrocolpopexy [183]	1.9
Mid-urethral sling (Transobturator route) [181]	1.61
Burch colposuspension [181, 184]	1.0-1.2
Native tissue colporrhaphy [182]	0.53

4.3.3 Diagnostic evaluation

The principal sign of bladder injury is visible haematuria [154, 155]. Absolute indications for bladder imaging include: visible haematuria and a pelvic fracture [154] or non-visible haematuria combined with high-risk pelvic fracture (disruption of the pelvic circle with displacement > 1 cm or diastasis of the pubic symphysis > 1 cm) or posterior urethral injury [154]. In the absence of these absolute indications, further imaging is based on clinical signs and symptoms including [154, 155, 161, 185]:

- inability to void or inadequate urine output;
- abdominal tenderness or distension due to urinary ascites, or signs of urinary ascites in abdominal imaging;
- uraemia and elevated creatinine level due to intraperitoneal re-absorption;
- entry/exit wounds at lower abdomen, perineum or buttocks in penetrating injuries.

Intra-operative signs of external iatrogenic bladder injury include: extravasation of urine, visible laceration, visible bladder catheter, and blood and/or gas in the urine bag during laparoscopy [174]. Direct inspection is the most reliable method of assessing bladder integrity [164]. Intravesical instillation of dye helps to detect smaller lesions [186]. If bladder perforation is close to the trigone, the ureteric orifices should be inspected [164, 174].

Internal bladder injury is recognised by cystoscopic identification of fatty tissue, dark space, or bowel [178]. It may also be detected by the inability to distend the bladder, low return of irrigation fluid, or abdominal distension [187].

Post-operatively, missed bladder trauma is diagnosed by haematuria, abdominal pain, abdominal distension, ileus, peritonitis, sepsis, urine leakage from the wound, decreased urinary output, or increased serum creatinine [164, 174]. An IBT during hysterectomy or caesarean delivery can result in vesico-vaginal or vesico-uterine fistulae [174, 188].

4.3.3.1 *Cystography*

Cystography is the preferred diagnostic modality for non-iatrogenic bladder injury and for a suspected IBT in the post-operative setting [188, 189]. Both plain and CT cystography have a comparable sensitivity (90-95%) and specificity (100%) [155, 190]. However, CT cystography is superior in the identification of bony fragments in the bladder and bladder neck injuries, as well as concomitant abdominal injuries [154, 157].

Cystography must be performed using retrograde filling of the bladder with a minimum volume of 300-350 mL of dilute contrast material [189, 191]. Passive bladder filling by clamping the urinary catheter during the excretory phase of CT or IVP is not sufficient to exclude bladder injury [155]. Intraperitoneal extravasation is visualised by free contrast medium in the abdomen outlining bowel loops or abdominal viscera [192]. Extraperitoneal bladder injury is typically diagnosed by flame-shaped areas of contrast extravasation in the peri-vesical soft tissues. Contrast medium in the vagina is a sign of vesico-vaginal fistula [188].

4.3.3.2 *Cystoscopy*

Cystoscopy is the preferred method for detection of intra-operative bladder injuries as it may directly visualise the laceration and can localise the lesion in relation to the position of the trigone and ureteral orifices [192]. A lack of bladder distension during cystoscopy suggests a large perforation. Cystoscopy is recommended to detect perforation of the bladder (or urethra) following retropubic sub-urethral sling operations [165, 184]. Routine intra-operative cystoscopy during other gynaecologic procedures is not recommended [193], although the threshold to perform it should be low in any suspected bladder injury.

4.3.3.3 *Ultrasound*

Ultrasound alone is insufficient in the diagnosis of bladder trauma, although it can be used to visualise intraperitoneal fluid or an extraperitoneal collection of fluid.

4.3.4 **Prevention**

The risk of bladder injury is reduced by emptying the bladder by urethral catheterisation in every procedure where the bladder is at risk [186, 194]. Furthermore, the catheter's balloon can aid in identification of the bladder [186]. For tumours at the lateral wall, obturator nerve block or general anaesthesia with adequate muscle relaxation can reduce the incidence of internal IBT during TURB [169]. There is conflicting evidence whether bipolar TURB can reduce the risk for an obturator jerk [168, 169]. The use of combat pelvic protection systems reduces the risk of bladder and other genitourinary injuries due to the blast mechanism of improvised explosive devices [163, 195].

4.3.5 **Disease management**

4.3.5.1 *Conservative management*

Conservative treatment, which comprises of clinical observation, continuous bladder drainage and antibiotic prophylaxis [167], is the standard treatment for an uncomplicated extraperitoneal injury due to blunt [154, 157, 159] or iatrogenic trauma [167].

Conservative treatment can also be chosen for uncomplicated intraperitoneal injury after TURB or other operations, only in the absence of peritonitis and ileus [179, 192]. Placement of an intraperitoneal drain is advocated, especially when the lesion is larger [187, 196]. Penetrating extraperitoneal bladder injuries (only if minor and isolated) can also be managed conservatively [160, 185, 197].

4.3.5.2 *Surgical management*

Bladder closure is performed with absorbable sutures [160, 164]. There is no evidence that two-layer is superior to watertight single-layer closure [157, 160].

4.3.5.2.1 Blunt non-iatrogenic trauma

Most extraperitoneal ruptures can be treated conservatively, however bladder neck involvement, bone fragments in the bladder wall, concomitant rectal or vaginal injury or entrapment of the bladder wall necessitate surgical intervention [154]. There is an increasing trend to treat pelvic ring fractures with open stabilisation and internal fixation with osteosynthetic material. During this procedure, an extraperitoneal rupture should be sutured concomitantly in order to reduce the risk of infection [154, 155]. Likewise, an extraperitoneal rupture should be sutured during surgical exploration for other injuries, in order to decrease the risk of complications and to reduce recovery time [159].

Intraperitoneal ruptures should always be managed by surgical repair [154, 157] because intraperitoneal urine extravasation can lead to peritonitis, intra-abdominal sepsis and death [156]. Abdominal organs should be inspected for possible associated injuries and urinomas must be drained if detected. Laparoscopic suturing of the intraperitoneal rupture is also possible [155].

4.3.5.2.2 Penetrating non-iatrogenic trauma

Penetrating bladder injury is managed by emergency exploration, debridement of devitalised bladder wall and primary bladder repair [161, 162]. A midline exploratory cystotomy is advised to inspect the bladder wall and the distal ureters [160, 161]. In gunshot wounds, there is a strong association with intestinal and rectal injuries, usually requiring faecal diversion [161, 185]. Most gunshot wounds are associated with two transmural injuries (entry and exit wounds) and the bladder should be carefully checked for these two lesions [161]. As the penetrating agent (bullet, knife) is not sterile, antibiotic treatment is advised [162].

4.3.5.2.3 Iatrogenic bladder trauma

Perforations recognised intra-operatively are primarily closed [198]. Bladder injuries not recognised during surgery or internal injuries should be managed according to their location. The standard of care for intraperitoneal injuries is surgical exploration and repair [192]. If surgical exploration is performed after TURB, the bowel must be inspected to rule out concomitant injury [166]. For extraperitoneal injuries, exploration is only needed for perforations complicated by symptomatic extravescical collections. It requires drainage of the collection, with or without closure of the perforation [199]. If bladder perforation is encountered during mid-urethral sling or transvaginal mesh procedures, sling re-insertion and urethral catheterisation (two to seven days) should be performed [200].

4.3.6 **Follow-up**

Continuous bladder drainage is required to prevent elevated intravesical pressure and to allow the bladder to heal [164, 201]. Conservatively treated bladder injuries (traumatic or external IBT) are followed up by cystography to rule out extravasation and ensure proper bladder healing [154]. The first cystography is planned approximately ten days after injury [160]. In case of ongoing leakage, cystoscopy should be performed to rule out bony fragments in the bladder, and a second cystography is warranted one week later [154].

After operative repair of a simple injury in a healthy patient, the catheter can be removed after five to ten days without cystography [160, 201]. In cases of complex injury (trigone involvement, ureteric re-implantation) or risk factors of impaired wound healing (e.g. steroids, malnutrition) cystography is advised [160, 201]. For conservatively treated internal IBT, catheter drainage, lasting five days for extraperitoneal and seven days for intraperitoneal perforations is proposed [167, 170].

4.3.7 **Summary of evidence and recommendations for bladder injury**

Summary of evidence	LE
The combination of pelvic fracture and visible haematuria is highly suggestive of bladder injury.	3
Cystography is the preferred diagnostic modality for non-iatrogenic bladder injury and for suspected IBT in the post-operative setting.	3
Cystography must be performed using retrograde filling of the bladder with a minimum volume of 300-350 mL of dilute contrast material. Passive bladder filling by clamping the urinary catheter during the excretory phase of CT or IVP is not sufficient to exclude bladder injury.	3
The risk of bladder perforation during mid-urethral sling operations for stress urinary incontinence is lower for the obturator route compared to the retropubic route.	1a
Conservative treatment, which comprises of clinical observation, continuous bladder drainage and antibiotic prophylaxis, is the standard treatment for an uncomplicated extraperitoneal injury due to blunt trauma.	3
In extraperitoneal bladder injury with either bladder neck involvement, bone fragments in the bladder wall, concomitant rectal or vaginal injury, or entrapment of the bladder wall, surgical intervention is necessary in order to decrease the risk of complications and to reduce recovery time.	3

Intraperitoneal bladder trauma is managed by surgical repair because intraperitoneal urine extravasation can lead to peritonitis, intra-abdominal sepsis and death.	3
Conservative treatment is suitable for uncomplicated intraperitoneal injury during endourological procedures, in the absence of peritonitis and ileus.	3
In cases of complex injury (trigone involvement, ureteric re-implantation) or risk factors of impaired wound healing (e.g. steroids, malnutrition) cystography is advised after bladder repair.	2a

Recommendations	Strength rating
Perform cystography in the presence of visible haematuria and pelvic fracture.	Strong
Perform cystography in case of suspected iatrogenic bladder injury in the post-operative setting.	Strong
Perform cystography with active retrograde filling of the bladder with dilute contrast (300-350 mL).	Strong
Perform cystoscopy to rule out bladder injury during retropubic sub-urethral sling procedures.	Strong
Manage uncomplicated blunt extraperitoneal bladder injuries conservatively.	Weak
Manage blunt extraperitoneal bladder injuries operatively in cases of bladder neck involvement and/or associated injuries that require surgical intervention.	Strong
Manage blunt intraperitoneal injuries by surgical exploration and repair.	Strong
Manage small uncomplicated intraperitoneal bladder injuries during endoscopic procedures conservatively.	Weak
Perform cystography to assess bladder wall healing after repair of a complex injury or in case of risk factors for wound healing.	Strong

4.4 Urethral Trauma

4.4.1 *Epidemiology, aetiology and pathophysiology*

4.4.1.1 *Anterior male urethral injury*

The bulbar urethra is the most common site affected by **blunt** trauma. In bulbar injuries, the bulb is compressed against the pubic symphysis, resulting in rupture of the urethra at the site of compression [202]. Possible mechanisms are straddle injuries or kicks to the perineum. A penile fracture can be complicated by a urethral injury in approximately 15% of cases [203, 204]. **Penetrating** anterior injuries are rare and are usually caused by gunshot wounds, stab wounds, dog bites, impalement or penile amputations [202, 205]. Depending on the affected segment, penetrating injuries are usually associated with penile, testicular and/or pelvic injuries [205, 206]. Insertion of **foreign bodies** is another rare cause of anterior injury. It is usually a result of autoerotic stimulation or may be associated with psychiatric disorders [207].

Iatrogenic injury is the most common type of urethral trauma [208, 209]. The incidence of urethral injury during transurethral catheterisation is 6.7 per 1,000 catheters inserted [210], and can occur due to creation of a false passage by the tip of the catheter, inadvertent inflation of the anchoring balloon in the urethra or removal of the catheter with the anchoring balloon not fully deflated [210]. A strict indication for every urethral catheterisation is an important preventive measure [208]. The importance of catheter insertion training programmes, to prevent urethral injury during transurethral catheterisation, have been demonstrated [211, 212]. Instrumentation of the urethra (TURP, cystoscopy, etc.) can traumatise all segments of it [208].

4.4.1.2 *Posterior male urethral injuries*

Blunt posterior urethral injuries are almost exclusively related to pelvic fractures with disruption of the pelvic ring [208, 209]. These injuries are referred to as pelvic fracture urethral injuries (PFUI) [202, 213], and are mainly caused by MVAs [214]. Pelvic fracture urethral injuries are divided into partial or complete ruptures [214]. In complete ruptures, there is a gap between the disrupted ends of the urethra, which fills up with scar tissue. There is no urethral wall in the scarred space and any lumen represents a fistulous tract between the urethral stumps [215]. Injuries of the bladder neck and prostate are rare and mostly occur at the anterior midline of both the bladder neck and prostatic urethra [216]. It is highly uncommon to find a complete transection of the bladder neck or an avulsion of the anterior part of the prostate [216]. Concomitant injuries to the head, thorax, abdomen and/or spine are frequent (up to 66%) [214].

Penetrating injuries of the pelvis, perineum or buttocks (mainly gunshot wounds) can also damage the posterior urethra, but are extremely rare in the civilian setting [217]. There is a high probability of associated injuries (80-90%), mainly intra-abdominal [161, 217].

The associated injuries which occur with both blunt and penetrating posterior urethral injuries can be life-threatening, and if so, will govern the patient's assessment and treatment [214]. Delayed morbidities of posterior urethral injuries include strictures, incontinence and erectile dysfunction, all of which

may have a detrimental effect on the patient's quality of life [218]. Erectile dysfunction occurs in 34% (25-45%) of patients with PFUI [219].

4.4.1.3 *Female urethral injuries*

Birth related injuries to the female urethra are rare and consist of minor (peri)urethral lacerations during vaginal delivery. Pelvic fractures are the main cause of blunt trauma [218, 220]; however, PFUIs in females are rare and less common than in males. This is usually attributed to the flexibility provided by the vagina and the greater inherent elasticity of the female urethra [218, 220], it may also be the result of less severe and more frequent stable pelvic fractures in females [154, 214]. In unstable pelvic fractures in females, a high suspicion for a urethral injury should be maintained [220]. Female urethral injuries are classified into two types: longitudinal or partial (most frequent) injuries and transverse or complete injuries [220]. Concomitant bladder or vaginal injury is possible; therefore, females are at risk of developing urinary incontinence and urethrovaginal fistula [214, 220].

Insertion of a synthetic sub-urethral sling for the treatment of female stress urinary incontinence is complicated by an intra-operative urethral injury in 0.2-2.5% of cases [221] and is an important cause of **iatrogenic** urethral injury.

4.4.2 **Evaluation**

4.4.2.1 *Clinical signs*

Blood at the meatus is the cardinal sign, but the absence of it doesn't rule out a urethral injury [154, 214]. Inability to void (with a palpable distended bladder) is another classic sign and is often associated with a complete rupture [214, 215]. Haematuria and pain on urination may be present in incomplete ruptures. Urinary extravasation and bleeding may result in scrotal, penile and/or perineal swelling and ecchymosis, depending on the location and extent of the trauma. The presentation of these clinical symptoms may be delayed (> 1 hour) [215].

Rectal examination should always be done to exclude an associated rectal injury (up to 5% of cases), and may reveal a 'high-riding' prostate, which is an unreliable finding [154, 215]. Failure to detect a rectal injury can cause significant morbidity and even mortality. A rectal injury is suggested by blood on the examining finger and/or a palpable laceration [154]. Another sign of urethral injury is difficulty or inability to pass a urethral catheter [154, 215].

A female urethral injury should be suspected from the combination of a (unstable) pelvic fracture with blood at the vaginal introitus, vaginal laceration, haematuria, urethrorrhagia, labial swelling, urinary retention or difficulties passing a urethral catheter [154, 218]. Vaginal examination is indicated to assess vaginal lacerations [154, 218].

4.4.2.2 *Urethrography*

Retrograde urethrography (RUG) is the standard in the early evaluation of a male urethral injury [154, 222] and is conducted by injecting 20-30 mL of contrast material while occluding the meatus. Films should be taken in a 30° oblique position. In patients with PFUI, it is important to move the X-ray beam to the 30° angle rather than the patient [214]. In an unstable patient, RUG should be postponed until the patient has been stabilised [154, 161].

During RUG, any extravasation outside the urethra is pathognomonic for urethral injury [215]. A typical image for incomplete rupture shows extravasation from the urethra which occurs while the bladder is still filling. A complete rupture is suggested by massive extravasation without bladder filling [214]. Although RUG is able to reliably identify the site of injury (anterior vs. posterior), the distinction between a complete and partial rupture is not always clear [213, 214]. Therefore, any proposed classification system based on RUG is not reliable [213, 214]. In females, the short urethra and vulvar oedema makes adequate urethrography nearly impossible [223].

Prior to deferred treatment, a combination of RUG and antegrade cysto-urethrography is the standard to evaluate site and extent of the urethral stenosis, and to evaluate the competence of the bladder neck [214].

4.4.2.3 *Cysto-urethroscopy*

Flexible cysto-urethroscopy is a valuable alternative to diagnose an acute urethral injury and may distinguish between complete and partial rupture [222]. Flexible cysto-urethroscopy is preferred to RUG in suspected penile fracture-associated urethral injury as RUG is associated with a high false-negative rate [224, 225]. In females, where the short urethra often precludes adequate radiological visualisation, cysto-urethroscopy and vaginoscopy are the diagnostic modalities of choice [154, 220]. If, prior to deferred treatment, the competence of the bladder neck is not clear upon antegrade cysto-urethrography, a suprapubic cystoscopy is advised [214].

4.4.2.4 *Ultrasound and magnetic resonance imaging*

In the acute phase, US scanning is used for guiding the placement of a suprapubic catheter [214]. In complex PFUIs, MRI before deferred treatment provides valuable additional information, which can help to determine the most appropriate surgical strategy [226]. This information includes a better estimation of the length of the distraction defect, degree of prostatic displacement and presence/absence of a false passage [226].

4.4.3 **Disease Management**

4.4.3.1 *Male anterior urethral injuries*

4.4.3.1.1 Immediate exploration and urethral reconstruction

This is indicated for penile fracture related injuries [204] and non-life threatening penetrating injuries [218]. Small lacerations can be repaired by simple closure [204]. Complete ruptures without extensive tissue loss are treated with anastomotic repair [204, 205]. In the case of longer defects or apparent infection (particularly bite wounds), a staged repair with urethral marsupialisation is needed [222].

Penetrating injuries require peri- and post-operative antibiotic treatment [227]. The role of immediate urethroplasty in blunt injuries is controversial. Patients (88.3% complete ruptures) who underwent immediate urethroplasty had a failure rate that was not significantly different compared to those who underwent delayed urethroplasty after initial suprapubic diversion (11.7% vs. 18.6%; $p=0.71$). The time to spontaneous voiding was significantly shorter in the immediate urethroplasty group (27 vs. 192 days) [228].

4.4.3.1.2 Urinary diversion

Blunt anterior urethral injuries are associated with spongiosal contusion. Evaluation of the limits of urethral debridement in the acute phase might be difficult and as a consequence, it is reasonable to start with urinary diversion only [222]. If urinary diversion is performed, the therapeutic options are suprapubic diversion or a trial of early endoscopic re-alignment with transurethral catheterisation [222]. One study reported a better outcome for suprapubic diversion compared to early endoscopic re-alignment and transurethral catheterisation [229]. Urinary diversion is maintained for one to two and three weeks for partial and complete ruptures, respectively [222, 229]. Satisfactory urethral luminal re-canalisation may occur in up to 68% after partial ruptures, but is rare (14%) after complete ruptures [229]. Transurethral or suprapubic urinary diversion are treatment options for iatrogenic or life-threatening penetrating injuries [218, 230]. Minor iatrogenic urethral injuries and urethral contusions do not require urinary diversion [3].

4.4.3.2 *Male posterior urethral injuries*

4.4.3.2.1 Emergency room management

As these injuries are usually associated with other severe injuries, resuscitation and immediate treatment of life-threatening injuries have absolute priority [214]. Penetrating injuries especially have a very high likelihood of associated injuries requiring immediate exploration [161, 217]. There is no urgency to treat the urethral injury and urinary diversion is not essential during the first hours after trauma [215]; however, it is preferable to establish early urinary diversion to:

- monitor urinary output, since this is a valuable sign of the haemodynamic condition and the renal function of the patient;
- treat symptomatic retention if the patient is still conscious;
- minimise urinary extravasation and its secondary effects, such as infection and fibrosis [214].

Insertion of a suprapubic catheter is an accepted practice in urgent situations [215, 217]. However, insertion of a suprapubic catheter is not without risk, especially in the unstable trauma patient where the bladder is often displaced by the pelvic haematoma or because of poor bladder filling due to haemodynamic shock or concomitant bladder injury. In these circumstances, an attempt at urethral catheterisation can be carried out by experienced personnel. It is extremely unlikely that the gentle passage of a urethral catheter will do any additional damage [214]. If there is any difficulty, a suprapubic catheter should be placed under US guidance or under direct vision for example, during laparotomy for associated injuries [214].

4.4.3.2.2 Early urethral management (less than six weeks after injury)

For partial injuries, urinary diversion (suprapubic or transurethral) is sufficient as these injuries can heal without significant scarring or obstruction [215, 218]. A complete injury will not heal, and formation of an obliterated segment is inevitable in case of suprapubic diversion alone [215, 218]. To avoid this obliteration and a long period of suprapubic diversion followed by deferred urethroplasty, the urethral ends can be sutured (urethroplasty) or approximated over a transurethral catheter (re-alignment).

4.4.3.2.2.1 Immediate urethroplasty

Urethroplasty within 48 hours after injury is difficult because of poor visualisation and the inability to accurately

assess the degree of urethral disruption, due to extensive swelling and ecchymosis, which may result in extensive unjustified urethral debridement. Another problem is the risk of severe bleeding (average 3 L) following entry into the pelvic haematoma [214]. In addition, with high rates of impotence (23%), incontinence (14%) and strictures (54%), urethroplasty within 48 hours is not indicated [214].

4.4.3.2.2 Early urethroplasty

Urethroplasty can be performed after two days and up to six weeks after the initial injury, if associated injuries have been stabilised, the distraction defect is short, the perineum is soft and the patient is able to lie down in the lithotomy position [231, 232]. This avoids a long period of suprapubic diversion with its discomfort and complications [231, 232]. As the results (complications, stricture recurrence, incontinence and impotence) are equivalent to delayed urethroplasty [232-234], early urethroplasty might be an option for patients fulfilling the above-mentioned conditions.

Lacerations (blunt or penetrating) at the bladder neck and prostatic urethra are a specific entity: they will never heal spontaneously, will cause local cavitation (and source of infection) and compromise the intrinsic sphincter mechanism (with risk of urinary incontinence) [216]. They must be reconstructed as soon as possible [213, 217, 218]. For penetrating injuries with severe lesions to the prostate, prostatectomy (with bladder neck sparing) must be performed [217].

4.4.3.2.3 Early re-alignment

Early re-alignment can be performed when a stable patient is on the operating table for other surgery or as a stand-alone procedure in the absence of concomitant injuries [161, 235]. In a partial injury, re-alignment, and transurethral catheterisation avoids extravasation of urine in the surrounding tissues reducing the inflammatory response. In complete injuries, the aim of re-alignment is to correct severe distraction injuries rather than to prevent a stricture [218, 236].

Re-alignment can be done by an open or endoscopic technique [236, 237], the endoscopic technique is preferred as it is less invasive and allows for direct visual control [236]. Using a flexible/rigid cystoscope and biplanar fluoroscopy, a guidewire is placed inside the bladder, over this a catheter is placed. If necessary, two cystoscopes can be used: one retrograde (per urethra) and one antegrade (suprapubic route through the bladder neck) [214]. The duration of catheterisation is three weeks for partial and six weeks for complete ruptures with voiding urethrography upon catheter removal [214]. It is important to avoid traction on the balloon catheter as it can damage the remaining sphincter mechanism at the bladder neck [214].

With contemporary endoscopic re-alignment procedures, stricture formation is reduced to 44-49% [236, 237] as compared to a 89-94% stricture rate with suprapubic diversion [237, 238]. There is no evidence that early re-alignment increases the risk of urinary incontinence (4.7-5.8%) or erectile dysfunction (16.7-20.5%) [237, 238].

Another potential benefit of early re-alignment is that when a stricture occurs it will be shorter and therefore, easier to treat. For short, non-obliterative strictures following re-alignment, direct vision urethrotomy can be performed. Approximately 50% of strictures after endoscopic re-alignment can be treated endoscopically [236]. However, repetitive endoscopic procedures in case of stricture formation might delay the time to definitive cure and can increase the incidence of adverse events (false passage, abscess formation) [239, 240]. In light of this, repetitive endoscopic treatments after failed re-alignment are not recommended; instead, urethroplasty must be performed.

Koraitim *et al.* found a shorter stricture length after early (open) re-alignment and as a consequence, a tendency for less complex manoeuvres to be needed to allow for a tension-free anastomosis during urethroplasty [241]. On the other hand, Tausch *et al.* reported an equal stricture length and no greater facilitation of urethroplasty after failed endoscopic re-alignment compared to suprapubic diversion only [239]. The proposed benefit is thus highly questionable. Furthermore, there is conflicting evidence as to whether failed early re-alignment jeopardises the success of definitive urethroplasty [214].

Differences between series in the rates of incontinence, impotence and re-stricture can be explained by differences in patient selection (severe vs. less severe trauma), a mix of partial and complete ruptures, and differences in follow-up duration. Furthermore, these differences make the comparison with other techniques difficult, especially with urethroplasty [154, 236].

4.4.3.2.3 Deferred management (greater than three months after injury)

The standard treatment remains deferred urethroplasty [13, 14]. In the case of a complete rupture, treated with an initial period of three months suprapubic diversion, obliteration of the posterior urethra is almost inevitable [215]. Endoscopic treatment of a complete obliteration is not successful [214]. After at least three months of suprapubic diversion, the pelvic haematoma is nearly always resolved, the prostate has descended into a more normal position, the scar tissue has stabilised [231] and the patient is clinically stable and able to lie down

in the lithotomy position [222, 231]. Associated life-threatening injuries often preclude early management of penetrating membranous urethral injuries. In those cases, suprapubic diversion with delayed urethroplasty is also advised [17, 25, 26]. Perineal anastomotic repair is the surgical technique of choice, but a combined abdominoperineal approach is necessary in rare cases of concomitant bladder neck injury or recto-urethral fistula [242].

The overall success rate for deferred urethroplasty is 86% [214]. Deferred urethroplasty does not significantly affect erectile function [243]. Although, a small proportion (<7%) of patients report *de novo* erectile dysfunction after delayed urethroplasty, others (6-20%) have recovery of erectile dysfunction after delayed urethroplasty [214]. Incontinence is rare with deferred urethroplasty (approximately 5%), and is usually due to incompetence of the bladder neck [214]. The assessment of sexual function and the decision on definitive treatment (e.g. penile prosthesis), should be undertaken two years after the trauma because of the potential return of potency within that time [213, 244].

4.4.3.3 Female urethral injuries

Emergency room management of PFUIs in females is the same as in males (section 4.4.3.2.1); however, subsequent management differs. Treatment options are [220]:

- **Early realignment:** This is associated with a high stricture and fistula rate.
- **Early repair (less than or equal to seven days):** Complication rate is the lowest with early repair; therefore, this strategy is preferred once the patient is hemodynamically stable [218, 220].
- **Delayed repair (greater than seven days):** Delayed repair often requires complex abdominal or combined abdominal-vaginal reconstruction with elevated risk of urinary incontinence and vaginal stenosis.

The approach (vaginal, abdominal or combined) for early repair depends on the location of the injury [220]. Proximal and mid-urethral disruptions require immediate exploration and primary repair using the retropubic and transvaginal routes, respectively, with primary suturing of the urethral ends or urethral laceration. Concomitant vaginal lacerations are repaired (2-layer closure) transvaginally at the same time [220]. Distal urethral injuries can be left hypospadiac since they do not disrupt the sphincter mechanism, but a concomitant vaginal laceration must be closed [154, 223]. In case of urethral injury during synthetic sub-urethral sling insertion, immediate repair is warranted with abortion of sling insertion [221].

Table 4.4.1: Complication rates for different treatment strategies for PFUIs in females [220]

Type of repair	Stricture (%)	Fistula (%)	Incontinence (%)	Vaginal stenosis (%)	Need for permanent urinary diversion (%)
Early realignment	59	13	0	0	0
Early repair	3	6	9	0	3
Delayed repair	3	4	31	4	7

4.4.4 **Summary of evidence and recommendations for the evaluation and management of urethral trauma**

Summary of evidence	LE
Implementing training programmes on urinary catheter insertion for personnel involved with urethral catheterisation significantly improves the rate of catheter-related complications.	2b
In males, a urethral injury is detected as contrast extravasation during urethrography or as a mucosal laceration during cysto-urethroscopy.	3
As opposed to cysto-urethroscopy, voiding cysto-urethrography will miss a female urethral injury in approximately 50% of cases.	3
Transurethral or suprapubic urinary diversion are the treatment options for iatrogenic injuries.	3
In males with blunt anterior urethral injuries, one study reported a worse outcome with transurethral catheterisation compared to suprapubic diversion.	3
If PFUIs are associated with life-threatening injuries, urethral management has no priority and urinary diversion with either urethral or suprapubic catheterisation is sufficient initially.	3
With early endoscopic re-alignment the stricture rate is reduced to 44-49% without increased risk of incontinence or erectile dysfunction.	3
Repetitive endoscopic treatments after failed re-alignment delays the time to definitive cure and increases the incidence of adverse events.	3
For partial posterior injuries, urinary diversion (suprapubic or transurethral) is sufficient as these injuries might heal without significant scarring or obstruction.	3
Immediate urethroplasty (< 48 hours) in male PFUI is associated with a higher risk of bleeding and stricture, incontinence and impotence rates compared to delayed urethroplasty.	3
Early urethroplasty (two days to six weeks) in selected patients for male PFUI is associated with similar stricture, incontinence and impotence rate compared to delayed urethroplasty.	3
Suprapubic diversion with delayed urethroplasty in male PFUI with complete urethral disruption is associated with a 86% stricture free success rate and with no significant impact on erectile function and urinary continence.	2a
Early repair in female PFUI has the lowest complication rate.	3

Recommendations	Strength rating
Provide appropriate training to reduce the risk of traumatic catheterisation.	Strong
Evaluate male urethral injuries with flexible cysto-urethroscopy and/or retrograde urethrography.	Strong
Evaluate female urethral injuries with cysto-urethroscopy and vaginoscopy.	Strong
Treat iatrogenic anterior urethral injuries by transurethral or suprapubic urinary diversion.	Strong
Treat blunt anterior urethral injuries in males by suprapubic diversion.	Weak
Treat pelvic fracture urethral injuries (PFUIs) in hemodynamically unstable patients by transurethral or suprapubic catheterisation initially.	Strong
Perform early endoscopic re-alignment in male PFUIs when feasible.	Weak
Do not repeat endoscopic treatments after failed re-alignment for male PFUI.	Strong
Treat partial posterior urethral injuries by suprapubic or transurethral catheter.	Strong
Do not perform immediate urethroplasty (< 48 hours) in male PFUIs.	Strong
Perform early urethroplasty (two days to six weeks) for male PFUIs with complete disruption in selected patients (stable, short gap, soft perineum, lithotomy position possible).	Weak
Manage complete posterior urethral disruption in male PFUIs with suprapubic diversion and deferred (at least three months) urethroplasty.	Strong
Perform early repair (within seven days) for female PFUIs (not delayed repair or early re-alignment).	Strong

4.4.5 Treatment algorithms

Management of anterior and posterior urethral injuries in men

Figure 4.4.1: Management of anterior urethral injuries in men





Figure 4.4.2: Management of posterior urethral injuries in men



RUG = retrograde urethrography; DVIU = direct visual internal urethrotomy.

4.5 Genital Trauma

4.5.1 Epidemiology, aetiology and pathophysiology

Of all urological injuries, 33-66% involve the external genitalia [245]. Genital trauma is much more common in males than in females, especially between the ages of 15 and 40 years. This is due to anatomical differences, increased frequency of road traffic accidents and increased participation in physical sports, war and crime. The risk of associated injuries to neighbouring organs (bladder, urethra, vagina, rectum and bowel), after blunt trauma is higher in females than in males.

Genital trauma is commonly caused by blunt injuries (80%). In males, blunt genital trauma frequently occurs unilaterally with approximately 1% presenting as bilateral scrotal or testicular injuries [246]. Any kind of contact sport, without the use of protective aids, may be associated with genital trauma. Off-road cycling, motor biking (especially on motorbikes with a dominant petrol tank), rugby, football and hockey are all activities associated with blunt testicular trauma [247-250].

Accidents during sexual intercourse can also cause genital trauma; men of younger age are the most affected. The major pathologies are penile fractures, strangulation and necrosis, and urethrovaginal foreign bodies resulting from autoeroticism practices [251].

The most important presentation of blunt penile trauma is penile fracture. The most common causes are sexual intercourse, forced flexion (taqaandan), masturbation and rolling over in 46%, 21%, 18 % and 8.2%, respectively [252]. The usual mechanism of injury is when the penis slips out of the vagina and strikes against the symphysis pubis or perineum. Sixty per cent of cases occur during consensual intercourse [253], with penile fracture more likely when the partner is on top. Penile fracture is caused by rupture of the cavernosal tunica albuginea, and may be associated with subcutaneous haematoma and lesions of the corpus spongiosum or urethra in 10-22% [254-256]. Genital injury is prevalent (42%) after sexual abuse [257].

Although animal bites are common, bites injuring the external genitalia are rare. Wounds are usually minor, but have a risk of wound infection. The most common bacterial infection caused by a dog bite is *Pasturella multocida*, which accounts for up to 50% of infections [258]. Other commonly involved organisms are *Escherichia coli*, *Streptococcus viridans*, *Staphylococcus aureus*, *Eikenella corrodens*, *Capnocytophaga canimorsus*, *Veillonella parvula*, *Bacteroides* and *Fusobacterium* spp. [258-260].

Gunshot injuries to the external genitalia are relatively uncommon and are usually not life-threatening; however, they can have a significant impact on quality of life. About 40-60% of all penetrating genito-urinary lesions involve the external genitalia [206, 261], 35% of these are gunshot wounds [246]. In a series of wartime injuries, the majority were caused by improvised explosive devices and other explosive ordinance, while smaller numbers of injuries were due to gunshot injuries [262]. In both males and females, penetrating injuries affect multiple organs in 70% of patients. In males, penetrating scrotal injuries affect both testes in 30% of cases compared with 1% in blunt injuries [246, 263]. Self-mutilation of the external genitalia has also been reported in psychotic patients and transsexuals [259]. Genital burns are rare in isolation and are usually due to industrial flames or chemicals [264]. Both male and female genital piercings increase the risk for unexpected genital trauma [265].

Traumatic dislocation of the testicle rarely occurs and is most common in victims of MVAs [266-269]. Bilateral dislocation of the testes has been reported in up to 25% of cases [267]. Testicular rupture is found in approximately 50% of cases of direct blunt scrotal trauma [270, 271]. It may occur under intense compression of the testis against the inferior pubic ramus or symphysis, resulting in a rupture of the tunica albuginea. A force of approximately 50 kg is necessary to cause testicular rupture [272]. Most penile avulsion injuries are self-inflicted, but some are a result of industrial accidents or assault.

Coital injury of the female genital tract can happen during consensual sexual intercourse. Up to 35% of all genital injuries in women are sustained during their first sexual contact. The most frequently found injuries are lacerations [273]. Blunt trauma to the vulva is rarely reported and usually presents as a large haematoma. The incidence of traumatic vulvar haematomas after vaginal deliveries has been reported as 1 in 310 deliveries [274]. The presence of a vulvar haematoma is closely related to an increased risk of associated vaginal, pelvic or abdominal injuries [275, 276]. Blunt injuries of the vulva and vagina are associated with pelvic trauma in 30%, after consensual intercourse in 25%, following sexual assault in 20%, and other blunt trauma in 15% [277].

4.5.2 **Diagnostic evaluation**

4.5.2.1 *Patient history and physical examination*

Penile fracture is associated with a sudden cracking or popping sound, pain and immediate detumescence. Local swelling of the penile shaft develops quickly, due to enlarging haematoma. Bleeding may spread along the fascial layers of the penile shaft and extend to the lower abdominal wall if Buck's fascia is also ruptured. Sometimes, the rupture of the tunica may be palpable. Less severe penile injuries can be distinguished from penile fracture, as they are not usually associated with detumescence [252].

Testicular rupture is associated with immediate pain, nausea, vomiting, and sometimes fainting. The hemiscrotum is tender, swollen, and ecchymotic. The testis itself may be difficult to palpate. **Blunt vulvar** or perineal trauma in women may be associated with bleeding, pain and voiding problems, bladder catheterisation is usually required.

In genital trauma, a urinalysis should be performed. The presence of visible haematuria requires a retrograde urethrogram in males. In females, flexible or rigid cystoscopy is recommended to exclude urethral and bladder injury [275, 277]. In women with genital injuries and blood at the vaginal introitus, further gynaecological investigation is needed [275].

4.5.3 **Imaging**

In cases of suspected penile fracture cavernosography, US or MRI [252, 278-280] can identify lacerations of the tunica albuginea in unclear cases [281], or provide reassurance that the tunica is intact. MRI is superior to US in diagnosing penile fracture [282]. If a concomitant urethral injury is suspected, manage as outlined in section 4.4.

Ultrasound should be performed to determine intra- and/or extra-testicular haematoma, testicular contusion, or rupture [271, 283-291]. However, the literature is contradictory as to the usefulness of US compared to clinical examination alone. Some studies have reported convincing findings with a specificity of up to 98.6% [292]. Heterogeneous echo pattern of the testicular parenchyma with the loss of contour definition is a highly sensitive and specific radiographic finding for testicular rupture [282]. Others reported poor specificity (78%) and sensitivity (28%) for the differentiation between testicular rupture and haematocele, while accuracy is as low as 56% [284]. Colour Doppler-duplex US may provide useful information when used to evaluate testicular perfusion. If scrotal US is inconclusive, testicular CT or MRI may be helpful [293]; however, these techniques did not specifically increase the detection rates of testicular rupture.

4.5.4 **Disease management**

4.5.4.1 *Animal bites*

Local wound management depends on the extent of tissue destruction. Antibiotics should be prescribed in accordance with local resistance patterns [294-296]. The possibility of rabies infection must be considered taking into account the geographical location, animal involved, specific nature of the wound and the type of attack (provoked/unprovoked). Elderly and immunosuppressed patients should be vaccinated with human rabies immunoglobulin and human diploid cell vaccine [297, 298].

4.5.4.2 *Human bites*

In cases of human bites, apart from wound management, infection should be considered since transmission of viral diseases may occur, Hepatitis B vaccine/immunoglobulin and/or immunodeficiency virus (HIV) post-exposure prophylaxis should be offered. For further details, see Guidelines for the Management of Human Bite Injuries [299].

4.5.4.3 *Blunt penile trauma*

Blunt trauma to the flaccid penis does not usually cause tearing of the tunica. Subcutaneous haematoma after sexual intercourse, without associated rupture of the cavernosal tunica albuginea, does not require surgical intervention. In these cases, non-steroidal analgesics and ice-packs are recommended [300].

4.5.4.4 *Penile fracture*

The thickness of the tunica albuginea in the flaccid state (approximately 2 mm) decreases in erection to 0.25-0.5mm, and is therefore more vulnerable to traumatic injury [292, 301]. When a penile fracture is diagnosed, surgical intervention with closure of the tunica albuginea is recommended; it ensures the lowest rate of negative long-term sequelae and has no negative effect on the psychological wellbeing of the patient [302]. The approach is usually through a circumferential incision proximal to the coronal sulcus which enables complete degloving of the penis. Increasingly, local longitudinal incisions centred on the area of fracture or ventral longitudinal approaches are currently used [224]. Further localisation may be gained with a flexible cystoscopy performed prior to incision, if urethral trauma is suspected and eventually proven. Surgical closure of the tunica should be carried out using absorbable sutures.

4.5.4.5 *Penetrating penile trauma*

In penetrating penile trauma non-operative management is recommended for small superficial injuries with intact Buck's fascia [206]. In more significant penetrating penile injuries, surgical exploration and debridement of necrotic tissue is recommended. Even in extended injuries of the penis, primary alignment of the disrupted tissues may allow for acceptable healing because of the robust penile blood supply [259].

The principles of care are debridement of devitalised tissue, with the preservation of as much viable tissues as possible, haemostasis, diversion of urine in selected cases and the removal of foreign bodies. Tissues of questionable viability may be left for subsequent definitive surgery. If a delayed repair is needed, depending on the type of injury and the extent of tissue damage, it usually takes place four to six weeks after the trauma has occurred.

The surgical approach depends upon the site and extent of the injury, but a subcoronal incision with penile degloving usually gives good exposure. Initially, a defect in the tunica albuginea should be closed after copious irrigation. If there has been too much tissue loss, the defect can be repaired either immediately or after delay with a patch (either from an autologous saphenous vein or xenograft).

The elasticity of genital skin means it is usually possible to manage the loss of a moderate amount of penile skin; however, management is more difficult in extensive injuries with significant skin loss. The tissue chosen for reconstruction following trauma needs to provide good coverage and must be suitable for reconstruction. Split-thickness skin grafting provides good coverage and a dependable take that is reproducible and durable. However, split-thickness grafts contract more than full-thickness grafts and their use on the penile shaft should be kept to a minimum. Skin grafts with thickness of at least 0.4 mm should be

used in order to reduce the risk of contraction [259]. Full-thickness skin grafting onto the penile shaft gives less contracture, a better cosmetic appearance and more resistance to trauma during intercourse, when re-established [300]. The donor site may be taken from the abdomen, buttock, thigh or axilla and is chosen according to surgeon's preference and the pattern of injury. In cases of extensive destruction of deeper tissues, or if later prosthetic placement is being considered, skin flaps, with their secure vascular supply, can be used.

4.5.4.6 *Penile avulsion injuries and amputation*

Acute management involves resuscitation of the patient, and preparation for surgical re-implantation of the penis if it has been recovered and is not too badly damaged. Surgical re-implantation should be considered for all patients and should be performed within 24 hours of amputation [303].

The severed penis should be washed with sterile saline, wrapped in saline-soaked gauze, placed in a sterile bag and immersed in iced water. The penis must not come into direct contact with the ice. A pressure dressing or a tourniquet should be placed around the penile stump to prevent excessive blood loss. Re-attachment can be achieved in a non-microsurgical way, but gives higher rates of post-operative urethral stricture and more problems with loss of sensation [304]. When operating microscopically, the corpora cavernosa and urethra are firstly aligned and repaired. Subsequently, the dorsal penile arteries, the dorsal vein and the dorsal nerves are anastomosed. The cavernosal arteries are generally too small to anastomose. The fascia and skin are closed in layers and both a urethral and a suprapubic catheter are placed.

If the severed penis cannot be found, or is unsuitable for re-attachment, then the end should be closed as it is done in partial penectomy. Later reconstruction may be employed to lengthen the penis (e.g. suspensory ligament division and V-Y plasty, pseudo-glans formation with split-thickness skin grafting, etc.). A delayed major reconstructive procedure, i.e. phalloplasty (either radial artery or pubic), is sometimes required for injuries which leave a very small or non-functioning penile stump [303].

4.5.4.7 *Testicular dislocation*

It can be either a subcutaneous dislocation with epifascial displacement of the testis or an internal dislocation. In the latter, the testis is positioned in the superficial external inguinal ring, inguinal canal or abdominal cavity. Traumatic dislocation of the testis is treated by manual replacement and secondary orchidopexy. If primary manual reposition cannot be performed, immediate orchidopexy is indicated.

4.5.4.8 *Haematocoele*

Conservative management is recommended in haematocoeles smaller than three times the size of the contralateral testis [305]. In large haematocoeles, non-operative management can fail, and delayed surgery (more than three days) is often required. Patients with large haematocoeles have a higher rate of orchiectomy than patients who undergo early surgery, even in non-ruptured testes [246, 259, 270, 306, 307]. Early surgical intervention results in preservation of the testis in more than 90% of cases compared to delayed surgeries which result in orchiectomy in 45-55% of patients [270]. In addition, non-operative management is also associated with prolonged hospital stays. Therefore, large haematocoeles should be treated surgically, irrespective of the presence of testicular contusion or rupture. At the very least, the blood clot should be evacuated from the tunica vaginalis sac to relieve disability and hasten recovery.

4.5.4.9 *Testicular rupture*

It is essential to surgically explore equivocal patients whenever imaging studies cannot definitively exclude testicular rupture. This involves exploration with evacuation of blood clots and haematoma, excision of any necrotic testicular tubules and closure of the tunica albuginea, usually with running 3.0-absorbable sutures.

4.5.4.10 *Penetrating scrotal trauma*

Penetrating injuries to the scrotum require surgical exploration with debridement of non-viable tissue. Depending on the extent of the injury, primary reconstruction of the testis and scrotum can usually be performed. In complete disruption of the spermatic cord, re-alignment without vaso-vasostomy may be considered if surgically feasible [308]. Staged secondary microsurgical vaso-vasostomy can be performed after rehabilitation; although, only a few cases have been reported [308]. If there is extensive destruction of the tunica albuginea, mobilisation of a free tunica vaginalis flap can be performed for testicular closure. If the patient is unstable or reconstruction cannot be achieved, orchiectomy is then indicated. Prophylactic antibiotics are recommended after scrotal penetrating trauma, although data to support this approach is lacking. Tetanus prophylaxis is mandatory.

Extended laceration of scrotal skin requires surgical intervention for skin closure. Due to the elasticity of the scrotum, most defects can be primarily closed, even if the lacerated skin is only minimally attached to the scrotum [259]. Local wound management with extensive initial wound debridement and washout is important for scrotal convalescence. In the case of extensive loss of genital tissue, e.g. improvised

explosive devices blast injury, complex and staged reconstructive surgical procedures are often required [262].

Table 4.5.1: Summary of key points for penile fracture and testicular trauma

Summary of key points:
Penile fracture
The most common causes of penile fracture are sexual intercourse, forced flexion, masturbation and rolling over.
Penile fracture is associated with a sudden cracking or popping sound, pain, immediate detumescence and local swelling.
Magnetic resonance imaging is superior to all other imaging techniques in diagnosing penile fracture.
Management of penile fracture is surgical intervention with closure of the tunica albuginea.
Testicular Trauma
Blunt testicular injury may occur under intense compression of the testis against the inferior pubic ramus or symphysis, resulting in a rupture of the tunica albuginea.
Testicular rupture is associated with immediate pain, nausea, vomiting, and sometimes fainting.
Scrotal ultrasound is the preferred imaging modality for the diagnosis of testicular trauma.
Surgical exploration in patients with testicular trauma ensures preservation of viable tissue when possible.

4.5.5 **Complications**

The possibility of complications from genital trauma, including psychological effects, erectile dysfunction, urethral stricture, and infertility is high. In patients with a history of penile fracture post-operative complications were reported in up to 20% of cases, development of plaques or nodules following surgery, post-operative curvature formation and erectile dysfunction occur in 13.9%, 2.8% and 1.9% of patients, respectively [252]. Conservative management of penile fracture increases complications, such as penile abscess, missed urethral disruption, penile curvature, and persistent haematoma requiring delayed surgical intervention [309]. Late complications after conservative management were fibrosis and angulations in 35% and impotence in up to 62% [253, 309].

Post-operative complications were reported in 8% of patients who underwent testicular repair after penetrating trauma [206]. Despite good management and regular follow up of external genital gunshot wounds, such wounds are fraught with the possibility of complications such as erectile dysfunction, urethral stricture, and infertility. Delayed complications include chronic pain and testicular atrophy. Haematoceles initially treated non-operatively may eventually need delayed surgery if they develop infection or undue pain. Genital injuries are rarely life threatening, but fertility and testosterone production often become the male trauma patient's chief concern once acute issues are resolved [310].

4.5.6 **Follow up**

In patients with genital trauma follow up should focus on diagnosis of and therapy for late complications. Erectile dysfunction, urethral stricture and assessment of fertility are the main concerns [256, 311].

4.5.7 **Summary of evidence and recommendations for evaluation and management of genital trauma.**

Summary of evidence	LE
A concomitant urethral injury complicates penile fractures and requires specialised management.	3
Ultrasound can determine intra- and/or extra-testicular haematoma, testicular contusion, or rupture with heterogeneous echo pattern parenchyma and loss of contour definition regarded as a highly sensitive and specific finding.	3
Surgical treatment of penile fracture ensures the lowest rate of negative long-term sequelae on functional and psychological wellbeing of the patient.	3
In patients with testicular rupture or equivocal imaging, surgical exploration can secure preservation of viable tissue.	3

Recommendations	Strength rating
Exclude urethral injury in the case of penile fracture.	Strong
Perform ultrasound (US) for the diagnosis of testis trauma.	Strong
Treat penile fractures surgically, with closure of tunica albuginea.	Strong
Explore the injured testis in all cases of testicular rupture and in those with inconclusive US findings.	Strong

5. REFERENCES

- Radmayr, C., *et al.*, EAU Guidelines on Paediatric Urology. In: EAU Guidelines, edition presented at the annual EAU Congress Barcelona 2019. ISBN 978-94-92671-04-2.
<https://uroweb.org/guideline/paediatric-urology/>
- Martinez-Pineiro, L., *et al.* EAU Guidelines on Urethral Trauma. *Eur Urol*, 2010. 57: 791.
<https://www.ncbi.nlm.nih.gov/pubmed/20122789>
- Summerton, D.J., *et al.* EAU guidelines on iatrogenic trauma. *Eur Urol*, 2012. 62: 628.
<https://www.ncbi.nlm.nih.gov/pubmed/22717550>
- Lumen, N., *et al.* Review of the current management of lower urinary tract injuries by the EAU Trauma Guidelines Panel. *Eur Urol*, 2015. 67: 925.
<https://www.ncbi.nlm.nih.gov/pubmed/25576009>
- Serafetinides, E., *et al.* Review of the current management of upper urinary tract injuries by the EAU Trauma Guidelines Panel. *Eur Urol*, 2015. 67: 930.
<https://www.ncbi.nlm.nih.gov/pubmed/25578621>
- Guyatt, G.H., *et al.* GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*, 2008. 336: 924.
<https://www.ncbi.nlm.nih.gov/pubmed/18436948>
- Guyatt, G.H., *et al.* What is "quality of evidence" and why is it important to clinicians? *BMJ*, 2008. 336: 995.
<https://www.ncbi.nlm.nih.gov/pubmed/18456631>
- Phillips B, *et al.* Oxford Centre for Evidence-based Medicine Levels of Evidence. Updated by Jeremy Howick March 2009. 1998.
<https://www.cebm.net/2009/06/oxford-centre-evidence-based-medicine-levels-evidence-march-2009/>
- Guyatt, G.H., *et al.* Going from evidence to recommendations. *BMJ*, 2008. 336: 1049.
<https://www.ncbi.nlm.nih.gov/pubmed/18467413>
- Soreide, K. Epidemiology of major trauma. *Br J Surg*, 2009. 96: 697.
<https://www.ncbi.nlm.nih.gov/pubmed/19526611>
- Middleton, P., The trauma epidemic. In: Major Trauma. Smith, J., Greaves, I., Porter, K. (2010) Oxford University Press: Oxford.
- Thornley, S., *et al.* Alcohol intake, marijuana use, and sleep deprivation on the risk of falls occurring at home among young and middle-aged adults: a case-crossover study. *N Z Med J*, 2014. 127: 32.
<https://www.ncbi.nlm.nih.gov/pubmed/25447247>
- Moore, E.E., *et al.* Organ injury scaling: spleen, liver, and kidney. *J Trauma*, 1989. 29: 1664.
<https://www.ncbi.nlm.nih.gov/pubmed/2593197>
- Monstrey, S.J., *et al.* Urological trauma and severe associated injuries. *Br J Urol*, 1987. 60: 393.
<https://www.ncbi.nlm.nih.gov/pubmed/3427315>
- MacKenzie, E.J., *et al.* A national evaluation of the effect of trauma-center care on mortality. *N Engl J Med*, 2006. 354: 366.
<https://www.ncbi.nlm.nih.gov/pubmed/16436768>
- Caterson, E.J., *et al.* Boston bombings: a surgical view of lessons learned from combat casualty care and the applicability to Boston's terrorist attack. *J Craniofac Surg*, 2013. 24: 1061.
<https://www.ncbi.nlm.nih.gov/pubmed/23851738>
- Feliciano DV, M.E., Mattox KL. , Trauma damage control, In: Trauma, F.D. Mattox KL, Moore EE, Editors. 2000, McGraw-Hill: New York.
- Hirshberg, A., *et al.* 'Damage control' in trauma surgery. *Br J Surg*, 1993. 80: 1501.
<https://www.ncbi.nlm.nih.gov/pubmed/8298911>
- Rignault, D.P. Recent progress in surgery for the victims of disaster, terrorism, and war--Introduction. *World J Surg*, 1992. 16: 885.
<https://www.ncbi.nlm.nih.gov/pubmed/1462624>

20. Rotondo, M.F., *et al.* 'Damage control': an approach for improved survival in exsanguinating penetrating abdominal injury. *J Trauma*, 1993. 35: 375.
<https://www.ncbi.nlm.nih.gov/pubmed/8371295>
21. Slater, M.S., *et al.* Terrorism in America. An evolving threat. *Arch Surg*, 1997. 132: 1059.
<https://www.ncbi.nlm.nih.gov/pubmed/9336502>
22. Frykberg, E.R. Medical management of disasters and mass casualties from terrorist bombings: how can we cope? *J Trauma*, 2002. 53: 201.
<https://www.ncbi.nlm.nih.gov/pubmed/12169923>
23. Jacobs, L.M., Jr., *et al.* An emergency medical system approach to disaster planning. *J Trauma*, 1979. 19: 157.
<https://www.ncbi.nlm.nih.gov/pubmed/458880>
24. Eberle, B.M., *et al.* Thromboembolic prophylaxis with low-molecular-weight heparin in patients with blunt solid abdominal organ injuries undergoing nonoperative management: current practice and outcomes. *J Trauma*, 2011. 70: 141.
<https://www.ncbi.nlm.nih.gov/pubmed/21217492>
25. Barrera, L.M., *et al.* Thromboprophylaxis for trauma patients. *Cochrane Database Syst Rev*, 2013: CD008303.
<https://www.ncbi.nlm.nih.gov/pubmed/23543562>
26. Meng, M.V., *et al.* Renal trauma: indications and techniques for surgical exploration. *World J Urol*, 1999. 17: 71.
<https://www.ncbi.nlm.nih.gov/pubmed/10367364>
27. Wessells, H., *et al.* Renal injury and operative management in the United States: results of a population-based study. *J Trauma*, 2003. 54: 423.
<https://www.ncbi.nlm.nih.gov/pubmed/12634519>
28. Santucci, R.A., *et al.* The literature increasingly supports expectant (conservative) management of renal trauma--a systematic review. *J Trauma*, 2005. 59: 493.
<https://www.ncbi.nlm.nih.gov/pubmed/16294101>
29. Sujenthiran, A., *et al.* Is Nonoperative Management the Best First-line Option for High-grade Renal trauma? A Systematic Review. *Eur Urol Focus*, 2017.
<https://www.ncbi.nlm.nih.gov/pubmed/28753890>
30. Mingoli, A., *et al.* Operative and nonoperative management for renal trauma: comparison of outcomes. A systematic review and meta-analysis. *Ther Clin Risk Manag*, 2017. 13: 1127.
<https://www.ncbi.nlm.nih.gov/pubmed/28894376>
31. Bjurlin, M.A., *et al.* Comparison of nonoperative management with renorrhaphy and nephrectomy in penetrating renal injuries. *J Trauma*, 2011. 71: 554.
<https://www.ncbi.nlm.nih.gov/pubmed/21610541>
32. Santucci, R.A., *et al.* Evaluation and management of renal injuries: consensus statement of the renal trauma subcommittee. *BJU Int*, 2004. 93: 937.
<https://www.ncbi.nlm.nih.gov/pubmed/15142141>
33. Kansas, B.T., *et al.* Incidence and management of penetrating renal trauma in patients with multiorgan injury: extended experience at an inner city trauma center. *J Urol*, 2004. 172: 1355.
<https://www.ncbi.nlm.nih.gov/pubmed/15371841>
34. Najibi, S., *et al.* Civilian gunshot wounds to the genitourinary tract: incidence, anatomic distribution, associated injuries, and outcomes. *Urology*, 2010. 76: 977.
<https://www.ncbi.nlm.nih.gov/pubmed/20605196>
35. Shariat, S.F., *et al.* Evidence-based validation of the predictive value of the American Association for the Surgery of Trauma kidney injury scale. *J Trauma*, 2007. 62: 933.
<https://www.ncbi.nlm.nih.gov/pubmed/17426551>
36. Santucci, R.A., *et al.* Validation of the American Association for the Surgery of Trauma organ injury severity scale for the kidney. *J Trauma*, 2001. 50: 195.
<https://www.ncbi.nlm.nih.gov/pubmed/11242281>
37. Malaeb, B., *et al.* Should blunt segmental vascular renal injuries be considered an American Association for the Surgery of Trauma Grade 4 renal injury? *J Trauma Acute Care Surg*, 2014. 76: 484.
<https://www.ncbi.nlm.nih.gov/pubmed/24458054>
38. Sierink, J.C., *et al.* Systematic review and meta-analysis of immediate total-body computed tomography compared with selective radiological imaging of injured patients. *Br J Surg*, 2012. 99 Suppl 1: 52.
<https://www.ncbi.nlm.nih.gov/pubmed/22441856>

39. Huber-Wagner, S., *et al.* Effect of whole-body CT during trauma resuscitation on survival: a retrospective, multicentre study. *Lancet*, 2009. 373: 1455.
<https://www.ncbi.nlm.nih.gov/pubmed/19321199>
40. Cachecho, R., *et al.* Management of the trauma patient with pre-existing renal disease. *Crit Care Clin*, 1994. 10: 523.
<https://www.ncbi.nlm.nih.gov/pubmed/7922736>
41. Cozar, J.M., *et al.* [Management of injury of the solitary kidney]. *Arch Esp Urol*, 1990. 43: 15.
<https://www.ncbi.nlm.nih.gov/pubmed/2331159>
42. Sebastia, M.C., *et al.* Renal trauma in occult ureteropelvic junction obstruction: CT findings. *Eur Radiol*, 1999. 9: 611.
<https://www.ncbi.nlm.nih.gov/pubmed/10354870>
43. Buchberger, W., *et al.* [Diagnosis and staging of blunt kidney trauma. A comparison of urinalysis, i.v. urography, sonography and computed tomography]. *Rofo*, 1993. 158: 507.
<https://www.ncbi.nlm.nih.gov/pubmed/8507839>
44. Carroll, P.R., *et al.* Renovascular trauma: risk assessment, surgical management, and outcome. *J Trauma*, 1990. 30: 547.
<https://www.ncbi.nlm.nih.gov/pubmed/2342137>
45. Eastham, J.A., *et al.* Radiographic evaluation of adult patients with blunt renal trauma. *J Urol*, 1992. 148: 266.
<https://www.ncbi.nlm.nih.gov/pubmed/1635113>
46. Schmidlin, F.R., *et al.* The higher injury risk of abnormal kidneys in blunt renal trauma. *Scand J Urol Nephrol*, 1998. 32: 388.
<https://www.ncbi.nlm.nih.gov/pubmed/9925001>
47. Chandhoke, P.S., *et al.* Detection and significance of microscopic hematuria in patients with blunt renal trauma. *J Urol*, 1988. 140: 16.
<https://www.ncbi.nlm.nih.gov/pubmed/3379684>
48. Heyns, C.F. Renal trauma: indications for imaging and surgical exploration. *BJU Int*, 2004. 93: 1165.
<https://www.ncbi.nlm.nih.gov/pubmed/15142132>
49. Sheth, S. *et al.* ACR Appropriateness Criteria® Renal trauma. 2012.
<https://acsearch.acr.org/docs/69373/Narrative/>
50. Morey, A.F., *et al.* Urotrauma: AUA guideline. *J Urol*, 2014. 192: 327.
<https://www.ncbi.nlm.nih.gov/pubmed/24857651>
51. McCombie, S.P., *et al.* The conservative management of renal trauma: a literature review and practical clinical guideline from Australia and New Zealand. *BJU Int*, 2014. 114 Suppl 1: 13.
<https://www.ncbi.nlm.nih.gov/pubmed/25124459>
52. Heller, M.T., *et al.* MDCT of renal trauma: correlation to AAST organ injury scale. *Clin Imaging*, 2014. 38: 410.
<https://www.ncbi.nlm.nih.gov/pubmed/24667041>
53. Fischer, W., *et al.* JOURNAL CLUB: Incidence of Urinary Leak and Diagnostic Yield of Excretory Phase CT in the Setting of Renal Trauma. *AJR Am J Roentgenol*, 2015. 204: 1168.
<https://www.ncbi.nlm.nih.gov/pubmed/26001225>
54. Colling, K.P., *et al.* Computed tomography scans with intravenous contrast: low incidence of contrast-induced nephropathy in blunt trauma patients. *J Trauma Acute Care Surg*, 2014. 77: 226.
<https://www.ncbi.nlm.nih.gov/pubmed/25058246>
55. Valentino, M., *et al.* Contrast-enhanced US evaluation in patients with blunt abdominal trauma(). *J Ultrasound*, 2010. 13: 22.
<https://www.ncbi.nlm.nih.gov/pubmed/23396012>
56. Mihalik, J.E., *et al.* The use of contrast-enhanced ultrasound for the evaluation of solid abdominal organ injury in patients with blunt abdominal trauma. *J Trauma Acute Care Surg*, 2012. 73: 1100.
<https://www.ncbi.nlm.nih.gov/pubmed/22832765>
57. Cagini, L., *et al.* Contrast enhanced ultrasound (CEUS) in blunt abdominal trauma. *Crit Ultrasound J*, 2013. 5 Suppl 1: S9.
<https://www.ncbi.nlm.nih.gov/pubmed/23902930>
58. Morey, A.F., *et al.* Single shot intraoperative excretory urography for the immediate evaluation of renal trauma. *J Urol*, 1999. 161: 1088.
<https://www.ncbi.nlm.nih.gov/pubmed/10081844>
59. Ku, J.H., *et al.* Is there a role for magnetic resonance imaging in renal trauma? *Int J Urol*, 2001. 8: 261.
<https://www.ncbi.nlm.nih.gov/pubmed/11389740>

60. Leppaniemi, A., *et al.* MRI and CT in blunt renal trauma: an update. *Semin Ultrasound CT MR*, 1997. 18: 129.
<https://www.ncbi.nlm.nih.gov/pubmed/9163832>
61. Wessells, H., *et al.* Preservation of renal function after reconstruction for trauma: quantitative assessment with radionuclide scintigraphy. *J Urol*, 1997. 157: 1583.
<https://www.ncbi.nlm.nih.gov/pubmed/9112481>
62. Schmidlin, F.R., *et al.* [The conservative treatment of major kidney injuries]. *Ann Urol (Paris)*, 1997. 31: 246.
<https://www.ncbi.nlm.nih.gov/pubmed/9480627>
63. Thall, E.H., *et al.* Conservative management of penetrating and blunt Type III renal injuries. *Br J Urol*, 1996. 77: 512.
<https://www.ncbi.nlm.nih.gov/pubmed/8777609>
64. Alsikafi, N.F., *et al.* Nonoperative management outcomes of isolated urinary extravasation following renal lacerations due to external trauma. *J Urol*, 2006. 176: 2494.
<https://www.ncbi.nlm.nih.gov/pubmed/17085140>
65. Buckley, J.C., *et al.* Selective management of isolated and nonisolated grade IV renal injuries. *J Urol*, 2006. 176: 2498.
<https://www.ncbi.nlm.nih.gov/pubmed/17085141>
66. Haas, C.A., *et al.* Use of ureteral stents in the management of major renal trauma with urinary extravasation: is there a role? *J Endourol*, 1998. 12: 545.
<https://www.ncbi.nlm.nih.gov/pubmed/9895260>
67. Moudouni, S.M., *et al.* Management of major blunt renal lacerations: is a nonoperative approach indicated? *Eur Urol*, 2001. 40: 409.
<https://www.ncbi.nlm.nih.gov/pubmed/11713395>
68. Keihani, S., *et al.* Contemporary management of high-grade renal trauma: Results from the American Association for the Surgery of Trauma Genitourinary Trauma study. *J Trauma Acute Care Surg*, 2018. 84: 418.
<https://www.ncbi.nlm.nih.gov/pubmed/29298242>
69. Elliott, S.P., *et al.* Renal arterial injuries: a single center analysis of management strategies and outcomes. *J Urol*, 2007. 178: 2451.
<https://www.ncbi.nlm.nih.gov/pubmed/17937955>
70. Sartorelli, K.H., *et al.* Nonoperative management of hepatic, splenic, and renal injuries in adults with multiple injuries. *J Trauma*, 2000. 49: 56.
<https://www.ncbi.nlm.nih.gov/pubmed/10912858>
71. Toutouzas, K.G., *et al.* Nonoperative management of blunt renal trauma: a prospective study. *Am Surg*, 2002. 68: 1097.
<https://www.ncbi.nlm.nih.gov/pubmed/12516817>
72. Dugi, D.D., 3rd, *et al.* American Association for the Surgery of Trauma grade 4 renal injury substratification into grades 4a (low risk) and 4b (high risk). *J Urol*, 2010. 183: 592.
<https://www.ncbi.nlm.nih.gov/pubmed/20018329>
73. Hammer, C.C., *et al.* Effect of an institutional policy of nonoperative treatment of grades I to IV renal injuries. *J Urol*, 2003. 169: 1751.
<https://www.ncbi.nlm.nih.gov/pubmed/12686825>
74. Jawas, A., *et al.* Management algorithm for complete blunt renal artery occlusion in multiple trauma patients: case series. *Int J Surg*, 2008. 6: 317.
<https://www.ncbi.nlm.nih.gov/pubmed/18590988>
75. Armenakas, N.A., *et al.* Indications for nonoperative management of renal stab wounds. *J Urol*, 1999. 161: 768.
<https://www.ncbi.nlm.nih.gov/pubmed/10022681>
76. Jansen, J.O., *et al.* Selective non-operative management of abdominal gunshot wounds: survey of practise. *Injury*, 2013. 44: 639.
<https://www.ncbi.nlm.nih.gov/pubmed/22341771>
77. Bernath, A.S., *et al.* Stab wounds of the kidney: conservative management in flank penetration. *J Urol*, 1983. 129: 468.
<https://www.ncbi.nlm.nih.gov/pubmed/6834529>
78. Wessells, H., *et al.* Criteria for nonoperative treatment of significant penetrating renal lacerations. *J Urol*, 1997. 157: 24.
<https://www.ncbi.nlm.nih.gov/pubmed/8976207>
79. DuBose, J., *et al.* Selective non-operative management of solid organ injury following abdominal gunshot wounds. *Injury*, 2007. 38: 1084.
<https://www.ncbi.nlm.nih.gov/pubmed/17544428>

80. Shefler, A., *et al.* [The role of nonoperative management of penetrating renal trauma]. *Harefuah*, 2007. 146: 345.
<https://www.ncbi.nlm.nih.gov/pubmed/17674549>
81. Hope, W.W., *et al.* Non-operative management in penetrating abdominal trauma: is it feasible at a Level II trauma center? *J Emerg Med*, 2012. 43: 190.
<https://www.ncbi.nlm.nih.gov/pubmed/22051843>
82. Raza, S.J., *et al.* Outcomes of renal salvage for penetrating renal trauma: a single institution experience. *Can J Urol*, 2018. 25: 9323.
<https://www.ncbi.nlm.nih.gov/pubmed/29900820>
83. Lanchon, C., *et al.* High Grade Blunt Renal Trauma: Predictors of Surgery and Long-Term Outcomes of Conservative Management. A Prospective Single Center Study. *J Urol*, 2016. 195: 106.
<https://www.ncbi.nlm.nih.gov/pubmed/26254724>
84. Shoobridge, J.J., *et al.* A 9-year experience of renal injury at an Australian level 1 trauma centre. *BJU Int*, 2013. 112 Suppl 2: 53.
<https://www.ncbi.nlm.nih.gov/pubmed/23418742>
85. van der Wilden, G.M., *et al.* Successful nonoperative management of the most severe blunt renal injuries: a multicenter study of the research consortium of New England Centers for Trauma. *JAMA Surg*, 2013. 148: 924.
<https://www.ncbi.nlm.nih.gov/pubmed/23945834>
86. Charbit, J., *et al.* What are the specific computed tomography scan criteria that can predict or exclude the need for renal angioembolization after high-grade renal trauma in a conservative management strategy? *J Trauma*, 2011. 70: 1219.
<https://www.ncbi.nlm.nih.gov/pubmed/23945834>
87. Lin, W.C., *et al.* Computed tomographic imaging in determining the need of embolization for high-grade blunt renal injury. *J Trauma Acute Care Surg*, 2013. 74: 230.
<https://www.ncbi.nlm.nih.gov/pubmed/23271099>
88. Huber, J., *et al.* Selective transarterial embolization for posttraumatic renal hemorrhage: a second try is worthwhile. *J Urol*, 2011. 185: 1751.
<https://www.ncbi.nlm.nih.gov/pubmed/21420122>
89. Hotaling, J.M., *et al.* Analysis of diagnostic angiography and angioembolization in the acute management of renal trauma using a national data set. *J Urol*, 2011. 185: 1316.
<https://www.ncbi.nlm.nih.gov/pubmed/21334643>
90. Saour, M., *et al.* Effect of renal angioembolization on post-traumatic acute kidney injury after high-grade renal trauma: a comparative study of 52 consecutive cases. *Injury*, 2014. 45: 894.
<https://www.ncbi.nlm.nih.gov/pubmed/24456608>
91. Moolman, C., *et al.* Nonoperative management of penetrating kidney injuries: a prospective audit. *J Urol*, 2012. 188: 169.
<https://www.ncbi.nlm.nih.gov/pubmed/22591960>
92. Davis, P., *et al.* Assessing the usefulness of delayed imaging in routine followup for renal trauma. *J Urol*, 2010. 184: 973.
<https://www.ncbi.nlm.nih.gov/pubmed/20643462>
93. Hadjipavlou, M., *et al.* Managing penetrating renal trauma: experience from two major trauma centres in the UK. *BJU Int*, 2018. 121: 928.
<https://www.ncbi.nlm.nih.gov/pubmed/29438587>
94. Husmann, D.A., *et al.* Major renal lacerations with a devitalized fragment following blunt abdominal trauma: a comparison between nonoperative (expectant) versus surgical management. *J Urol*, 1993. 150: 1774.
<https://www.ncbi.nlm.nih.gov/pubmed/24072011>
95. McAninch, J.W., *et al.* Renal reconstruction after injury. *J Urol*, 1991. 145: 932.
<https://www.ncbi.nlm.nih.gov/pubmed/2016804>
96. Robert, M., *et al.* Management of major blunt renal lacerations: surgical or nonoperative approach? *Eur Urol*, 1996. 30: 335.
<https://www.ncbi.nlm.nih.gov/pubmed/8931966>
97. Nash, P.A., *et al.* Nephrectomy for traumatic renal injuries. *J Urol*, 1995. 153: 609.
<https://www.ncbi.nlm.nih.gov/pubmed/7861494>
98. Gonzalez, R.P., *et al.* Surgical management of renal trauma: is vascular control necessary? *J Trauma*, 1999. 47: 1039.
<https://www.ncbi.nlm.nih.gov/pubmed/10608530>
99. Rostas, J., *et al.* Intraoperative management of renal gunshot injuries: is mandatory exploration of Gerota's fascia necessary? *Am J Surg*, 2016. 211: 783.
<https://www.ncbi.nlm.nih.gov/pubmed/26867480>

100. Davis, K.A., *et al.* Predictors of the need for nephrectomy after renal trauma. *J Trauma*, 2006. 60: 164.
<https://www.ncbi.nlm.nih.gov/pubmed/16456451>
101. Wright, J.L., *et al.* Renal and extrarenal predictors of nephrectomy from the national trauma data bank. *J Urol*, 2006. 175: 970.
<https://www.ncbi.nlm.nih.gov/pubmed/16469594>
102. DiGiacomo, J.C., *et al.* The role of nephrectomy in the acutely injured. *Arch Surg*, 2001. 136: 1045.
<https://www.ncbi.nlm.nih.gov/pubmed/11529828>
103. Brandes, S.B., *et al.* Reconstructive surgery for trauma of the upper urinary tract. *Urol Clin North Am*, 1999. 26: 183.
<https://www.ncbi.nlm.nih.gov/pubmed/10086060>
104. Shekarriz, B., *et al.* The use of fibrin sealant in urology. *J Urol*, 2002. 167: 1218.
<https://www.ncbi.nlm.nih.gov/pubmed/11832701>
105. Knudson, M.M., *et al.* Outcome after major renovascular injuries: a Western trauma association multicenter report. *J Trauma*, 2000. 49: 1116.
<https://www.ncbi.nlm.nih.gov/pubmed/11130498>
106. Tillou, A., *et al.* Renal vascular injuries. *Surg Clin North Am*, 2001. 81: 1417.
<https://www.ncbi.nlm.nih.gov/pubmed/11766183>
107. Tasian, G.E., *et al.* Evaluation of renal function after major renal injury: correlation with the American Association for the Surgery of Trauma Injury Scale. *J Urol*, 2010. 183: 196.
<https://www.ncbi.nlm.nih.gov/pubmed/19913819>
108. Fiard, G., *et al.* Long-term renal function assessment with dimercapto-succinic acid scintigraphy after conservative treatment of major renal trauma. *J Urol*, 2012. 187: 1306.
<https://www.ncbi.nlm.nih.gov/pubmed/22341289>
109. Montgomery, R.C., *et al.* Posttraumatic renovascular hypertension after occult renal injury. *J Trauma*, 1998. 45: 106.
<https://www.ncbi.nlm.nih.gov/pubmed/9680021>
110. Heyns, C.F., *et al.* Increasing role of angiography and segmental artery embolization in the management of renal stab wounds. *J Urol*, 1992. 147: 1231.
<https://www.ncbi.nlm.nih.gov/pubmed/1569655>
111. Monstrey, S.J., *et al.* Renal trauma and hypertension. *J Trauma*, 1989. 29: 65.
<https://www.ncbi.nlm.nih.gov/pubmed/2911106>
112. Lebech, A., *et al.* [Hypertension following blunt kidney injury]. *Ugeskr Laeger*, 1990. 152: 994.
<https://www.ncbi.nlm.nih.gov/pubmed/2183457>
113. Wang, K.T., *et al.* Late development of renal arteriovenous fistula following gunshot trauma--a case report. *Angiology*, 1998. 49: 415.
<https://www.ncbi.nlm.nih.gov/pubmed/9591535>
114. Elliott, S.P., *et al.* Ureteral injuries: external and iatrogenic. *Urol Clin North Am*, 2006. 33: 55.
<https://www.ncbi.nlm.nih.gov/pubmed/16488280>
115. Blackwell, R.H., *et al.* Complications of Recognized and Unrecognized Iatrogenic Ureteral Injury at Time of Hysterectomy: A Population Based Analysis. *J Urol*, 2018. 199: 1540.
<https://www.ncbi.nlm.nih.gov/pubmed/29408429>
116. Pereira, B.M., *et al.* A review of ureteral injuries after external trauma. *Scand J Trauma Resusc Emerg Med*, 2010. 18: 6.
<https://www.ncbi.nlm.nih.gov/pubmed/20128905>
117. McGeady, J.B., *et al.* Current epidemiology of genitourinary trauma. *Urol Clin North Am*, 2013. 40: 323.
<https://www.ncbi.nlm.nih.gov/pubmed/23905930>
118. Siram, S.M., *et al.* Ureteral trauma: patterns and mechanisms of injury of an uncommon condition. *Am J Surg*, 2010. 199: 566.
<https://www.ncbi.nlm.nih.gov/pubmed/20359576>
119. Serkin, F.B., *et al.* Combat urologic trauma in US military overseas contingency operations. *J Trauma*, 2010. 69 Suppl 1: S175.
<https://www.ncbi.nlm.nih.gov/pubmed/20622614>
120. Brandes, S., *et al.* Diagnosis and management of ureteric injury: an evidence-based analysis. *BJU Int*, 2004. 94: 277.
<https://www.ncbi.nlm.nih.gov/pubmed/15291852>
121. Chou, M.T., *et al.* Prophylactic ureteral catheterization in gynecologic surgery: a 12-year randomized trial in a community hospital. *Int Urogynecol J Pelvic Floor Dysfunct*, 2009. 20: 689.
<https://www.ncbi.nlm.nih.gov/pubmed/19165412>

122. Delacroix, S.E., Jr., *et al.* Urinary tract injuries: recognition and management. *Clin Colon Rectal Surg*, 2010. 23: 104.
<https://www.ncbi.nlm.nih.gov/pubmed/21629628>
123. Visco, A.G., *et al.* Cost-effectiveness of universal cystoscopy to identify ureteral injury at hysterectomy. *Obstet Gynecol*, 2001. 97: 685.
<https://www.ncbi.nlm.nih.gov/pubmed/11339916>
124. Halabi, W.J., *et al.* Ureteral injuries in colorectal surgery: an analysis of trends, outcomes, and risk factors over a 10-year period in the United States. *Dis Colon Rectum*, 2014. 57: 179.
<https://www.ncbi.nlm.nih.gov/pubmed/24401879>
125. Johnson, D.B., *et al.* Complications of ureteroscopy. *Urol Clin North Am*, 2004. 31: 157.
<https://www.ncbi.nlm.nih.gov/pubmed/15040412>
126. Petersen, S.S., *et al.* Rate of Urologic Injury with Robotic Hysterectomy. *J Minim Invasive Gynecol*, 2018. 25: 867.
<https://www.ncbi.nlm.nih.gov/pubmed/29337210>
127. Schimpf, M.O., *et al.* Universal ureteral stent placement at hysterectomy to identify ureteral injury: a decision analysis. *BJOG*, 2008. 115: 1151.
<https://www.ncbi.nlm.nih.gov/pubmed/18518875>
128. Hesselman, S., *et al.* Effect of remote cesarean delivery on complications during hysterectomy: a cohort study. *Am J Obstet Gynecol*, 2017. 217: 564 e1.
<https://www.ncbi.nlm.nih.gov/pubmed/28735704>
129. Gilmour, D.T., *et al.* Rates of urinary tract injury from gynecologic surgery and the role of intraoperative cystoscopy. *Obstet Gynecol*, 2006. 107: 1366.
<https://www.ncbi.nlm.nih.gov/pubmed/16738165>
130. Wu, H.H., *et al.* The detection of ureteral injuries after hysterectomy. *J Minim Invasive Gynecol*, 2006. 13: 403.
<https://www.ncbi.nlm.nih.gov/pubmed/16962522>
131. Pokala, N., *et al.* A randomized controlled trial comparing simultaneous intra-operative vs sequential prophylactic ureteric catheter insertion in re-operative and complicated colorectal surgery. *Int J Colorectal Dis*, 2007. 22: 683.
<https://www.ncbi.nlm.nih.gov/pubmed/17031654>
132. Jhaveri, J.K., *et al.* Ureteral injuries sustained during robot-assisted radical prostatectomy. *J Endourol*, 2014. 28: 318.
<https://www.ncbi.nlm.nih.gov/pubmed/24147874>
133. Kunkle, D.A., *et al.* Delayed diagnosis of traumatic ureteral injuries. *J Urol*, 2006. 176: 2503.
<https://www.ncbi.nlm.nih.gov/pubmed/17085143>
134. Parpala-Sparman, T., *et al.* Increasing numbers of ureteric injuries after the introduction of laparoscopic surgery. *Scand J Urol Nephrol*, 2008. 42: 422.
<https://www.ncbi.nlm.nih.gov/pubmed/18609278>
135. Medina, D., *et al.* Ureteral trauma: preoperative studies neither predict injury nor prevent missed injuries. *J Am Coll Surg*, 1998. 186: 641.
<https://www.ncbi.nlm.nih.gov/pubmed/9632150>
136. Lucarelli, G., *et al.* Delayed relief of ureteral obstruction is implicated in the long-term development of renal damage and arterial hypertension in patients with unilateral ureteral injury. *J Urol*, 2013. 189: 960.
<https://www.ncbi.nlm.nih.gov/pubmed/23017525>
137. Alabousi, A., *et al.* Multi-modality imaging of the leaking ureter: why does detection of traumatic and iatrogenic ureteral injuries remain a challenge? *Emerg Radiol*, 2017. 24: 417.
<https://www.ncbi.nlm.nih.gov/pubmed/28451770>
138. Speicher, P.J., *et al.* Ureteral stenting in laparoscopic colorectal surgery. *J Surg Res*, 2014. 190: 98.
<https://www.ncbi.nlm.nih.gov/pubmed/24656474>
139. Coakley, K.M., *et al.* Prophylactic Ureteral Catheters for Colectomy: A National Surgical Quality Improvement Program-Based Analysis. *Dis Colon Rectum*, 2018. 61: 84.
<https://www.ncbi.nlm.nih.gov/pubmed/29215477>
140. Hassinger, T.E., *et al.* Ureteral stents increase risk of postoperative acute kidney injury following colorectal surgery. *Surg Endosc*, 2018. 32: 3342.
<https://www.ncbi.nlm.nih.gov/pubmed/29340810>
141. Smith, T.G., 3rd, *et al.* Damage control maneuvers for urologic trauma. *Urol Clin North Am*, 2013. 40: 343.
<https://www.ncbi.nlm.nih.gov/pubmed/23905932>

142. Koukouras, D., *et al.* Percutaneous minimally invasive management of iatrogenic ureteral injuries. *J Endourol*, 2010. 24: 1921.
<https://www.ncbi.nlm.nih.gov/pubmed/20964484>
143. El Abd, A.S., *et al.* Immediate and late management of iatrogenic ureteric injuries: 28 years of experience. *Arab J Urol*, 2015. 13: 250.
<https://www.ncbi.nlm.nih.gov/pubmed/26609443>
144. Png, J.C., *et al.* Principles of ureteric reconstruction. *Curr Opin Urol*, 2000. 10: 207.
<https://www.ncbi.nlm.nih.gov/pubmed/10858898>
145. Khan, F., *et al.* Management of ureteropelvic junction obstruction in adults. *Nat Rev Urol*, 2014. 11: 629.
<https://www.ncbi.nlm.nih.gov/pubmed/25287785>
146. Burks, F.N., *et al.* Management of iatrogenic ureteral injury. *Ther Adv Urol*, 2014. 6: 115.
<https://www.ncbi.nlm.nih.gov/pubmed/24883109>
147. Wenske, S., *et al.* Outcomes of distal ureteral reconstruction through reimplantation with psoas hitch, Boari flap, or ureteroneocystostomy for benign or malignant ureteral obstruction or injury. *Urology*, 2013. 82: 231.
<https://www.ncbi.nlm.nih.gov/pubmed/23642933>
148. Chung, B.I., *et al.* The use of bowel for ureteral replacement for complex ureteral reconstruction: long-term results. *J Urol*, 2006. 175: 179.
<https://www.ncbi.nlm.nih.gov/pubmed/16406903>
149. Armatys, S.A., *et al.* Use of ileum as ureteral replacement in urological reconstruction. *J Urol*, 2009. 181: 177.
<https://www.ncbi.nlm.nih.gov/pubmed/19013597>
150. Meng, M.V., *et al.* Expanded experience with laparoscopic nephrectomy and autotransplantation for severe ureteral injury. *J Urol*, 2003. 169: 1363.
<https://www.ncbi.nlm.nih.gov/pubmed/12629362>
151. Decaestecker, K., *et al.* Robot-assisted Kidney Autotransplantation: A Minimally Invasive Way to Salvage Kidneys. *Eur Urol Focus*, 2018. 4: 198.
<https://www.ncbi.nlm.nih.gov/pubmed/30093358>
152. Zhao, L.C., *et al.* Robotic Ureteral Reconstruction Using Buccal Mucosa Grafts: A Multi-institutional Experience. *Eur Urol*, 2017.
<https://www.ncbi.nlm.nih.gov/pubmed/29239749>
153. Pereira, B.M., *et al.* Bladder injuries after external trauma: 20 years experience report in a population-based cross-sectional view. *World J Urol*, 2013. 31: 913.
<https://www.ncbi.nlm.nih.gov/pubmed/22544337>
154. Figler, B.D., *et al.* Multi-disciplinary update on pelvic fracture associated bladder and urethral injuries. *Injury*, 2012. 43: 1242.
<https://www.ncbi.nlm.nih.gov/pubmed/22592152>
155. Wirth, G.J., *et al.* Advances in the management of blunt traumatic bladder rupture: experience with 36 cases. *BJU Int*, 2010. 106: 1344.
<https://www.ncbi.nlm.nih.gov/pubmed/20438556>
156. Deibert, C.M., *et al.* The association between operative repair of bladder injury and improved survival: results from the National Trauma Data Bank. *J Urol*, 2011. 186: 151.
<https://www.ncbi.nlm.nih.gov/pubmed/21575961>
157. Matlock, K.A., *et al.* Blunt traumatic bladder rupture: a 10-year perspective. *Am Surg*, 2013. 79: 589.
<https://www.ncbi.nlm.nih.gov/pubmed/23711268>
158. Johnsen, N.V., *et al.* Epidemiology of Blunt Lower Urinary Tract Trauma With and Without Pelvic Fracture. *Urology*, 2017. 102: 234.
<https://www.ncbi.nlm.nih.gov/pubmed/28043650>
159. Johnsen, N.V., *et al.* Evaluating the Role of Operative Repair of Extraperitoneal Bladder Rupture Following Blunt Pelvic Trauma. *J Urol*, 2016. 195: 661.
<https://www.ncbi.nlm.nih.gov/pubmed/26318983>
160. Urry, R.J., *et al.* The incidence, spectrum and outcomes of traumatic bladder injuries within the Pietermaritzburg Metropolitan Trauma Service. *Injury*, 2016. 47: 1057.
<https://www.ncbi.nlm.nih.gov/pubmed/26854075>
161. Cinman, N.M., *et al.* Gunshot wounds to the lower urinary tract: a single-institution experience. *J Trauma Acute Care Surg*, 2013. 74: 725.
<https://www.ncbi.nlm.nih.gov/pubmed/23425728>
162. Al-Azzawi, I.S., *et al.* Lower genitourinary trauma in modern warfare: the experience from civil violence in Iraq. *Injury*, 2014. 45: 885.
<https://www.ncbi.nlm.nih.gov/pubmed/24485550>

163. Williams, M., *et al.* Management of combat-related urological trauma in the modern era. *Nat Rev Urol*, 2013. 10: 504.
<https://www.ncbi.nlm.nih.gov/pubmed/23877722>
164. Cordon, B.H., *et al.* Iatrogenic nonendoscopic bladder injuries over 24 years: 127 cases at a single institution. *Urology*, 2014. 84: 222.
<https://www.ncbi.nlm.nih.gov/pubmed/24857278>
165. Ford, A.A., *et al.* Mid-urethral sling operations for stress urinary incontinence in women. *Cochrane Database Syst Rev*, 2017. 7: CD006375.
<https://www.ncbi.nlm.nih.gov/pubmed/28756647>
166. Golan, S., *et al.* Transurethral resection of bladder tumour complicated by perforation requiring open surgical repair - clinical characteristics and oncological outcomes. *BJU Int*, 2011. 107: 1065.
<https://www.ncbi.nlm.nih.gov/pubmed/20860654>
167. El Hayek, O.R., *et al.* Evaluation of the incidence of bladder perforation after transurethral bladder tumor resection in a residency setting. *J Endourol*, 2009. 23: 1183.
<https://www.ncbi.nlm.nih.gov/pubmed/19530900>
168. Sugihara, T., *et al.* Comparison of perioperative outcomes including severe bladder injury between monopolar and bipolar transurethral resection of bladder tumors: a population based comparison. *J Urol*, 2014. 192: 1355.
<https://www.ncbi.nlm.nih.gov/pubmed/24893311>
169. Venkatramani, V., *et al.* Monopolar versus bipolar transurethral resection of bladder tumors: a single center, parallel arm, randomized, controlled trial. *J Urol*, 2014. 191: 1703.
<https://www.ncbi.nlm.nih.gov/pubmed/24333244>
170. Collado, A., *et al.* Early complications of endoscopic treatment for superficial bladder tumors. *J Urol*, 2000. 164: 1529.
<https://www.ncbi.nlm.nih.gov/pubmed/11025697>
171. Shazly, S.A., *et al.* Robotic radical hysterectomy in early stage cervical cancer: A systematic review and meta-analysis. *Gynecol Oncol*, 2015. 138: 457.
<https://www.ncbi.nlm.nih.gov/pubmed/26056752>
172. Brummer, T.H., *et al.* FINHYST, a prospective study of 5279 hysterectomies: complications and their risk factors. *Hum Reprod*, 2011. 26: 1741.
<https://www.ncbi.nlm.nih.gov/pubmed/21540244>
173. Billfeldt, N.K., *et al.* A Swedish population-based evaluation of benign hysterectomy, comparing minimally invasive and abdominal surgery. *Eur J Obstet Gynecol Reprod Biol*, 2018. 222: 113.
<https://www.ncbi.nlm.nih.gov/pubmed/29408741>
174. Tarney, C.M. Bladder Injury During Cesarean Delivery. *Curr Womens Health Rev*, 2013. 9: 70.
<https://www.ncbi.nlm.nih.gov/pubmed/24876830>
175. Honore, C., *et al.* HIPEC for peritoneal carcinomatosis: does an associated urologic procedure increase morbidity? *Ann Surg Oncol*, 2012. 19: 104.
<https://www.ncbi.nlm.nih.gov/pubmed/21638092>
176. Sawkar, H.P., *et al.* Frequency of lower urinary tract injury after gastrointestinal surgery in the nationwide inpatient sample database. *Am Surg*, 2014. 80: 1216.
<https://www.ncbi.nlm.nih.gov/pubmed/25513920>
177. Kockerling, F., *et al.* TEP versus TAPP: comparison of the perioperative outcome in 17,587 patients with a primary unilateral inguinal hernia. *Surg Endosc*, 2015. 29: 3750.
<https://www.ncbi.nlm.nih.gov/pubmed/25805239>
178. Balbay, M.D., *et al.* The actual incidence of bladder perforation following transurethral bladder surgery. *J Urol*, 2005. 174: 2260.
<https://www.ncbi.nlm.nih.gov/pubmed/16280794>
179. Nieder, A.M., *et al.* Transurethral bladder tumor resection: intraoperative and postoperative complications in a residency setting. *J Urol*, 2005. 174: 2307.
<https://www.ncbi.nlm.nih.gov/pubmed/16280830>
180. Welk, B.K., *et al.* Are male slings for post-prostatectomy incontinence a valid option? *Curr Opin Urol*, 2010. 20: 465.
<https://www.ncbi.nlm.nih.gov/pubmed/20838219>
181. Novara, G., *et al.* Updated systematic review and meta-analysis of the comparative data on colposuspensions, pubovaginal slings, and midurethral tapes in the surgical treatment of female stress urinary incontinence. *Eur Urol*, 2010. 58: 218.
<https://www.ncbi.nlm.nih.gov/pubmed/20434257>
182. Maher, C., *et al.* Transvaginal mesh or grafts compared with native tissue repair for vaginal prolapse. *Cochrane Database Syst Rev*, 2016. 2: CD012079.
<https://www.ncbi.nlm.nih.gov/pubmed/26858090>

183. Maher, C.F., *et al.* Laparoscopic sacral colpopexy versus total vaginal mesh for vaginal vault prolapse: a randomized trial. *Am J Obstet Gynecol*, 2011. 204: 360 e1.
<https://www.ncbi.nlm.nih.gov/pubmed/21306698>
184. Ogah, J., *et al.* Minimally invasive synthetic suburethral sling operations for stress urinary incontinence in women: a short version Cochrane review. *Neurourol Urodyn*, 2011. 30: 284.
<https://www.ncbi.nlm.nih.gov/pubmed/21412819>
185. Pereira, B.M., *et al.* Penetrating bladder trauma: a high risk factor for associated rectal injury. *Adv Urol*, 2014. 2014: 386280.
<https://www.ncbi.nlm.nih.gov/pubmed/24527030>
186. Clarke-Pearson, D.L., *et al.* Complications of hysterectomy. *Obstet Gynecol*, 2013. 121: 654.
<https://www.ncbi.nlm.nih.gov/pubmed/23635631>
187. Manikandan, R., *et al.* Percutaneous peritoneal drainage for intraperitoneal bladder perforations during transurethral resection of bladder tumors. *J Endourol*, 2003. 17: 945.
<https://www.ncbi.nlm.nih.gov/pubmed/14744369>
188. Patel, B.N., *et al.* Imaging of iatrogenic complications of the urinary tract: kidneys, ureters, and bladder. *Radiol Clin North Am*, 2014. 52: 1101.
<https://www.ncbi.nlm.nih.gov/pubmed/25173661>
189. Lehnert, B.E., *et al.* Lower male genitourinary trauma: a pictorial review. *Emerg Radiol*, 2014. 21: 67.
<https://www.ncbi.nlm.nih.gov/pubmed/24052083>
190. Quagliano, P.V., *et al.* Diagnosis of blunt bladder injury: A prospective comparative study of computed tomography cystography and conventional retrograde cystography. *J Trauma*, 2006. 61: 410.
<https://www.ncbi.nlm.nih.gov/pubmed/16917459>
191. Ramchandani, P., *et al.* Imaging of genitourinary trauma. *AJR Am J Roentgenol*, 2009. 192: 1514.
<https://www.ncbi.nlm.nih.gov/pubmed/19457813>
192. Alperin, M., *et al.* Conservative management of postoperatively diagnosed cystotomy. *Urology*, 2009. 73: 1163 e17.
<https://www.ncbi.nlm.nih.gov/pubmed/18514295>
193. Teeluckdharry, B., *et al.* Urinary Tract Injury at Benign Gynecologic Surgery and the Role of Cystoscopy: A Systematic Review and Meta-analysis. *Obstet Gynecol*, 2015. 126: 1161.
<https://www.ncbi.nlm.nih.gov/pubmed/26551173>
194. Stember, D.S., *et al.* Outcomes of abdominal wall reservoir placement in inflatable penile prosthesis implantation: a safe and efficacious alternative to the space of Retzius. *J Sex Med*, 2014. 11: 605.
<https://www.ncbi.nlm.nih.gov/pubmed/24286533>
195. Oh, J.S., *et al.* Effectiveness of the combat pelvic protection system in the prevention of genital and urinary tract injuries: An observational study. *J Trauma Acute Care Surg*, 2015. 79: S193.
<https://www.ncbi.nlm.nih.gov/pubmed/26406430>
196. Pansadoro, A., *et al.* Conservative treatment of intraperitoneal bladder perforation during transurethral resection of bladder tumor. *Urology*, 2002. 60: 682.
<https://www.ncbi.nlm.nih.gov/pubmed/12385934>
197. Inaba, K., *et al.* Selective nonoperative management of torso gunshot wounds: when is it safe to discharge? *J Trauma*, 2010. 68: 1301.
<https://www.ncbi.nlm.nih.gov/pubmed/22341771>
198. Lee, J.S., *et al.* Urologic complications following obstetric and gynecologic surgery. *Korean J Urol*, 2012. 53: 795.
<https://www.ncbi.nlm.nih.gov/pubmed/23185673>
199. Traxer, O., *et al.* Technique and complications of transurethral surgery for bladder tumours. *BJU Int*, 2004. 94: 492.
<https://www.ncbi.nlm.nih.gov/pubmed/15329099>
200. MacDonald, S., *et al.* Complications of Transvaginal Mesh for Pelvic Organ Prolapse and Stress Urinary Incontinence: Tips for Prevention, Recognition, and Management. *Eur Urol Focus*, 2016. 2: 260.
<https://www.ncbi.nlm.nih.gov/pubmed/28723371>
201. Inaba, K., *et al.* Prospective evaluation of the utility of routine postoperative cystogram after traumatic bladder injury. *J Trauma Acute Care Surg*, 2013. 75: 1019.
<https://www.ncbi.nlm.nih.gov/pubmed/24256676>
202. Latini, J.M., *et al.* SIU/ICUD Consultation On Urethral Strictures: Epidemiology, etiology, anatomy, and nomenclature of urethral stenoses, strictures, and pelvic fracture urethral disruption injuries. *Urology*, 2014. 83: S1.
<https://www.ncbi.nlm.nih.gov/pubmed/24210733>

203. Falcone, M., *et al.* Current Management of Penile Fracture: An Up-to-Date Systematic Review. *Sex Med Rev*, 2017.
<https://www.ncbi.nlm.nih.gov/pubmed/28874325>
204. Barros, R., *et al.* Primary urethral reconstruction results in penile fracture. *Ann R Coll Surg Engl*, 2018. 100: 21.
<https://www.ncbi.nlm.nih.gov/pubmed/29022780>
205. Bjurlin, M.A., *et al.* Clinical characteristics and surgical outcomes of penetrating external genital injuries. *J Trauma Acute Care Surg*, 2013. 74: 839.
<https://www.ncbi.nlm.nih.gov/pubmed/23425745>
206. Phonsombat, S., *et al.* Penetrating external genital trauma: a 30-year single institution experience. *J Urol*, 2008. 180: 192.
<https://www.ncbi.nlm.nih.gov/pubmed/18499189>
207. Ratkal, J.M., *et al.* Electric Wire as Foreign Body in the Bladder and Urethra-a Case Report and Review of Literature. *Indian J Surg*, 2015. 77: 1323.
<https://www.ncbi.nlm.nih.gov/pubmed/27011559>
208. Lumen, N., *et al.* Etiology of urethral stricture disease in the 21st century. *J Urol*, 2009. 182: 983.
<https://www.ncbi.nlm.nih.gov/pubmed/19616805>
209. Palminteri, E., *et al.* Contemporary urethral stricture characteristics in the developed world. *Urology*, 2013. 81: 191.
<https://www.ncbi.nlm.nih.gov/pubmed/23153951>
210. Davis, N.F., *et al.* Incidence, Cost, Complications and Clinical Outcomes of Iatrogenic Urethral Catheterization Injuries: A Prospective Multi-Institutional Study. *J Urol*, 2016. 196: 1473.
<https://www.ncbi.nlm.nih.gov/pubmed/27317985>
211. Bhatt, N.R., *et al.* A prospective audit on the effect of training and educational workshops on the incidence of urethral catheterization injuries. *Can Urol Assoc J*, 2017. 11: E302.
<https://www.ncbi.nlm.nih.gov/pubmed/28761592>
212. Kashefi, C., *et al.* Incidence and prevention of iatrogenic urethral injuries. *J Urol*, 2008. 179: 2254.
<https://www.ncbi.nlm.nih.gov/pubmed/18423712>
213. Gomez, R.G., *et al.* SIU/ICUD Consultation on Urethral Strictures: Pelvic fracture urethral injuries. *Urology*, 2014. 83: S48.
<https://www.ncbi.nlm.nih.gov/pubmed/24210734>
214. Barratt, R.C., *et al.* Pelvic fracture urethral injury in males-mechanisms of injury, management options and outcomes. *Transl Androl Urol*, 2018. 7: S29.
<https://www.ncbi.nlm.nih.gov/pubmed/29644168>
215. Mundy, A.R., *et al.* Urethral trauma. Part I: introduction, history, anatomy, pathology, assessment and emergency management. *BJU Int*, 2011. 108: 310.
<https://www.ncbi.nlm.nih.gov/pubmed/19874306>
216. Mundy, A.R., *et al.* Pelvic fracture-related injuries of the bladder neck and prostate: their nature, cause and management. *BJU Int*, 2010. 105: 1302.
<https://www.ncbi.nlm.nih.gov/pubmed/19874306>
217. Tausch, T.J., *et al.* Gunshot wound injuries of the prostate and posterior urethra: reconstructive armamentarium. *J Urol*, 2007. 178: 1346.
<https://www.ncbi.nlm.nih.gov/pubmed/17706720>
218. Mundy, A.R., *et al.* Urethral trauma. Part II: Types of injury and their management. *BJU Int*, 2011. 108: 630.
<https://www.ncbi.nlm.nih.gov/pubmed/19874306>
219. Benson, C.R., *et al.* Sexual Dysfunction in Urethral Reconstruction: A Review of the Literature. *Sex Med Rev*, 2017.
<https://www.ncbi.nlm.nih.gov/pubmed/29108977>
220. Patel, D.N., *et al.* Female urethral injuries associated with pelvic fracture: a systematic review of the literature. *BJU Int*, 2017. 120: 766.
<https://www.ncbi.nlm.nih.gov/pubmed/28805298>
221. Gomes, C.M., *et al.* Update on complications of synthetic suburethral slings. *Int Braz J Urol*, 2017. 43: 822.
<https://www.ncbi.nlm.nih.gov/pubmed/28266818>
222. Brandes, S. Initial management of anterior and posterior urethral injuries. *Urol Clin North Am*, 2006. 33: 87.
<https://www.ncbi.nlm.nih.gov/pubmed/16488283>
223. Black, P.C., *et al.* Urethral and bladder neck injury associated with pelvic fracture in 25 female patients. *J Urol*, 2006. 175: 2140.
<https://www.ncbi.nlm.nih.gov/pubmed/16697821>

224. Mazaris, E.M., *et al.* Penile fractures: immediate surgical approach with a midline ventral incision. *BJU Int*, 2009. 104: 520.
<https://www.ncbi.nlm.nih.gov/pubmed/16488283>
225. Kamdar, C., *et al.* Penile fracture: preoperative evaluation and surgical technique for optimal patient outcome. *BJU Int*, 2008. 102: 1640.
<https://www.ncbi.nlm.nih.gov/pubmed/18710448>
226. Horiguchi, A., *et al.* Pubourethral Stump Angle Measured on Preoperative Magnetic Resonance Imaging Predicts Urethroplasty Type for Pelvic Fracture Urethral Injury Repair. *Urology*, 2018. 112: 198.
<https://www.ncbi.nlm.nih.gov/pubmed/29158171>
227. Kunkle, D.A., *et al.* Evaluation and management of gunshot wounds of the penis: 20-year experience at an urban trauma center. *J Trauma*, 2008. 64: 1038.
<https://www.ncbi.nlm.nih.gov/pubmed/18404072>
228. Gong, I.H., *et al.* Comparison of immediate primary repair and delayed urethroplasty in men with bulbous urethral disruption after blunt straddle injury. *Korean J Urol*, 2012. 53: 569.
<https://www.ncbi.nlm.nih.gov/pubmed/22950003>
229. Elgammal, M.A. Straddle injuries to the bulbar urethra: management and outcome in 53 patients. *Int Braz J Urol*, 2009. 35: 450.
<https://www.ncbi.nlm.nih.gov/pubmed/19719861>
230. Maheshwari, P.N., *et al.* Immediate endoscopic management of complete iatrogenic anterior urethral injuries: a case series with long-term results. *BMC Urol*, 2005. 5: 13.
<https://www.ncbi.nlm.nih.gov/pubmed/16281970>
231. Lumen, N., *et al.* Perineal anastomotic urethroplasty for posttraumatic urethral stricture with or without previous urethral manipulations: a review of 61 cases with long-term followup. *J Urol*, 2009. 181: 1196.
<https://www.ncbi.nlm.nih.gov/pubmed/19152939>
232. Scarberry, K., *et al.* Delayed Posterior Urethroplasty Following Pelvic Fracture Urethral Injury: Do We Have to Wait 3 Months? *Urology*, 2018.
<https://www.ncbi.nlm.nih.gov/pubmed/29545047>
233. Aboutaieb, R., *et al.* [Surgical treatment of traumatic ruptures of the posterior urethra]. *Prog Urol*, 2000. 10: 58.
<https://www.ncbi.nlm.nih.gov/pubmed/10785920>
234. Sfaxi, M., *et al.* [Surgical treatment of post-traumatic complete urethral rupture: deferred urgent urethral suture or delayed repair?]. *Prog Urol*, 2006. 16: 464.
<https://www.ncbi.nlm.nih.gov/pubmed/17069041>
235. Leddy, L.S., *et al.* Outcomes of endoscopic realignment of pelvic fracture associated urethral injuries at a level 1 trauma center. *J Urol*, 2012. 188: 174.
<https://www.ncbi.nlm.nih.gov/pubmed/22591965>
236. Elshout, P.J., *et al.* Outcomes of Early Endoscopic Realignment Versus Suprapubic Cystostomy and Delayed Urethroplasty for Pelvic Fracture-related Posterior Urethral Injuries: A Systematic Review. *Eur Urol Focus*, 2017. 3: 545.
<https://www.ncbi.nlm.nih.gov/pubmed/28753868>
237. Warner, J.N., *et al.* The management of the acute setting of pelvic fracture urethral injury (realignment vs. suprapubic cystostomy alone). *Arab J Urol*, 2015. 13: 7.
<https://www.ncbi.nlm.nih.gov/pubmed/26019971>
238. Barrett, K., *et al.* Primary realignment vs suprapubic cystostomy for the management of pelvic fracture-associated urethral injuries: a systematic review and meta-analysis. *Urology*, 2014. 83: 924.
<https://www.ncbi.nlm.nih.gov/pubmed/24680459>
239. Tausch, T.J., *et al.* Unintended negative consequences of primary endoscopic realignment for men with pelvic fracture urethral injuries. *J Urol*, 2014. 192: 1720.
<https://www.ncbi.nlm.nih.gov/pubmed/24972309>
240. Horiguchi, A., *et al.* Primary Realignment for Pelvic Fracture Urethral Injury Is Associated With Prolonged Time to Urethroplasty and Increased Stenosis Complexity. *Urology*, 2017. 108: 184.
<https://www.ncbi.nlm.nih.gov/pubmed/28606774>
241. Koraitim, M.M. Effect of early realignment on length and delayed repair of postpelvic fracture urethral injury. *Urology*, 2012. 79: 912.
<https://www.ncbi.nlm.nih.gov/pubmed/22342415>
242. Mundy, A.R. Anastomotic urethroplasty. *BJU Int*, 2005. 96: 921.
<https://www.ncbi.nlm.nih.gov/pubmed/16153236>

243. Hosseini, J., *et al.* Effects of Anastomotic Posterior Urethroplasty (Simple or Complex) on Erectile Function: a Prospective Study. *Urol J*, 2018. 15: 33.
<https://www.ncbi.nlm.nih.gov/pubmed/29299889>
244. Koraitim, M.M. Predictors of erectile dysfunction post pelvic fracture urethral injuries: a multivariate analysis. *Urology*, 2013. 81: 1081.
<https://www.ncbi.nlm.nih.gov/pubmed/23465164>
245. Brandes, S.B., *et al.* External genitalia gunshot wounds: a ten-year experience with fifty-six cases. *J Trauma*, 1995. 39: 266.
<https://www.ncbi.nlm.nih.gov/pubmed/7674395>
246. Monga, M., *et al.* Testicular Trauma. *Adolesc Med*, 1996. 7: 141.
<https://www.ncbi.nlm.nih.gov/pubmed/10359963>
247. Frauscher, F., *et al.* US findings in the scrotum of extreme mountain bikers. *Radiology*, 2001. 219: 427.
<https://www.ncbi.nlm.nih.gov/pubmed/11323467>
248. de Peretti, F., *et al.* [Fuel tanks of motorcycles. Role in severe trauma of the pelvis]. *Presse Med*, 1993. 22: 61.
<https://www.ncbi.nlm.nih.gov/pubmed/8493205>
249. Herrmann, B., *et al.* Genital injuries in prepubertal girls from inline skating accidents. *Pediatrics*, 2002. 110: e16.
<https://www.ncbi.nlm.nih.gov/pubmed/12165615>
250. Lawson, J.S., *et al.* Catastrophic injuries to the eyes and testicles in footballers. *Med J Aust*, 1995. 163: 242.
<https://www.ncbi.nlm.nih.gov/pubmed/7565208>
251. Gaspar, S.S., *et al.* Sexual Urological Emergencies. *Sexual Medicine Reviews*, 2015. 3: 93.
<https://www.ncbi.nlm.nih.gov/pubmed/27784550>
252. Amer, T., *et al.* Penile Fracture: A Meta-Analysis. *Urol Int*, 2016. 96: 315.
<https://www.ncbi.nlm.nih.gov/pubmed/26953932>
253. Haas, C.A., *et al.* Penile fracture and testicular rupture. *World J Urol*, 1999. 17: 101.
<https://www.ncbi.nlm.nih.gov/pubmed/10367369>
254. Nicolaisen, G.S., *et al.* Rupture of the corpus cavernosum: surgical management. *J Urol*, 1983. 130: 917.
<https://www.ncbi.nlm.nih.gov/pubmed/6632099>
255. Tsang, T., *et al.* Penile fracture with urethral injury. *J Urol*, 1992. 147: 466.
<https://www.ncbi.nlm.nih.gov/pubmed/1732623>
256. De Luca, F., *et al.* Functional outcomes following immediate repair of penile fracture: a tertiary referral centre experience with 76 consecutive patients. *Scand J Urol*, 2017. 51: 170.
<https://www.ncbi.nlm.nih.gov/pubmed/28125311>
257. McGregor, M.J., *et al.* Sexual assault forensic medical examination: is evidence related to successful prosecution? *Ann Emerg Med*, 2002. 39: 639.
<https://www.ncbi.nlm.nih.gov/pubmed/12023707>
258. Donovan, J.F., *et al.* The therapy of genital trauma by dog bite. *J Urol*, 1989. 141: 1163.
<https://www.ncbi.nlm.nih.gov/pubmed/2651716>
259. McAninch, J.W., *et al.* Major traumatic and septic genital injuries. *J Trauma*, 1984. 24: 291.
<https://www.ncbi.nlm.nih.gov/pubmed/6368854>
260. Presutti, R.J. Prevention and treatment of dog bites. *Am Fam Physician*, 2001. 63: 1567.
<https://www.ncbi.nlm.nih.gov/pubmed/11327433>
261. Selikowitz, S.M. Penetrating high-velocity genitourinary injuries. Part I. Statistics mechanisms, and renal wounds. *Urology*, 1977. 9: 371.
<https://www.ncbi.nlm.nih.gov/pubmed/855062>
262. Hudak, S.J., *et al.* Operative management of wartime genitourinary injuries at Balad Air Force Theater Hospital, 2005 to 2008. *J Urol*, 2009. 182: 180.
<https://www.ncbi.nlm.nih.gov/pubmed/19450817>
263. Cass, A.S., *et al.* Bilateral testicular injury from external trauma. *J Urol*, 1988. 140: 1435.
<https://www.ncbi.nlm.nih.gov/pubmed/3193512>
264. Michielsen, D., *et al.* Burns to the genitalia and the perineum. *J Urol*, 1998. 159: 418.
<https://www.ncbi.nlm.nih.gov/pubmed/9649253>
265. Nelius, T., *et al.* Genital piercings: diagnostic and therapeutic implications for urologists. *Urology*, 2011. 78: 998.
<https://www.ncbi.nlm.nih.gov/pubmed/22054364>
266. Lee, J.Y., *et al.* Traumatic dislocation of testes and bladder rupture. *Urology*, 1992. 40: 506.
<https://www.ncbi.nlm.nih.gov/pubmed/1466102>

267. Nagarajan, V.P., *et al.* Traumatic dislocation of testis. *Urology*, 1983. 22: 521.
<https://www.ncbi.nlm.nih.gov/pubmed/6649208>
268. Pollen, J.J., *et al.* Traumatic dislocation of the testes. *J Trauma*, 1982. 22: 247.
<https://www.ncbi.nlm.nih.gov/pubmed/7069812>
269. Shefi, S., *et al.* Traumatic testicular dislocation: a case report and review of published reports. *Urology*, 1999. 54: 744.
<https://www.ncbi.nlm.nih.gov/pubmed/10754145>
270. Cass, A.S., *et al.* Testicular injuries. *Urology*, 1991. 37: 528.
<https://www.ncbi.nlm.nih.gov/pubmed/2038785>
271. Wang, Z., *et al.* Diagnosis and management of testicular rupture after blunt scrotal trauma: a literature review. *Int Urol Nephrol*, 2016. 48: 1967.
<https://www.ncbi.nlm.nih.gov/pubmed/27567912>
272. Wasko, R., *et al.* Traumatic rupture of the testicle. *J Urol*, 1966. 95: 721.
<https://www.ncbi.nlm.nih.gov/pubmed/5935538>
273. Tchounzou, R., *et al.* Retrospective Analysis of Clinical Features, Treatment and Outcome of Coital Injuries of the Female Genital Tract Consecutive to Consensual Sexual Intercourse in the Limbe Regional Hospital. *Sex Med*, 2015. 3: 256.
<https://www.ncbi.nlm.nih.gov/pubmed/26797059>
274. Sotto, L.S., *et al.* Perigenital hematomas; analysis of forty-seven consecutive cases. *Obstet Gynecol*, 1958. 12: 259.
<https://www.ncbi.nlm.nih.gov/pubmed/13578292>
275. Husmann, D.A. Editorial Comment. *J Urol* 1998. 159: 959.
<https://www.jurology.com/doi/full/10.1016/S0022-5347%2801%2963782-0>
276. Okur, H., *et al.* Genitourinary tract injuries in girls. *Br J Urol*, 1996. 78: 446.
<https://www.ncbi.nlm.nih.gov/pubmed/8881959>
277. Goldman, H.B., *et al.* Traumatic injuries of the female external genitalia and their association with urological injuries. *J Urol*, 1998. 159: 956.
<https://www.ncbi.nlm.nih.gov/pubmed/9474191>
278. Karadeniz, T., *et al.* Penile fracture: differential diagnosis, management and outcome. *Br J Urol*, 1996. 77: 279.
<https://www.ncbi.nlm.nih.gov/pubmed/8800899>
279. Fedel, M., *et al.* The value of magnetic resonance imaging in the diagnosis of suspected penile fracture with atypical clinical findings. *J Urol*, 1996. 155: 1924.
<https://www.ncbi.nlm.nih.gov/pubmed/8618289>
280. Pretorius, E.S., *et al.* MR imaging of the penis. *Radiographics*, 2001. 21 Spec No: S283.
<https://www.ncbi.nlm.nih.gov/pubmed/11598264>
281. Uder, M., *et al.* MRI of penile fracture: diagnosis and therapeutic follow-up. *Eur Radiol*, 2002. 12: 113.
<https://www.ncbi.nlm.nih.gov/pubmed/11868085>
282. Buckley, J.C., *et al.* Diagnosis and management of testicular ruptures. *Urol Clin North Am*, 2006. 33: 111.
<https://www.ncbi.nlm.nih.gov/pubmed/16488285>
283. Andipa, E., *et al.* Magnetic resonance imaging and ultrasound evaluation of penile and testicular masses. *World J Urol*, 2004. 22: 382.
<https://www.ncbi.nlm.nih.gov/pubmed/15300391>
284. Corrales, J.G., *et al.* Accuracy of ultrasound diagnosis after blunt testicular trauma. *J Urol*, 1993. 150: 1834.
<https://www.ncbi.nlm.nih.gov/pubmed/8080482>
285. Fournier, G.R., Jr., *et al.* Scrotal ultrasonography and the management of testicular trauma. *Urol Clin North Am*, 1989. 16: 377.
<https://www.ncbi.nlm.nih.gov/pubmed/2652862>
286. Kratzik, C., *et al.* Has ultrasound influenced the therapy concept of blunt scrotal trauma? *J Urol*, 1989. 142: 1243.
<https://www.ncbi.nlm.nih.gov/pubmed/2681835>
287. Martinez-Pineiro, L., Jr., *et al.* Value of testicular ultrasound in the evaluation of blunt scrotal trauma without haematocele. *Br J Urol*, 1992. 69: 286.
<https://www.ncbi.nlm.nih.gov/pubmed/1568102>
288. Micallef, M., *et al.* Ultrasound features of blunt testicular injury. *Injury*, 2001. 32: 23.
<https://www.ncbi.nlm.nih.gov/pubmed/11164397>

289. Mulhall, J.P., *et al.* Emergency management of blunt testicular trauma. *Acad Emerg Med*, 1995. 2: 639.
<https://www.ncbi.nlm.nih.gov/pubmed/8521212>
290. Patil, M.G., *et al.* The value of ultrasound in the evaluation of patients with blunt scrotal trauma. *Injury*, 1994. 25: 177.
<https://www.ncbi.nlm.nih.gov/pubmed/8168890>
291. Churukanti, G.R., *et al.* Role of Ultrasonography for Testicular Injuries in Penetrating Scrotal Trauma. *Urology*, 2016. 95: 208.
<https://www.ncbi.nlm.nih.gov/pubmed/27132505>
292. Lee, S.H., *et al.* Trauma to male genital organs: a 10-year review of 156 patients, including 118 treated by surgery. *BJU Int*, 2008. 101: 211.
<https://www.ncbi.nlm.nih.gov/pubmed/17922859>
293. Muglia, V., *et al.* Magnetic resonance imaging of scrotal diseases: when it makes the difference. *Urology*, 2002. 59: 419.
<https://www.ncbi.nlm.nih.gov/pubmed/11880084>
294. Talan, D.A., *et al.* Bacteriologic analysis of infected dog and cat bites. Emergency Medicine Animal Bite Infection Study Group. *N Engl J Med*, 1999. 340: 85.
<https://www.ncbi.nlm.nih.gov/pubmed/11880084>
295. Presutti, R.J. Bite wounds. Early treatment and prophylaxis against infectious complications. *Postgrad Med*, 1997. 101: 243.
<https://www.ncbi.nlm.nih.gov/pubmed/9126216>
296. Lewis, K.T., *et al.* Management of cat and dog bites. *Am Fam Physician*, 1995. 52: 479.
<https://www.ncbi.nlm.nih.gov/pubmed/7625323>
297. Dreesen, D.W., *et al.* Current recommendations for the prophylaxis and treatment of rabies. *Drugs*, 1998. 56: 801.
<https://www.ncbi.nlm.nih.gov/pubmed/9829154>
298. Anderson, C.R. Animal bites. Guidelines to current management. *Postgrad Med*, 1992. 92: 134.
<https://www.ncbi.nlm.nih.gov/pubmed/1614928>
299. Gee, S., *et al.* On behalf of the North West Policy Group. Guidance for the Management of Human Bite Injuries. 2010.
https://webarchive.nationalarchives.gov.uk/20140714113432/http://www.hpa.org.uk/webc/HPAwebFile/HPAweb_C/1194947350692
300. Summerton, D.J., *et al.* Reconstructive surgery in penile trauma and cancer. *Nat Clin Pract Urol*, 2005. 2: 391.
<https://www.ncbi.nlm.nih.gov/pubmed/16474736>
301. Mydlo, J.H., *et al.* Urethrography and cavernosography imaging in a small series of penile fractures: a comparison with surgical findings. *Urology*, 1998. 51: 616.
<https://www.ncbi.nlm.nih.gov/pubmed/9586616>
302. Penbegul, N., *et al.* No evidence of depression, anxiety, and sexual dysfunction following penile fracture. *Int J Impot Res*, 2012. 24: 26.
<https://www.ncbi.nlm.nih.gov/pubmed/21918532>
303. Virasoro, R., *et al.* Penile Amputation: Cosmetic and Functional Results. *Sexual Medicine Reviews*, 2015. 3: 214.
<https://www.ncbi.nlm.nih.gov/pubmed/27784611>
304. Babaei, A.R., *et al.* Penile replantation, science or myth? A systematic review. *Urol J*, 2007. 4: 62.
<https://www.ncbi.nlm.nih.gov/pubmed/17701923>
305. Tiguert, R., *et al.* Management of shotgun injuries to the pelvis and lower genitourinary system. *Urology*, 2000. 55: 193.
<https://www.ncbi.nlm.nih.gov/pubmed/10688077>
306. Altarac, S. Management of 53 cases of testicular trauma. *Eur Urol*, 1994. 25: 119.
<https://www.ncbi.nlm.nih.gov/pubmed/8137851>
307. Cass, A.S., *et al.* Value of early operation in blunt testicular contusion with hematocele. *J Urol*, 1988. 139: 746.
<https://www.ncbi.nlm.nih.gov/pubmed/3352037>
308. Altarac, S. A case of testicle replantation. *J Urol*, 1993. 150: 1507.
<https://www.ncbi.nlm.nih.gov/pubmed/8411440>
309. Orvis, B.R., *et al.* Penile rupture. *Urol Clin North Am*, 1989. 16: 369.
<https://www.ncbi.nlm.nih.gov/pubmed/2652861>
310. Etabbal, A.M., *et al.* War-related penile injuries in Libya: Single-institution experience. *Arab J Urol*, 2018. 16: 250.
<https://www.ncbi.nlm.nih.gov/pubmed/29892491>

311. Starmer, B.Z., *et al.* Considerations in fertility preservation in cases of testicular trauma. *BJU Int*, 2018. 121: 466.
<https://www.ncbi.nlm.nih.gov/pubmed/29164757>

6. CONFLICT OF INTEREST

All members of the Urological Trauma Guidelines working group have provided disclosure statements of all relationships that they have that might be perceived as a potential source of a conflict of interest. This information is publically accessible through the European Association of Urology website: <http://uroweb.org/guideline>. This guidelines document was developed with the financial support of the European Association of Urology. No external sources of funding and support have been involved. The EAU is a non-profit organisation and funding is limited to administrative assistance and travel and meeting expenses. No honoraria or other reimbursements have been provided.

7. CITATION INFORMATION

The format in which to cite the EAU Guidelines will vary depending on the style guide of the journal in which the citation appears. Accordingly, the number of authors or whether, for instance, to include the publisher, location, or an ISBN number may vary.

The compilation of the complete Guidelines should be referenced as:

EAU Guidelines. Edn. presented at the EAU Annual Barcelona Spain 2019. ISBN 978-94-92671-04-2.

If a publisher and/or location is required, include:

EAU Guidelines Office, Arnhem, The Netherlands. <http://uroweb.org/guidelines/compilations-of-all-guidelines/>

References to individual guidelines should be structured in the following way:

Contributors' names. Title of resource. Publication type. ISBN. Publisher and publisher location, year.